

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (75-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.

59144-6

DATE OF ISSUANCE

MAY 28, 1991

TERM OF ISSUANCE

Until Reregistration

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

Green Ona Liquid Sevin Insect
Spray

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Green Ona, Inc.
P.O. Box 200
Madison, GA 30650

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Under the authority vested in me by section 3(c)(7)(B) of the Act, I have determined that you:

1. are in compliance with the Act and the registration of your product under section 3(c)(7)(B) of the Act is hereby approved when the agency receives all required information and you are in compliance with the Act.

2. make the labeling changes listed below before you release the product for shipment:

a. Add the following to the label: "EPA Reg. No. 59144-6."

b. Change the preharvest interval for grapes to 7 days.

3. Submit five (5) copies of your revised label to the agency before you release the product for shipment. Refer to the 5-79 enclosure for a further description of label requirements.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

Dennis H. Edwards

DATE

5/28/91

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (E7505C)

Enclosures

1 1 2 5

1 1 2 5

GREEN CHARM LIQUID SEVIN[®] INSECT SPRAY

Brand Carbaryl Insecticide
CONTROLS INSECTS ON VEGETABLES, LAWNS,
ORNAMENTALS AND FRUIT

ACTIVE INGREDIENT:

Carbaryl: 1-Naphthyl N-methylcarbamate - - - - 21.3 %

INERT INGREDIENTS: - - - - - 78.7 %

Contains 2 lbs. Carbaryl per gallon Total 100.0 %
SEVIN is a registered trademark of Rhone-Poulenc for Carbaryl insecticides.

KEEP OUT OF REACH OF CHILDREN CAUTION

"PRECAUTIONAL USUARIO Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicada ampliamente"

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED Call a physician of Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES** Flush eyes with plenty of water. Get medical attention immediately. **IF ON SKIN** Wash thoroughly with soap and water. **IF INHALED** Remove victim to fresh air. Apply Artificial Respiration if indicated. **NOTE TO PHYSICIAN** Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Over exposure to this substance may cause toxic signs and symptoms due to the stimulation of the cholinergic nervous system. These effects of over exposure are spontaneously and rapidly reversible. See NOTE TO PHYSICIAN below.
SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION May be harmful if swallowed. Avoid breathing spray mist or vapors. Do not take internally. Avoid contact with skin, eyes or clothing. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling. **NOTE TO PHYSICIAN** Also observe NOTE in Statements of Practical Treatment. Specific treatment consists of periteneal atropine sulfate. Caution should be maintained to prevent over-atropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes to maintain the effect for at least 12 hours. Dosage for children should be appropriately reduced. Complete recovery from over exposure is to be expected within 24 hour. Narcotics and other sedatives should not be used. Further, drugs like 2-PAM are not recommended unless organophosphate intoxication is also suggested.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands, except under forest canopy. Do not apply when weather conditions favor drift from area treated. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until 1 week after application or to take equally effective precautions.
10-01-80

EPA Reg. No. 59144- EPA Est. No. 59144-GA-1

Net Contents 1 Pint



P.O. BOX 290 - MADISON, GA 30650

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas until sprays have dried unless appropriate protective clothing is worn. Protective clothing means at least a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric, covering the body, including the arms and legs) shoes and socks.

Because certain States may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Written or oral warnings must be given to workers who are expected to be in treated areas or in an area to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information: "CAUTION: Area treated with CARBARYL (name) on (Date). Do not enter treated area without appropriate clothing until sprays have dried. See Statement of Practical Treatment and Precautionary Statements for action to be taken in case of accidental exposure."

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original labeled container in a cool, dry, locked place out of reach of children. Do not store in areas where temperatures frequently exceed 100° F. The product may be used following exposure to several freeze/thaw cycles. **DISPOSAL:** Do not reuse empty container. Wrap empty container in several layers of newspaper and discard in trash.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system. Shake containers prior to use. Do not reuse empty containers or measuring devices for other purposes. Apply with a pressure, power, pump or hose-end sprayer to give thorough coverage when insects or damage appear. Repeat as necessary to maintain control, unless spray interval is specified. **Plant Response Precautions:** Carbaryl will injure Boston Ivy, Virginia Creeper and Maidenhair fern. During early season it may injure Virginia and Sand Pine species.

VEGETABLE CROPS

All spray rates are underlined as; - (x) tps./gal. Each rate follows the specific insect names as listed. Do not over dose.

ASPARAGUS: Asparagus Beetle, Grasshoppers - 4 to 8 tps./gal. Pre-harvest interval - 1 day. Apache Cicada, Asparagus Beetle - 8 to 16 tps./gal. Post harvest application only. Treat ferns or brush growth. Do not treat more than once every 3 days.

BEANS: BLACK EYED PEAS, COWPEAS, CROWDER OR SOUTHERN PEAS, DRY BEANS, GREEN BEANS, LIME BEANS, NAVY BEANS, AND SNAP BEANS: Blister Beetle, Mexican Bean Beetles, - 2 to 4 tps./gal.; Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetle, Flea Beetles, Green Cloverworm, Japanese Beetle, Leafhoppers, Three Cornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar, Western Bean Cutworms - 4 tps./gal. Armyworms, Cutworms, European Corn Borer, Fall Armyworms, Grasshoppers, Tarnished Plant Bug, Webworms - 4 to 6 tps./gal. Corn Earworm, Cowpea Curculio, Lima Bean Pod Borer, Lygus Bugs, Stink Bugs - 6 tps./gal. Pre-harvest intervals - 0 days (all except cowpeas), 3 days for cowpeas.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CARROTS, CAULIFLOWER, CELERY, CHINESE CABBAGE, COLLARDS, DANDELION, ENDIVE (ESCAROLE), GARDEN BEET, HANOVER SALAD, HORSE RADISH, KALE, KOHLRABI, LETTUCE, MUSTARD GREENS, OKRA, PARSNIPS, PARSLEY, RADISHES, RUTABAGAS, SALSIFY, SPINACH, SWISS CHARD, TURNIPS Flea Beetles, Harlequin Bugs, Leafhoppers - 2 to 4 tps./gal. Aster Leafhopper, Grasshoppers, Lygus Bugs, Spittlebugs - 4 to 6 tps./gal. Armyworm, Corn Earworm, Fall Armyworm, Imported Cabbage worm, Stink Bug, Tarnished Plant Bug - 4 to 8 tps./gal. Pre-harvest interval days: 0 - Carrots; 2 - Broccoli, brussels sprouts, Cabbage, Cauliflower, Garden Beet Roots, Head Lettuce, Horseradish, Kohlrabi, Parsnips, Radishes, Rutabagas, Salsify Roots, and Turnip Roots; 14 days - Chinese Cabbage, Celery, Collards, Dandelion, Endive (Escarole).

See the back of label for statement

role), Garden Beet tops, Hanover Salad, Kale, Mustard Greens, Leaf Lettuce, Parsley, Salsify tops, Spinach, Swiss Chard, and Turnip tops, NOT FOR USE ON CELERY IN CALIFORNIA. Observe plant response precautions. Lettuce: treat on a 5 to 7 days schedule after heads begin to form.

CORN: Armyworm, Chinch Bug, Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Flea Beetles, Grasshoppers, Japanese Beetle, Leafhoppers, Sap Beetle, Southern Corn Borer - 4 to 8 tps./gal. OBSERVE BEE CAUTION. For insects attacking silks and ears, apply at 1 to 6 days intervals starting when first silks appear and continuing until silks begin to dry. For larvae in whorl and foliage feeders, apply as necessary. Optimum timing and good coverage are essential for effective control. Western Bean Cutworms - 8 tps./gal. Treat when infestation averages 15% and at 90 to 100% tassel emergence. Treatment after 100% silk emergence will reduce effectiveness. Pre-harvest interval - 0 days.

CUCUMBER, MELONS, PUMPKINS, SQUASH: Pickleworm, Melonworms - 2 to 4 tps./gal.; Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs - 4 tps./gal. Pre-harvest intervals - 0 days.

POTATO, TOMATO, EGGPLANT, PEPPER, PEANUTS, PEAS: Bean Leaf Beetles, Blister Beetles, Colorado Potato Beetle, Cucumber Beetles, Flea Beetles, Green Cloverworm, Japanese Beetles, Leafhoppers, Mexican Bean Beetles, Red-necked Peanutoworm, Three Cornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar - 4 tps./gal.; Alfalfa Caterpillar, Armyworm, Corn Earworms, Grasshoppers, Pea Leaf Weevil, Pea Weevil, Webworms, Yellowstriped Armyworm - 6 tps./gal. European Corn Borer, Fall Armyworm, Lace Bug, Stink Bugs, Tarnished Plant Bug, Tomato Hornworm, Tomato Pinworm - 4 to 8 tps./gal. Cutworms, White fringed Beetle adults - 6 to 8 tps./gal. Pre-harvest intervals - 0 days.

SWEET POTATO: Corn Earworm, Cucumber Beetles, Flea Beetles, Grasshoppers, Sweet Potato Hornworm, Tortoise Beetles - 4 to 8 tps./gal. NOT FOR USE IN CALIFORNIA. Apply as a foliar spray as needed. Yellowstriped Armyworm - 8 tps./gal. Sweet Potato Weevil - 4 to 8 tps./gal. Full coverage is essential. Use lower rate on young plants and higher rate on mature plants. Pre-harvest interval - 0 days.

FRUIT AND NUT CROPS

APPLE THINNING: Apply 1 full coverage dilute spray between 10 and 25 days after full bloom. If factors such as tree age, variety, nutrition, previous crop, pruning, bloom and degree of set, favor excessive fruit thinning with carbaryl, exercise caution to avoid possible yield reduction. Pre-harvest interval - 1 day. **APPLES ONLY** Apply 1 to 2 tps./gal. for easily thinned varieties including Cortland, Grimes, Jonathan, McIntosh, Orleans, Rome Beauty, Puritan, Red Delicious, Winesap, Yellow Newton. Apply 2 to 4 tps./gal. for difficult to thin varieties including Baldwin, Northern Spy, Rhode Island Greening, Steve Red, Turley, Wealthy, Yellow Transparent and York Imperial.

ALMOND: Peach Twig Borer, San Jose Scale, Fruitree Leafroller - 4 tps./gal. Apply in popcorn or petal fall stages and again when the May brood of the peach Twig Borer begins to hatch or thereafter as needed. Navel Orangeworm - 4 tps./gal. Time early and mid season applications to correspond to moth flight peaks. Make a late season application at initiation of hull split or up to 10% hull split.

APPLE, PEARS: Apple Aphid, Apple Maggot, Apple Mealybug, Apple Rust Mite, Apple Sucker, Bagworms, California Pear Slug, (Pear Sawfly), Codling Moth, Eastern Tent Caterpillar, European Pear Sawfly, Eyespotted Bud Moth, Forbes Scale,

Fruitree Leafroller, Gracianum Scale, Lesser Apple Scale, Pear Leaf Blister Beetle, Pear Leafminer, Pear Leaf Cicada, Red Banded Scale, Tarnished Plant Leafhoppers, Woolly Apple Aps. /gal. Pre-harvest interval - 0 days. Thinning, delay use until psylla control, apply when present. For optimum present.

BLACKBERRIES, RASPBERRIES, LOGANBERRIES, GRACIANUM SCALE, LESSER APPLE SCALE, PEAR LEAF BLISTER BEETLE, PEAR LEAF MINER, PEAR LEAF CICA, RED BANDED SCALE, TARNISHED PLANT LEAFHOPPERS, WOOLLY APPLE APT. /gal. Pre-harvest interval - 0 days. Thinning, delay use until psylla control, apply when present. For optimum present.
BLACKBERRIES, RASPBERRIES, LOGANBERRIES, GRACIANUM SCALE, LESSER APPLE SCALE, PEAR LEAF BLISTER BEETLE, PEAR LEAF MINER, PEAR LEAF CICA, RED BANDED SCALE, TARNISHED PLANT LEAFHOPPERS, WOOLLY APPLE APT. /gal. Pre-harvest interval - 0 days. Thinning, delay use until psylla control, apply when present. For optimum present.

BLUEBERRIES: Blueberry Fruitworm, European Pear Sawfly. Apply 3 weeks before harvest. Pre-harvest interval - 0 days.

CITRUS FRUITS: (Grapefruit, Tangerines, Citrus Leafroller, California Orange Weevil, Fruitree Leafroller, Moth, Western Indian Black Scale, Brown Scale, Citrus Snow Scale) dilute spray mixture petroleum oils commonly used.

CRANBERRIES: Cutworm, Fruitworm, Japanese Beetle. 12 tps./gal. Pre-harvest interval - 12 days. Shrimp and crabs. Do not use. Apply in late intervals.

FILBERTS: Filbert Apple Sawfly. Pre-harvest interval - 12 days. Eggs are hatching. Reapply when Moth and again 3 to 4 weeks later.

FRUIT AND NUT CROPS

ALSO, TREE LIME DISMORPHIC PEST CONTROL

GREEN CHARM LIQUID SEVIN INSECT SPRAY

Brand Carbaryl Insecticide
CONTROLS INSECTS ON VEGETABLES, LAWNS,
ORNAMENTALS AND FRUIT

ACTIVE INGREDIENT:

Carbaryl: 1-Naphthyl N-methylcarbamate - - - - 21.3%

INERT INGREDIENTS: - - - - - 78.7%

Total 100.0%

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

EPA Reg. No. 59144- EPA Est. No. 59144-GA-1

Net Contents 1 Pint

GRO TEC, INC.

P.O. BOX 290 - MADISON, GA 30650

FRUIT AND NUT CROPS (Cont.)

APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES: Apple Pandemis, Black Cherry Aphid, Brown Soft Scale, Cherry Fruitworm, Cherry Maggot, Cucumber Beetles, European Earwig, Eyespotted Bud Moth, Forbes Scale, Fruittree Leafroller, Green Fruitworm, Japanese Beetles, June Beetles, Lecanium Scales, Lesser Peachtree Borer, Mealy Plum Aphids, Olive Scale, Oriental Fruit Moth, Oystershell Scale, Peach Twig Borer, Periodical Cicada, Plum Curculio, Prune Leafhopper, Redbanded Leafroller, Rose Chafer, San Jose Scale, Tarnished Plant Bug, Variegated Leafroller - 4 tps./gal. Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Tussocks Moth - 3 to 4 tps./gal. For optimum scale control, apply when crawlers are present. For lesser peach tree borer control spray limbs and tree trunks thoroughly, weekly, during moth flight. Pre-harvest interval - 1 day, cherries, peaches, plums, and prunes; 3 days apricots and nectarines.

PECANS: Black Margined Aphid, Fall Webworms, Hickory Shuckworm, Lesser Webworm, Pecan Leaf Phylloxera, Pecan Nut Casebearer, Pecan Spittlebugs, Pecan Weevil, Twig Girdler, Walnut Caterpillar - 4 to 10 tps./gal. Pre-harvest interval - 0 days.

WALNUT: Calico Scale, Codling Moth, European Fruit Lecanium, Filbertworm, Fruittree Leafroller, Frosted Scale - 2 tps./gal. For Codling Moth apply when average cross sectional diameters of nuts are 1/2 to 3/4 inch. Repeat during middle of later June as needed. European Earwig - 8 tps./gal. Spray tree trunks to point of run-off. Pre-harvest interval - 0 days.

TREES AND ORNAMENTALS

For dilute spray applications to trees, (including shade trees, shelter belts, plantations, parks and recreational areas), ornamental, woody plants and shrubs, apply the specified dosage per gallon of water in sufficient spray volume to provide thorough coverage. (6 teaspoonfuls equal 1 fluid ounce.)

FOR TREES, WOODY PLANTS & SHRUBS apply with: ressur, power, pump, or hose-end sprayer. Do not treat Boston Ivy, Virginia Creeper, Maiden Hair fern; nor Virginia or Sand Pine early in season.

INSECTS CONTROLLED ON TREES AND ORNAMENTALS

Ants, Apple Aphid, Armyworms, Azalea Leafminer, Bagworms, Birch leafminer, Blister Beetles, Boxelder Bugs, Boxwood Leafminer, Browntail Moth, Cankerworms, Catalpa Sphinx, Chiggers, Cooley Spruce Gall Aphid, Cutworms, Cypress Tip Moth, Douglas Fir Tussock Moth, Eastern Spruce Gall Aphid, Elm Leaf Aphid, Elm Leaf Beetles, Elm Span Worm, Eriophyid Mites, European Pine Shoot Moth, Fall Armyworm, Flea Beetles, Fuller Rose Beetle, Gall Midges, Gall Wasps, Green Striped Mapleworm, Grasshoppers, Gypay Moth, Hackberry Nipplegall, Inaker, Holly Bud Moth, Holly Leafminers, Jackpine Budworm, Japanese Beetles, Jeffrey Pine Needleminer, June Beetles, Lace Bugs, Leafhoppers, Leafrollers, Locusts Borer, Maple Leafcutter, Mealy Bugs, Mimosa Webworms, Nantucket Pine Tip Moth, Oak Leafminers, Oak Leaf Skeletonizer, Oakworm complex, Oleander Caterpillar, Olive Ash Borer, Orange Striped Oakworm, Orange Tortrix, Periodical Cicada, Pine Sawfly, Pine Spittlebug, Pitch Pine Tip Moth, Plant Bugs, Poinsettia Hornworm, Psyllids, Puss Caterpillar, Redhumped Oakworm, Rose Aphid, Rose Chafer, Rose Slug, Saddled Prominent, Sawflies (exposed), Scale Insects, Sowbugs, Spiny Elm Caterpillar, Springtails, Spruce Budworm, Spruce Needleminer, Subtropical Pine Tip Moth, Tent Caterpillar, Thorn Bugs, Thrips (exposed), Ticks, Walnut Caterpillar, Webworms, Western Hemlock Looper, Western Spruce Budworm, Willow Leaf Beetles, Yellow Poplar Weevil - Use 4 tps./gal.

Observe Plant response precautions. Apply dilute spray to obtain thorough coverage of upper and lower leaf surfaces. To control scale insects, treat trunks, stems and twigs in addition to plant foliage. For optimum control treat when worms are small. Addition of a sticker may improve residual control.

For Ips Engraver Beetles, Mountain Pine Beetle, Roundheaded Pine Beetle, Western Pine Beetle - use 13.3 fluid ounces per gallon of water. Effective as a preventive treatment only. Repeat annually as required to prevent beetle attacks. Apply 1 gallon of spray per 50 square feet of bark in late May to early June, or prior to beetle attack. Treat tree trunk from ground level up until trunk diameter is less than 5 inches.

LAWN AND RECREATIONAL AREAS TURF GRASSES

INSECTS: Ants, Armyworms, Bluegrass Billbugs, Centipedes, Chiggers, Chinch Bugs, Cutworms, Earwigs, Essex Skipper, European Chafer, Fall Armyworm, Fiery Skipper, Fleas, Grasshoppers, June Beetles, Leafhoppers, Lucerne Moth, Millipedes, Mosquitoes, Sod Webworms, Sow bugs, Springtails, Ticks, White Grubs, Yellowstriped Armyworm - use 12 fl. oz. per 1000 sq. ft. (16 qts per acre) of turf grass. Make application in sufficient spray volume for thorough coverage and turf hatch penetration. Repeat as necessary.

For Armyworms, Cutworms, Fall Armyworms, and Sod Webworms control do not irrigate treated areas following insecticide application.

For Chinch Bugs control treat entire turf grass area rather than just damaged areas. Irrigation of turf grass area before insecticide application will aid in penetration into turf grass.

For White Grub control applications should be made when grubs are feeding near the soil surface, usually during late March through May, or July to early September, or as recommended by local Agricultural Extension Service agents. Water or irrigate turf grasses thoroughly soon after treatment.

For IMPORTED FIRE ANTS - use 2 tablespoons (1 fluid ounce) per gallon of water. Apply a total of 2 gallons of the diluted solution over the surface of each mound, or at least 1 quart per 6 inches of mound diameter, using bucket, can or other appropriate equipment. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12 sq.ft.). Do not disturb mounds prior to treatment. Pour solution from a height of about 3 feet to give sufficient force to break mound apex and flow into ant tunnels. For best results apply in cool weather, 65 to 80 degrees F, or in early morning or late evening hours. Repeat application if mound activity resumes after 10 days. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.

CONTROL OF TICKS THAT VECTOR LYME DISEASE

May be used for control of juvenile and adult Ixodes spp. (Deer Tick and Black Legged Tick) and Amblyomma spp. ticks (Lone Star Ticks) which vector Lyme Disease. Mix 2 tps./gal. and begin spray applications to lawns, shrub beds, ornamental plantings, wooded areas and around outside perimeters of home and other buildings in late spring or early summer (May/June) to control juvenile ticks (larvae and nymphs). Do not make spot applications. Treat entire lawn and perimeter wooded areas and property boundaries where exposure to ticks may occur. Control of adult ticks can be achieved with late summer and fall applications (Aug/Sept) at the same rate. Ticks may be reintroduced from surrounding areas on host animals. Repeat treatment at 3 to 4 week intervals to maintain adequate control levels.

MOSQUITO CONTROL

YARDS and RECREATIONAL AREAS: Adult Mosquitoes - Precautions - May kill shrimp and crabs. Do not use in areas where these are important resource. See Bee Precautionary Statement - Avoid direct application to lakes, streams and ponds.

For dilute spray ground applications to trees (including shade trees, shelter belts, plantations, parks and recreational areas), ornamental, woody plants and shrubs - apply 1 to 2 tps./gal. Treat shrubbery and areas where adult mosquitoes congregate. Treat when adult mosquitoes are active in early morning or late evening. Repeat at 7 to 10 day intervals. For residual control in subtropical regions use 16 fl. oz. per gallon of water and apply 4 gallons of prepared spray per 2000 sq. ft. of surface area. Repeat in 3 to 6 months or when necessary.

PEST CONTROL IN AND AROUND BUILDINGS

INSECTS: Brown Dog Ticks, Earwigs, Fleas, Millipedes, Silverfish - 7 ozs./gal. Ants, Cockroaches - 17 ozs./gal. For use in and around buildings such as homes, apartments, warehouses, barns and municipal and recreational areas. Apply as coarse wet spray or with a paint brush to outdoor sleeping quarters of pets, outside perimeter of dwellings and other areas where insects tend to congregate.

Do not use this product in commercial food areas of food handling establishments, restaurants, or the places where food is prepared and processed. Avoid application to surfaces where visible spray residues are objectionable.

POULTRY I

CHICKENS, DUCKS, GESE: Chicken Mites, Fleas as a direct spray on bird Machine: Mix 20 ounces c 1/2 gallons per 1000 hens Repeat in 4 weeks if necessary or cylinder type compress gallons of spray. Use 1 gal on slatted floor. Repeat in Direct "mist spraying" for c to spraying insects and b not apply to poultry and g

POULTRY ROX

INSECTS: Bed Bugs, Chick Mite - 5 tps./gal. Fowl Tick of spray mixture per 100 sq feet spray into cracks. Repeat nests, eggs and feeding spraying. Do not treat pre Lesser Mealworm - 33 ozs. per 100 sq. ft. of floor space Ventilate while spraying. C slaughter.

NOTICE: This product is reasonably fit for the use in accordance with direct Manufacturer is not responsible for conditions not reseller and/or seller and buy