

NOV 25 1992

Mr. H. R. McLane  
H. R. McLane, Inc.  
7210 Red Road, Suite 206  
Miami, Florida 33143-5321

Dear Mr. McLane:

Subject: Product Name Change & New Labeling  
Crawling Insect Granules 100  
EPA Registration No. 59144-9  
Your Application Dated September 2, 1992

We have no objection to the change in the name of this product from "Crawling Insect Granules 100" to "Green Charm Dursban 1½ Granules." Our files will be changed to recognize this as the official name of record for this product.

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely,

DHE

Dennis H. Edwards, Jr.  
Product Manager 19  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosure

# GREEN CHARM®


# DURSBAN®

## 1% GRANULES

*For Control of Fleas, Ticks, Chiggers and  
other Crawling Insects as Listed in Lawns,  
Turf and around the Outside of Homes  
Kill's Fire Ants*

1 POUND TREATS UP TO 400 SQ.FT

INSECT CONTROL OUTSIDE OF HOMES,  
APARTMENTS, CONDOMINIUMS, WAREHOUSES,  
HOSPITALS, SCHOOLS, FOOD HANDLING  
ESTABLISHMENTS & COMMERCIAL BUILDINGS



**ACCEPTED**

NOV 25 1992

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 59144-9

ACTIVE INGREDIENT:  
Chlorpyrifos: [O,O,-diethyl O-(3,5,6-trichloro  
-2-pyridyl) phosphorothioate] - - 1.0%

INERT INGREDIENTS: - - - - - 99.0%

Total 100.0%

DURSBAN® is a Reg. Trademark of DowElanco  
KEEP OUT OF REACH OF CHILDREN  
CAUTION

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT

EPA Reg No 59144-9 EPA EST. NO 5-144-GA-1<sub>a</sub>; 6720-FL-1<sub>a</sub>; 51038 GA-1<sub>c</sub>; 788-GA-1<sub>a</sub>; 44816-MO-1<sub>a</sub>; 19713-GA-1,  
Subscript is first letter of run code on container (59-9BAG PM4/ 03-01-92)

NET WEIGHT 10 POUNDS



P.O. BOX 290 - MADISON, GA 30650

# GREEN CHARM®

## DURSBAN®

### 1% GRANULES

NET WEIGHT 10 POUNDS

#### IMPORTED FIRE ANT CONTROL

**MOUND TREATMENT:** For treatment of Fire Ant mounds occurring in lawns, turf and premise areas, treat individual ant mounds measuring from 3 to 12 inches in diameter by applying 4 tablespoons (8 oz.) of granules evenly in a circle around the outer edges (1 to 2 feet from the center) of the mound. Finish the treatment then by applying 7 tablespoons (1.2 oz.) sprinkled evenly over the center of the mound. After application of the granules, gently sprinkle the treated area with a minimum of 1 quart of water per 6 inches of mound diameter. Apply in cool weather (65 to 80 degrees F) or in early morning or late evening. Best results will be obtained if mound is not disturbed after treatment thus allowing ants to come and go and to be exposed to the insecticide.

#### FOR LAWNS, GOLF COURSES, SOD FARMS AND OTHER ORNAMENTAL / RECREATIONAL TURF

Apply DURSBAN 1% GRANULES uniformly to home lawns, golf courses, sod farms, and other ornamental and recreational turf and non-crop land areas using any hand or power operated gravity or rotary type spreader designed for application of granules. Treat when insects first appear, using GRANULES at the rate of 100 lbs./acre for Surface Insects. Increase this rate as indicated for control of certain Soil Insects as outlined in separate paragraphs. Immediately after application, sprinkle the turf with 30 gallons of water per 1,000 sq. ft. (1300 gals./acre) to move the granules into the zone of insect infestation. The following table gives amounts of GRANULES and water required for small areas. Repeat treatment as necessary.

\*Observe Watering limitation in paragraph for SOD WEBWORMS.

#### FOR SURFACE INSECTS

For control of Ants, Armyworms (Beet, Fall & Yellowstriped), Brown Dog Ticks, Centipedes, Chiggers, Chinch Bugs, Cutworms, Crickets, Earwigs, Flary Skipper, Fleas, Grasshoppers, Leafhoppers, Millipedes, Clover Mites, Sod Webworms (Lawn Moth), Sowbugs, Springtails, and Ticks use 2.5 lbs. per 1,000 sq. ft.

Areas to be Treated	Pounds of GRANULES	Gallons of Water to Sprinkle on Area after Treatment
400 sq. ft.	1 lb.	5 gals.
2,000 sq. ft.	5 lbs.	30 gals.
10,000 sq. ft.	25 lbs.	150 gals.
20,000 sq. ft.	50 lbs.	300 gals.

#### FOR SOIL FEEDING INSECTS

Refer to specific paragraph for TURFGRASS WEEVIL; WHITE GRUBS; CRANE FLY (larvae) and BLUEGRASS BILLBUG in left panel for rates to control soil insects which feed below the soil surface.

Observe proper Spreader Settings for these higher rates.

#### PERIMETER TREATMENT AROUND DWELLINGS

USE FOR CRAWLING INSECT CONTROL OUTSIDE OF HOMES, APARTMENTS, CONDOMINIUMS, WAREHOUSES, HOSPITALS, SCHOOLS, FOOD HANDLING ESTABLISHMENTS AND COMMERCIAL BUILDINGS

To protect against insects entering the indoors, a band of granules may be placed around the outside of buildings to kill Ants, Brown Dog Ticks, Crickets, Earwigs, Fleas, Millipedes, Sowbugs, Springtails and Spiders before they attempt to enter. Use 1 pound to treat 400 sq. ft. (as a band 5 ft. x 80 ft. long) around the structure. Apply when insects have become troublesome or before they cause problems in the spring or early fall. Repeat one month later and then as needed for control but not more than once every 7 days. Keep children and pets off treated area until granules are sprinkled in and the area is dry.

#### SPREADER SETTINGS

SPREADER SETTINGS are approximate. Your spreader may need recalibrating due to age and use. Test proper working condition by placing 2 1/2 lbs. of granules into your spreader using the setting of the 5 lbs./1,000 sq. ft. rate. Apply over a pre-measured area of 500 sq. ft. (5 ft. x 100 ft. long). If the spreader contains GRANULES at end of test, open the spreader 1/2 to 3/4 setting; if granules have run-out before the end of test, close (shut) the spreader 1/2 to 3/4 setting. Repeat test if indicated insects may not be killed and control may be "spotty" if your spreader does not apply the exact recommended amount/1,000 sq. ft. as recommended!

SPREADER	Pounds of GRANULES to Use per 1,000 Sq. Ft.		
	2.25 - 2.50 lbs.	3.5 lbs.	4.5 - 4.75 lbs.
Scott Lawn	#4	#5	#6
Cyclone 81	#3.5	#4.5	#5.5

**WARRANTY AND DISCLAIMER:** Subject to the limitation of liability herein set forth, Gro-Tec, Inc. warrants that the product sold is as described on the container, with recognized tolerances. Liability on this warranty is limited to the purchase price of the product. In no event shall Gro-Tec, Inc. be liable for indirect, consequential, or other damages in excess of the purchase price. Gro-Tec, Inc. makes no other warranty, either expressed or implied as to merchantability or fitness for a particular purpose of the product. The product is sold at the price quoted only upon the condition that the above disclaimer of warranty and the limitation of damages are part of the basis of the bargain. Products not accepted under these terms should be returned within 20 days in the original, unopened container and Buyer will receive his choice of either return of purchase price or a price quotation which includes such warranties as desired on the basis of the bargain (provided such quote is obtainable which is dependent upon the nature of the products sold) (50-98AG2 PM4/4 of 03/01/83)

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. DO NOT induce vomiting or give anything by mouth to an unconscious or convulsing person. Get medical attention. **IF IN EYES:** Flush eyes with plenty of water. **IF ON SKIN:** In case of contact, remove contaminated clothing and wash affected area with soap and water. **NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote only by injection.

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** May be harmful if swallowed. Avoid contact with eyes and clothing. Avoid breathing dust. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before reuse. Keep away from food, feedstuffs, and water supplies. Keep children and pets off treated areas until this material is washed into the soil and the grass is dry.

##### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Use only as directed.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original labeled container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not re-use empty container. Wrap container and put in trash.

Controls Ants, Armyworms (Beet, Fall, Yellowstriped), Brown Dog Ticks, Centipedes, Chiggers\*, Chinch Bugs\*, Clover Mites, Cutworms, Crickets, Earwigs, Flary Skipper, Fleas\*, Grasshoppers, Greenbugs, Leafhoppers, Millipedes, Sod Webworms\* (Lawn Moth), Sow Bugs, Springtails, Turfgrass Weevil\* (Hyperodes), European Crane Fly larvae\*, White Grubs\* (Black Turfgrass Ateanius, European Chafer, Southern Chafer, Japanese Beetle, June Beetle larvae), Bluegrass Billbugs\*, Fire Ants\*.

#### \*SPECIFIC DIRECTIONS

The following paragraphs correspond to insects followed by asterisk(\*) on the front panel (above). Insects not followed by an Asterisk(\*) may be controlled by following the DIRECTIONS FOR USE ON LAWNS ....(etc.). Repeat treatments as needed for control.

#### CHIGGERS, TICKS, & FLEAS

Use 1 lb / 400 sq. ft. (2.25 to 2.50 lbs./1,000 sq. ft.) for area control of these insects, infesting non-crop land areas such as roadsides, footpaths, and trails, picnic and camping sites, parks, recreational areas and around dwellings where these pests are present and create a nuisance or may act as a disease vector. DO NOT allow use of treated area during applications. For Fleas (out-of-doors) also treat sleeping quarters and other areas frequented by pets such as beneath porches and along fencing.

#### CHINCH BUGS

Apply 12 to 17 lbs./5,000 sq. ft. when insects appear or their damage is imminent. To prevent damage apply as soon as daily temperatures reach 80 degrees and repeat every 2 months. Use low rate for minimum infestations; higher rate for heavy infestations.

#### SOD WEBWORMS

For Sod Webworms, delay watering or mowing the treated area for 12 to 24 hours after treatment. Treat in the spring or when insects first appear troublesome. Additional treatments at 30 day intervals may be needed for new infestations. Apply at rate of 2.5 lbs. per 1,000 sq. ft.

#### TURFGRASS WEEVIL

Use 12.5 lbs. per 2,500 sq. ft. Treat suspected problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service.

#### WHITE GRUBS; CRANE FLY LARVAE

To control White Grubs (Black Turfgrass Ateanius, European Chafer, Southern Masked Chafer, Japanese Beetle and June Beetle Larvae), use 12.5 lbs to treat 2,500 sq. ft. After applications apply enough water to thoroughly wash the insecticide into the turf. For best results remove thatch first and treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service Specialist for best time to treat. European Crane Fly larvae will be controlled using 12.5 lbs per 3,600 sq. ft. or will be controlled simultaneously with the same application being applied for control of "White Grubs" (12.5 lbs./2,500 sq. ft.). Consult your Agricultural Agent for proper time to apply. For best results, first remove thatch. Water the granules into the soil.

#### BLUEGRASS BILLBUGS

Use 12.5 lbs./2,500 sq. ft. for Billbug grubs. Remove thatch before treatment. Wash granules into soil.

**GRO-TEC**

**BEST AVAILABLE COPY**