

59144-12

09-29-2010

1/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Kim Davis
Reg West Company, LLC
8202 West 20th St.
Suite A
Greeley, CO 80634-4696

SEP 29 2010

Subject: EPA Reg. 59144-12 / Pennington Crabgrass 43 with Fertilizer Preemergence Herbicide Notification

Dear Ms. Davis:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 7-27-10 for the product EPA Reg. 59144-12 / Pennington Crabgrass 43 with Fertilizer Preemergence Herbicide. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov or Kathryn Montague at 703-305-1243 or Montague.Kathryn@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn Montague
Team Leader
Herbicide Branch, Team 23
Registration Division (7504P)

| | | |
|--|--|--|
| United States Environmental Protection Agency 1200 Pennsylvania Avenue, N.W. Washington, DC 20460-0001 | <input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other | OPP Identifier Number Notification |
|--|--|--|

Application for Pesticide – Section I

| | | |
|---|---|--|
| 1. Company/Product Number EPA Reg. No. 59144-12 | 2. EPA Product Manager Richard Gebken - K. Montague | 3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/ Product (Name) Pennington Crabgrass Preventer 43 with Fertilizer Preemergence Herbicide | PM # 10 23 | |
| 5. Name and Address of Applicant (Include ZIP Code) Gro Tec, Inc. c/o RegWest Company, LLC 8203 West 20th St., Suite A Greeley, CO 80634-4696 | | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i) my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____ |

Check if this is a new address

Section – II

| | |
|--|---|
| <input type="checkbox"/> Amendment – Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification – Explain below. | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other – Explain below. |
|--|---|

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)

Submit Label Revision Notification per PR Notices 98-10 and 2007-4.

This notification is consistent with the provisions of PR Notices 98-10 and 2007-4 and EPA regulations at 40 CFR § 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notices 98-10 and 2007-4 and 40 CFR § 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section – III

| | | | |
|---|---|---|---|
| 1. Material This Product Will Be Packaged In: | | | |
| Child Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No | Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No | Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No | 2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify): _____ |
| *Certification must be submitted If "Yes" Unit Packaging Wt. No. Per Container | | If "Yes" Packaging Wt. No. Per Container | |
| 3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container | | 4. Size(s) Retail Container | |
| | | 5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling Accompanying Product | |
| 6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Stenciled <input type="checkbox"/> Paper glued <input type="checkbox"/> Other _____ | | | |

Section - IV

| | | |
|--|---|--|
| 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application). | | |
| Name Kim Davis | Title Consultant/Agent | Telephone No. (Include Area Code) 970.353.0611 kim@regwest.com |
| Certification | | 6. Date Application Received _____ (Stamped) |
| I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. | | |
| 2. Signature | 3. Title Consultant/Agent | |
| 4. Typed Name Kim Davis | 5. Date July 27, 2010 | |

Label Revision Notification



July 27, 2010

Mr. Richard Gebken, Product Manager 10
Document Processing Desk - **NOTIF**
Office of Pesticide Programs – **7504P**
U.S. Environmental Protection Agency
One Potomac Yard (South Building), Room S-4900
2777 South Crystal Drive
Arlington, VA 22202

Dear Mr. Gebken:

Subject: Gro Tec, Inc.
Pennington Crabgrass Preventer 43 with Fertilizer Preemergence Herbicide
EPA Reg. No. 59144-12

On behalf of **Gro Tec, Inc.**, RegWest Company, LLC is submitting the attached documents per PR Notices 98-10 and 2007-4. RegWest Company will act as sole agent in this endeavor.

In addition to the 2007-4 PR Notice changes, we are deleting the "Crabgrass Seed Germination Dates" and replacing the text with a graphic. Please refer to the *Reference Label* for an illustration of all revisions.

Please contact me at 970.353.0611 or kim@regwest.com if you have any questions or comments concerning this submission.

Regards,
RegWest Company, LLC

Kim Davis,
Consultant/Agent

Attachments

cc: GTI – Pennington, D.

8202 West 20th St., Suite A
Greely, CO 80634-4696
p 970-353-0611
f 970-353-0613

Pennington Crabgrass Preventer 43 with Fertilizer Preemergence Herbicide

For Selective Preemergent Control of Grass and Broadleaf Weeds in:
Established Turf Grasses (excluding Golf Course Putting Greens) and Lawns
Landscape Ornamentals (including Established Perennials and Wildflower Plantings)

20 Pounds Treats up to 5,714 sq. ft. [an area approximately 57' x 100']
[40 Pounds Treats up to 11,428 sq. ft. [an area approximately 114' x 100']]
[50 Pounds Treats up to 14,285 sq. ft. [an area approximately 142' x 100']]

NOTIFICATION
SEP 29 2010

Contains 0.43% Barricade® Preemergent Herbicide
Feeds up to 4 Months
Crabgrass Preventer Plus Fertilizer with Barricade® 0.43%
Controls Crabgrass
Quality Fertilizers from The Lawn Seed Experts
Crabgrass Preventer + Lawn Fertilizer 32-4-6
Controls Grass and Broadleaf Weeds Before they Germinate
Contains an Effective Preemergent Herbicide for Control of Crabgrass
and Many Other Nuisance Grasses and Broadleaf Weeds
Also Controls Barnyardgrass, Annual Bluegrass (*Poa annua*), Crabgrass,
Chickweed, Foxtails, Goosegrass, Henbit, Pigweed and Spurge
Feeds Your Desirable Grass up to 4 Months with a Complete Fertilizer
Featuring Pennington's Coated Slow Release Nitrogen

| | |
|---------------------------------------|---------------|
| Active Ingredient: | |
| Prodiamine (CAS No. 29091-21-2) | 0.43% |
| Other Ingredients | <u>99.57%</u> |
| Total | 100.00% |



Keep Out of Reach of Children
CAUTION

See back panel for First Aid and additional Precautionary Statements.


EPA Reg. No. 59144-12

EPA Est. _____

Net Wt: _____ lbs.

 1-888-648-8955
 1-800-424-9300

Barricade® is a Registered Trademark of Syngenta Crop Protection



5
15

Guaranteed Fertilizer Analysis 32-4-6
[Code]

Total Nitrogen (N) 32.00%
 ____% Ammoniacal Nitrogen
 ____% Urea Nitrogen
 Available Phosphate (P₂O₅) 4.00%
 Potassium (K₂O) 6.00%
 Derived from: Urea, Ammonium Phosphate, Muriate of Potash.

{Back Panel}

First Aid

| | |
|---|---|
| If in Eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If on Skin: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Immediately rinse skin with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| If Inhaled: | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. [You may also contact 1-800-424-9300 for emergency medical treatment information.] | |

{Note: The First Aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Thoroughly wash with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

Environmental Hazards


This product has low solubility in water. At the limits of solubility this product is not toxic to fish. However, at concentrations above the level of water solubility it may be toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk or tree, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergent herbicide that provides residual control of many grass and broadleaf weeds in

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals



- established perennials and wildflower plantings.

This product controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control is obtained when this product is activated by at least 1/2 inch of rainfall or irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown for sale or other commercial use, commercial seed production or research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses (excluding putting greens) or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product. Do not apply this product through any type of irrigation system. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury or illegal residues.

Application Directions



Uniformly apply using suitable, calibrated application equipment.

Established Turf

This product is a selective preemergent herbicide that, when properly applied, controls certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of product that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label.

Most effective weed control in turf grasses is obtained when this product is activated by at least 1/2 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See map for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES



Use Precautions

The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut treated sod within 120 days after application. Do not apply to newly set sod until the following year. (4) Application to turf stressed by drought, low fertility or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (6) Do not apply this product to putting greens or areas where dichondra, Colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

Rates of Application

Apply this product as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Base the maximum use rate selection on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

| Maximum Application Rate of This Product per Calendar Year By Turf Grass Species ¹ | | | |
|--|-----------------|--------------------------|-----------|
| Turf Species | Lbs. Product /A | Lbs. Product/100 Sq. Ft. | Lbs. AI/A |
| Bermudagrass ² , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia | 349 | 0.80 | 1.5 |
| Buffalograss, Kentucky Bluegrass, Perennial Ryegrass | 233 | 0.54 | 1.0 |
| Creeping Red Fescue | 174 | 0.40 | 0.75 |
| Creeping Bentgrass | 151 | 0.35 | 0.65 |

/A = per Acre A.I. = Active Ingredient

¹These are the maximum rates per calendar year by species limitations.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 121 lbs./A (0.3 lbs./100 sq. ft.). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Only suppresses Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

- Do not apply more than 349 lbs./A (0.8 lbs./100 sq. ft.) of this product per calendar year.
- Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the **Maximum Annual Rates** table.

Weeds Controlled

When used in accordance with this label this product controls the following weeds:

| | | |
|--|---------------------------------|--------------------------------|
| Barnyardgrass | Henbit | Signalgrass, Broadleaf |
| Bluegrass, Annual (<i>Poa annua</i>) | Itchgrass | Speedwell, Persian |
| Carpetweed | Johnsongrass (from seed) | Purslane, Common |
| Chickweed, Common | Junglerice | Pusley, Florida |
| Chickweed, Mouseear (from seed) | Knotweed | Rescuegrass ² |
| Crabgrass (Large, Smooth) | Kochia | Shepherd's Purse ³ |
| Crowfootgrass | Lambsquarter, Common | Sprangletop |
| Cupgrass, Woolly | Lovegrass | Spurge, Prostate |
| Foxtails, Annual | Panicum (Texas, Fall, Browntop) | Witchgrass |
| Goosegrass ¹ | Pigweed | Woodsorrel, Yellow (from seed) |

¹In many areas a single application of 151 to 349 lbs./A (0.35 to 0.80 lbs./100 sq. ft.) of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 151 to 233 lbs./A (0.35 to 0.54 lbs./100 sq. ft.) followed 60 to 90 days later by a second application that does not exceed the maximum annual rate. Do not exceed the maximum rate for turf grass species listed in the **Maximum Annual Rate** table above.

²Suppression only.

³Apply in late summer, fall or winter prior to germination.

Sequential applications may be made provided the total amount of product applied does not exceed the recommended maximum annual application rate for each turf species. All applications must be made prior to germination of the weed seeds.

When to Apply After Overseeding Turf

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

When to Overseed after Application

This product inhibits the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding results.

| Rate of This Product | | | Months Before Overseeding | | |
|----------------------|--------------------|--------|---------------------------|------------|-------|
| Lbs. / A | Lbs. / 100 Sq. Ft. | AI / A | North | Transition | South |
| 116 | 0.27 | 0.50 | 4 | 4 | 4 |
| 151 | 0.35 | 0.65 | 5 | 4 | 4 |
| 174 | 0.40 | 0.75 | 6 | 5 | 5 |
| 186 | 0.43 | 0.80 | | 6 | 6 |
| 233 | 0.54 | 1.00 | | 7 | 7 |
| 265 | 0.61 | 1.14 | | | 9 |
| 302 | 0.69 | 1.30 | | | 10 |
| 349 | 0.80 | 1.50 | | | 12 |

Landscape Ornamentals (including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergent weed control in ornamentals.

Use Rates

Apply this product at 151 to 349 lbs./A (0.35 to 0.80 lbs./100 sq. ft.) in fall and/or spring. Use higher application rates for longer control periods. Sequential or single applications may be made provided the total amount of product applied does not exceed the maximum annual application rate of 349 lbs./A.

Application Timing and Information

This product is a preemergent herbicide and will not control emerged weeds. Most effective weed control in ornamentals is obtained when this product is activated by at least 1/2 inch of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when this product is applied to soil free of clods, weeds and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergent herbicide.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

Newly-Transplanted Landscape Ornamentals

This product may be applied to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle around new transplants and thoroughly water before applying this product.

Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species

This product will not harm most ornamental trees, shrubs, vines and flowers. The species listed in Table 1 are tolerant to this product. This product is also approved for application, except in California, to the species in Table 2. Best results are obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots as this may prevent uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress, such as heat, drought or frost damage, some cultivars of listed plants may be sensitive to this product.

Table 1 – Tolerant Ornamental Species (All States)

| Scientific Name | Common Name |
|----------------------------------|---|
| <i>Abies</i> spp. | Fir species** (Balsam, Fraser, Noble, etc.) |
| <i>Acer palmatum</i> | Japanese Maple |
| <i>Acer platanoides</i> | Norway Maple |
| <i>Actinidia chinensis</i> | Kiwi* |
| <i>Agapanthus africanus</i> | Lily-of-the-Nile (African Lily) |
| <i>Arctostaphylos densiflora</i> | Vine Hill Manzanita |
| <i>Arctotheca calendula</i> | Cape Weed |
| <i>Aucuba japonica</i> | Japanese Aucuba |
| <i>Barberis gladwynensis</i> | Barberry |
| <i>Barberis julianae</i> | Wintergreen Barberry |
| <i>Barberis mentorensis</i> | Mentor Barberry |
| <i>Barberis thunbergii</i> | Japanese Barberry |
| <i>Barberis verruculosa</i> | Warty Barberry |
| <i>Buxus microphylla</i> | Japanese Boxwood |
| <i>Callistemon viminalis</i> | Weeping Bottlebrush |
| <i>Calluna vulgaris</i> | Scotch Heather |
| <i>Carpobrotus edulis</i> | Hottentot Fig (Ice Plant) |
| <i>Cassia artemisoides</i> | Feathery Cassia |
| <i>Ceanothus rigidus</i> | Wild Lilac |
| <i>Chamaecyparis pisifera</i> | False Cypress |
| <i>Cleyera japonica</i> | Cleyera |
| <i>Citrus</i> spp. | Citrus species* |
| <i>Cornus florida</i> | Flowering Dogwood |
| <i>Cornus stolonifera</i> | American Dogwood |
| <i>Cortaderia sellonana</i> | Pampas Grass |
| <i>Cotoneaster apiculatus</i> | Cranberry Cotoneaster |
| <i>Cotoneaster buxifolius</i> | Cotoneaster |
| <i>Cotoneaster dammeri</i> | Bearberry Cotoneaster |

| Scientific Name | Common Name |
|---------------------------------|--|
| <i>Cotoneaster microphyllus</i> | Rockspray Cotoneaster |
| <i>Cretaeagus</i> spp. | Hawthorne |
| <i>Cupressus sempervirens</i> | Italian Cypress |
| <i>Delasperma alba</i> | White Trailing Ice Plant |
| <i>Dodonea viscosa</i> | Hop Bush |
| <i>Elaeagnus pungens</i> | Silverberry |
| <i>Euonymus fortunei</i> | Wintercreeper |
| <i>Euonymus japonica</i> | Japanese Spindle Tree (Evergreen Euonymus) |
| <i>Euonymus kiautschovicks</i> | Spreading Euonymus |
| <i>Fatsia japonica</i> | Japanese Aralia |
| <i>Forsythia intermedia</i> | Border Forsythia |
| <i>Forsythia viridissima</i> | Greenstem Forsythia |
| <i>Gardenia jasminoides</i> | Gardenia, Cape-Jasmine |
| <i>Gladiolus</i> spp. | Gladiolus species** |
| <i>Hedera helix</i> | English Ivy |
| <i>Hibiscus</i> | Rose of Sharon** |
| <i>Hibiscus Rosa-sinensis</i> | Chinese Hibiscus |
| <i>Ilex cornuta</i> | Chinese Holly |
| <i>Ilex crenata</i> | Japanese Holly |
| <i>Ilex opaca</i> | American Holly |
| <i>Ilex pemyi</i> | Holly |
| <i>Ilex vomitoria</i> | Yaupon Holly |
| <i>Iris</i> spp. | Iris species** |
| <i>Jasminium nudiflorum</i> | Winter Jasmine |
| <i>Juniperus chinensis</i> | Chinese Juniper |
| <i>Juniperus conferta</i> | Shore Juniper |
| <i>Juniperus horizontalis</i> | Creeping Juniper |
| <i>Juglans</i> spp. | Walnut* |
| <i>Justicia brandegeana</i> | Shrimp Plant |
| <i>Lagerstromia indica</i> | Crepe Myrtle |
| <i>Ligustrum amurense</i> | Amur Privet |
| <i>Ligustrum japonicum</i> | Japanese Privet |
| <i>Ligustrum lucidum</i> | Glossy Privet (Wax-leaf) |
| <i>Liriope muscari</i> | Big Blue Lilyturf |
| <i>Lonicera japonica</i> | Japanese Honeysuckle |
| <i>Lonicera tatarica</i> | Tatarian Honeysuckle |

11
 15

| Scientific Name | Common Name |
|---------------------------------|--|
| <i>Magnolia</i> spp. | Magnolia species** |
| <i>Malephora luteola</i> | Ice Plant |
| <i>Malus</i> spp. | Crabapple* |
| <i>Nandina domestica</i> | Heavenly Bamboo |
| <i>Narcissus</i> spp. | Narcissus species** |
| <i>Nerium</i> spp. | Oleander |
| <i>Olea europaea</i> | Olive* |
| <i>Ophiopogon japonicus</i> | Mondo Grass |
| <i>Osteospermum fruticosum</i> | Trailing African Daisy |
| <i>Oxydendrum arboreum</i> | Sourwood |
| <i>Persea americana</i> | Avodaco* |
| <i>Photinia fraseri</i> | Frasier's Photinia (Redtip) |
| <i>Picea</i> spp. | Spruce species** (Colorado Blue, Norway, etc.) |
| <i>Pieris japonica</i> | Lily-of-the-Valley Shrub |
| <i>Pinus brutia</i> | Calabrian Pine |
| <i>Pinus canariensis</i> | Canary Island Pine |
| <i>Pinus elliottii</i> | Slash Pine |
| <i>Pinus halepensis</i> | Aleppo Pine |
| <i>Pinus nigra</i> | Austrian Black Pine |
| <i>Pinus palustris</i> | Longleaf Pine |
| <i>Pinus radiata</i> | Monterey Pine |
| <i>Pinus strobus</i> | Eastern White Pine |
| <i>Pinus sylvestris</i> | Scotch Pine |
| <i>Pinus taeda</i> | Loblolly Pine |
| <i>Pinus thunbergiana</i> | Japanese Black Pine |
| <i>Pinus virginiana</i> | Virginia Pine |
| <i>Pistachio</i> spp. | Pistachio* |
| <i>Pittosporum rhombifolium</i> | Queensland Pittosporum |
| <i>Pittosporum tobira</i> | Japanese Pittosporum |
| <i>Podocarpus macrophyllus</i> | Japanese Yew |
| <i>Prunus laurocerasus</i> | English Laurel |
| <i>Prunus</i> spp. | Almond, Apricot, Nectarine, Peach, Plum and Prune* |
| <i>Pseudotsuga menziesii</i> | Douglas Fir** |
| <i>Pyracantha coccinea</i> | Firethorn Scarlet |
| <i>Pyracantha fortuneana</i> | Firethorn |
| <i>Pyracantha koidzumii</i> | Firethorn |

| Scientific Name | Common Name |
|--|---|
| <i>Pyrus</i> spp. | Bradford Pear spp. |
| <i>Quercus rubra</i> | Oak species |
| <i>Raphiolepis indica</i> | Indian Hawthorne |
| <i>Rhododendron</i> (including Azalea) | "Coral Bells," "Formosa," "Hino-crimson," "PJM," "Roseum Elegans" |
| <i>Rosa banksiae</i> | Lady Bank's Rose |
| <i>Rosmarinus officinalis</i> | Rosemary* |
| <i>Rumohra adiantiformis</i> | Leatherleaf Fern |
| <i>Santolina virens</i> | |
| <i>Sedum album</i> | Stonecrop |
| <i>Syzygium paniculatum</i> | Japanese Boxcherry |
| <i>Taxus cuspidata</i> | Japanese Yew |
| <i>Taxus media</i> | Yew |
| <i>Thuja occidentalis</i> | American Arborvitae |
| <i>Trachelospermum asiaticum</i> | Star Jasmine |
| <i>Tsuga canadensis</i> | Canada Hemlock |
| <i>Tulipa</i> spp. | Tulip species |
| <i>Viburnum japonicum</i> | Japanese Viburnum |
| <i>Viburnum odoratissimum</i> | Sweet Viburnum |
| <i>Viburnum plicatum</i> | Japanese Snowball |
| <i>Viburnum rigidum</i> | Canary island Viburnum |
| <i>Viburnum tinus</i> | Laurustinus |
| <i>Viburnum trilobum</i> | Cranberry Bush |
| <i>Viburnum wrightii</i> | Leatherleaf Viburnum |
| <i>Vinca major</i> | Vinca |
| <i>Vinca minor</i> | Dwarf Periwinkle |
| <i>Vitis</i> spp. | Grape* |
| <i>Weigela florida</i> | Old Fashioned Weigela |
| <i>Yucca aloifolia</i> | Spanish Bayonet |
| <i>Yucca filamentosa</i> | Yucca, Adam's Needle |

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.

Table 2 – Tolerant Ornamental Species/Varieties (All States Except California)

| Scientific Name | Common Name |
|------------------------------|---------------------|
| <i>Abelia grandiflora</i> | Abelia: Sherwood |
| <i>Achillea</i> spp. | Yarrow: King Edward |
| <i>Agapanthus orientalis</i> | |

| Scientific Name | Common Name |
|------------------------------------|---|
| <i>Akebia quinata</i> | Five-Leaf or Chocolate Vine |
| <i>Allium cernuum</i> | Lady's Leek, Nodding Onion |
| <i>Anemone hybrida</i> | Japanese Anemone |
| <i>Aquilegia</i> spp. | Aquilegia: Red and Gold |
| <i>Artemisia</i> spp. | Wormwood: Silver Mound, Castle |
| <i>Aster</i> spp. | Aster: Bonny Blue, Purple Dome |
| <i>Aster X frikartii</i> | |
| <i>Athyrium filix-femina</i> | Lady Fern; Fern Lady |
| <i>Begonia</i> spp. | Fibrous Begonia: Hardy Grandis |
| <i>Bergenia cordifolia</i> | |
| <i>Boltonia asteroides</i> | Snowbank |
| <i>Bougainvillea</i> spp. | Bougainvillea |
| <i>Buddleia davidii</i> | Butterfly-Brush (Dwarf Blue); Royal Red |
| <i>Callistemon citrinus</i> | Crimson Bottlebrush |
| <i>Campanula carpatica</i> | Tussock Bellflower (White Clips) |
| <i>Campis X tagliabuana</i> | Trumpet Creeper, Trumpet Flower; Madame Galen |
| <i>Ceratostigma plumbaginoides</i> | |
| <i>Crhysanthemum nipponicum</i> | |
| <i>Coreopsis</i> spp. | Coreopsis (Calliopsis): Early Sunrise, Moonbeam |
| <i>Crocosmia</i> spp. | Lucifer |
| <i>Delosperma</i> spp. | Cooperi Pink |
| <i>Delphinium</i> spp. | Larkspur; Blue Elf |
| <i>Dianthus deltoides</i> | Dianthus, maiden Pinks "Zing" |
| <i>Dianthus gratianopolitanus</i> | Cheddar Pink |
| <i>Echinacea pupurea</i> | Coneflower, Purple; Magnus |
| <i>Forsythia suspensa</i> | Weeping Forsythia |
| <i>Gaillardia</i> spp. | Gaillardia, Blanket Flower: "Goblin" |
| <i>Gentiana dahurica</i> | Gentian |
| <i>Geranium cinereum</i> | Cranesbill |
| <i>Gypsophila repens</i> | Baby's Breath |
| <i>Helianthemum</i> spp. | Sunrose |
| <i>Hemerocallis</i> spp. | Daylily: Aztec Gold, Stella De Oro, Tender Love |
| <i>Heucherella</i> spp. | Coral Bell; Bridget Bloom |
| <i>Hibiscus</i> spp. | Mallow; Disco Belle White |
| <i>Hosta plantaginea</i> | Hosta, Plantain Lily (Fragrant) |
| <i>Hosta sieboldiana</i> | Hosta, "Searsucker" |

14
 15

| Scientific Name | Common Name |
|---|--|
| <i>Houttuynia cordata</i> var. <i>variegata</i> | |
| <i>Hydrangea macrophylla</i> | Bigleaf Hydrangea |
| <i>Inula ensifolia</i> | |
| <i>Iris ensata</i> | Sword-Leaved Iris; Jodlesong |
| <i>Iris siberica</i> | Siberian Iris; Cabernet |
| <i>Juniperus davurica</i> | Parsoni |
| <i>Lagerstromia indica</i> x <i>fauriel</i> | Crape Myrtle; Tuscarora |
| <i>Lantana montevidensis</i> | Weeping Lantana |
| Lavender spp. | Lavender; Munstead |
| <i>Leontopodium alpinum</i> | Edelweiss |
| <i>Ligstrum sinense</i> | Chinese Privet; Variegata |
| <i>Lilium</i> spp. | Lily: Jazz |
| <i>Lirope muscari</i> var. <i>variegata</i> | Lirope, Variegated |
| <i>Lirope spicata</i> | Lirope, Creeping |
| <i>Lobelia cardinalis</i> | Cardinal Flower, Indian Pink |
| <i>Loropetalum chinense</i> q | Burgundy |
| <i>Lythrum</i> spp. | Loosestrife; Modern Pink |
| <i>Miscanthus sinensis</i> | Yaku Jima, Silberfeder |
| <i>Oenothera missourensis</i> | Evening Primrose |
| <i>Osmanthus heterophyllus</i> | Osmanthus (False Holly): Gulf Tide |
| <i>Paeonia suffruticosa</i> | Tree Peony |
| <i>Pennisetum setaceum</i> | Fountain Grass (Dwarf) |
| <i>Perovskia atriplicifolia</i> | |
| <i>Physostegia virginiana</i> | Dragonhead, False; Vivid |
| <i>Quercus shumardii</i> | Oak, Shumard's Red |
| <i>Rhaphiolepis umbellata</i> | Yedda Hawthorne |
| <i>Rhododendron</i> (including azalea) | "Delaware Valley White," "Flame Creeper," "Girard Crimson," "George L. Tabor," Wakeiebisu, White Gumpo |
| <i>Rudbeckia</i> spp. | Black-Eyes Susan: Goldstrum |
| <i>Saxifraga</i> spp. | Saxifrage; Purple Dome |
| <i>Scabiosa</i> spp. | Pincushion Flower |
| <i>Sedum cauticola</i> | Stonecrop; Lidakense |
| <i>Sedum dasyphyllum</i> | Stonecrop |
| <i>Sedum spurium</i> | Stonecrop; Dragon's Blood |
| <i>Spiraea bumalda</i> | Spiraea: Anthony Waterer |
| <i>Syzyglum paniculatum</i> | Australian Brushcherry |
| <i>Teucrium</i> spp. | Germander |

15
 15

| Scientific Name | Common Name |
|---------------------------------|-----------------------------------|
| <i>Thalictrum dipterocarpum</i> | Meadow Rue |
| <i>Veronica</i> spp. | Veronica, Speedwell; Sunny Border |
| <i>Viburnum suspensum</i> | Arrowood Virburmum |

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.

Spreader Settings

Rotary spreaders are easier to use and reduce streaking that's often caused by drop spreaders. While drop spreaders work well, they require more care to avoid gaps and streaks that appear as light green stripes about a week after fertilizer application. Always apply with a properly calibrated rotary or drop type fertilizer spreader. Never apply by hand. Refer to and follow the spreader manufacturer's calibration and setting instructions.

The above spreader settings are approximate. Calibrate your spreader on 100 square feet and adjust to apply at the rate of 0.8 pound of product per 100 square feet.

{For household/domestic use product in bags up to 50 pounds:}

[Storage and Disposal]

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original bag in a cool, dry (preferably locked) storage area inaccessible to children and pets. Store away from feed or food stuffs and separated from other fertilizers and pesticides. Keep bag closed when product is not being used. **Pesticide/Container Disposal: If empty:** Non-refillable container; do not reuse or refill this container. Offer for recycling, if available; otherwise, discard in trash. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

{Per PR Notice 2007-4 the batch code / lot number will appear on the label or container.}

Warranty and Disclaimer: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. To the extent consistent with applicable law, the manufacturer is not responsible for the use of this product contrary to the label instructions, under abnormal conditions or under conditions not reasonably foreseeable to the manufacturer and/or seller, and buyer assumes the risk of any such use. If you decide, after reading this label, that you can't use this product immediately return the unopened package to the point of purchase.

GRO TEC, INC.
 P.O. Box 290
 Madison, GA 30650
 1-800-285-SEED

[] Denotes alternate/optional verbiage
 { } Denotes verbiage not printed on market labeling