MASTER LABEL

AGRISPON®


A biologically derived biostimulant for soil and plants.

<u>Ingredients</u>

Active Ingredient:	
Plant Extract*	0.56%
Other Ingredient:	99.44%
Total	

^{*} The plant extract is derived from <u>Quercus falcata</u>, <u>Opuntia lindheimeri</u>, <u>Rhus aromatica</u>, and <u>Rhizophoria mangle</u> tissues.

Keep out of reach of children CAUTION



Manufactured by: Agriculture Sciences, Inc. 3601 Garden Brook Dallas, TX 75234

EPA Reg. No. 59174-3 EPA Est. No. 59174-TX-1

Agrispon <u>Agricultural/Commercial</u> Use Sublabel. This label contains application rates specific to 32 fl. oz., 1, 2.5, 30, 55, 275, and 300 gallon containers.

AGRISPON®

A biologically derived biostimulant for soil and plants.

<u>Ingredients</u>	
Other Ingredient:	
* The plant extract is derived from Quercus fa aromatica, and Rhizophoria mangle tissues.	alcata, <u>Opuntia lindheimeri, Rhus</u>
Keep out of rea	ach of children
CAUT	TION
See Side Pane	el for First Aid
CHEMIGATION: Refer to supplement use directions for chemigation. Do irrigation system unless the supple followed.	not apply this product through any
Use this product only in accordance of Protection Standard, 40 CFR Part 11 entitled AGRICULTURAL USE RECEIVED FOR USE section of the labeling for	70. Refer to supplemental labeling RUIREMENTS in the DIRECTIONS
Manufactured by: Agriculture Sciences, Inc. 3601 Garden Brook	EPA Reg. No. 59174-3
Dallas, TX 75234	EPA Est. No. 59174-TX-1
Net contentsfl. oz.	This container will treat acres.

Agrispon Ag Sublabel Page 2

AGRISPON®

Agrispon stimulates root development and beneficial soil micro-organisms, increasing the efficiency of nutrient uptake from soil and improving soil structure over time. Agrispon can also hasten certain plant growth processes and improve a plant's ability to withstand pest and environmental stresses. Agrispon enables the soil to supply nutrients to the plant more efficiently.

See attachment for DIRECTIONS for USE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION Avoid contact with skin or clothing. Wash throughly with soap and water after handling.

FIRST AID

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

long-sleeved shirt and long pants waterproof gloves shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately, if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Last Edited March 6, 2000

WARRANTY

The Manufacturer warrants that this product conforms to the original formulation and is fit for use as directed. Neither Manufacturer nor seller shall be liable for any injury, loss or damage direct or indirect arising from the misuse of the product. Agriculture Sciences, Inc. and its various sellers' only obligation shall be to replace such quantity of the product that is proven defective before purchase.

ATTACHMENT EPA REG. No. 59174-3

AGRISPON®

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements in this labeling about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

coveralls waterproof gloves shoes plus socks.

USES

Agrispon may be used on all food and feed crops, consistent with the use directions and restrictions stated below. Agrispon may also be used for horticultural applications, as noted below.

EQUIPMENT AND MIXING

Apply Agrispon via boom, fan jet, hand-carried, or backpack spray equipment. This product may be applied through drip and overhead irrigation systems. Follow all chemigation directions below. Dilute Agrispon in water as specified in the table below:

Treatment Area	Agripson	Water (minimum)*
5,000 square feet	1.5 fluid ounces	10 gallons
1 acre	13 fluid ounces	90 gallons
10 acres	1 gallon	900 gallons

^{*}The table specifies the minimum amount of water to be used; Agrispon may diluted in a larger volume of water if desired. The amount of water used in the mix will vary according to equipment, type of nozzle used, number of nozzles, ground speed, system pressure, and calibration. If soil is covered with plant material, mulch, or thatch, use sufficient water to transport it to the soil during application or lightly irrigate after application.

Shake well before using. Agitate solutions either during or immediately after dilution. Apply solution within eight hours of mixing.

APPLICATION INSTRUCTIONS

ANNUALS

Short cycle (plants having a growth cycle of 60 days or less)

Make a single application at, or as soon after planting as possible. Maximum number of applications/area/year: 6

Mid season (plants having a growth cycle of 60 to 120 days)

Make the first application at planting or within two weeks of germination with a subsequent application 45 to 60 days after. Maximum number of applications/area/year: 6

Last Edited March 6, 2000

Long-season (plants having a growth cycle longer than 120 days)

Make an initial application at planting or within two weeks of germination. Make subsequent applications at 45 to 60 day intervals during the period when the plant is growing vigorously. Maximum number of applications/area/year: 8

PERENNIALS

Trees, Excluding Commercial Forestry Use:

Apply to soil surface under the tree extending out 3 to 4 times the distance of the trunk to drip line to insure root coverage.

Evergreen: For newly planted trees, initial application should be made at, or as soon after planting as possible. For established trees, initial application should be made when soil temperature warms to 50°F. For climates where soil temperatures are ≥50°F year round, initial application may be made at any time. Application should be repeated every 60 days as weather permits. Maximum number of applications/area/year: 6

Deciduous: Make first application at bud break or leaf flush in the spring. Make subsequent application every 60 days until dormancy occurs. Maximum number of applications/area/year: 6

GREENHOUSE/GARDEN

Apply every 6 to 8 weeks during active growth. Number of applications will depend on the growth cycle length as explained above.

CHEMIGATION OF AGRISPON

General Information

Apply this product only through drip (trickle) or sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

MIXING AND APPLICATION

)

The following instructions apply to all chemigation methods discussed on this labeling.

Determine the number of acres to be treated by the chemigation system. Prepare a premix by adding 13 fluid ounces of AGRISPON and a minimum of one half gallon of water for each acre to be treated into a reservoir container. The use of larger quantities of water to dilute the premix may make calibration of the application easier. Meter the premix into the chemigation system at a rate that will consume the entire premix within the period of chemigation or within 8 hours, whichever is less. Maintain agitation in the reservoir during the period of chemigation to keep material in suspension.

OBSERVE THE FOLLOWING PRECAUTIONS IF YOUR CHEMIGATION SYSTEM IS CONNECTED TO A PUBLIC WATER SYSTEM

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of a year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in the cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

STATEMENTS CONCERNING THE OPERATION OF SPRINKLER CHEMIGATION; UTILIZING A PRESSURIZED WATER AND PESTICIDE INJECTION SYSTEM.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quickclosing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

STATEMENTS CONCERNING THE OPERATION OF DRIP (TRICKLE) CHEMIGATION; UTILIZING A PRESSURIZED WATER AND PESTICIDE INJECTION SYSTEM.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-

operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

STORAGE AND DISPOSAL

)

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container only.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed

of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or

reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of

smoke.

Household Use Sublabel

AGRISPON®

A biologically derived biostimulant for soil and plants. For residential use only

ing	redi	er	nts

Active Ingredient:	
Plant Extract*	0.56%
Other Ingredient:	99.449
Total	100.00%

Keep out of reach of children CAUTION

See Side Panel for First Aid

Manufactured by: Agriculture Sciences, Inc. 3601 Garden Brook Dallas, TX 75234

EPA Reg. No. 59174-3 EPA Est. No. 59174-TX-1

Net contents	fl. oz.
This container will treat	square feet.

^{*} The plant extract is derived from <u>Quercus falcata</u>, <u>Opuntia lindheimeri</u>, <u>Rhus aromatica</u>, and <u>Rhizophoria mangle</u> tissues.

AGRISPON®

Agrispon stimulates root development and beneficial soil micro-organisms, increasing the efficiency of nutrient uptake from soil and improving soil structure over time. Agrispon can also hasten certain plant growth processes and improve a plant's ability to withstand pest and environmental stresses. Agrispon enables the soil to supply nutrients to the plant more efficiently.

See attachment for DIRECTIONS for USE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

1

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

WARRANTY

The Manufacturer warrants that this product conforms to the original formulation and is fit for use as directed. Neither Manufacturer nor seller shall be liable for any injury, loss or damage direct or indirect arising from the misuse of the product. Agriculture Sciences, Inc. and its various sellers' only obligation shall be to replace such quantity of the product that is proven defective before purchase.

ATTACHMENT EPA Reg. No. 59174-3 AGRISPON®

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Keep children and pets out of the treated area until sprays have dried.

USES

Agrispon may be used on all food and feed crops, consistent with the use directions and restrictions stated below. Agrispon may also be used for horticultural applications, as noted below.

EQUIPMENT AND MIXING

<u>For application via watering can, hand trigger sprayer, or pressurized sprayer</u>: Dilute Agrispon in water according to the table below.

Treatment Area	Agripson	Water (minimum)*
50 square feet	5 drops	1 quart
500 square feet	1 teaspoon	1 gallon
1,000 square feet	2 teaspoons	2 gallons

^{*} If soil is covered with plant material, mulch, or thatch, dilute Agrispon in a larger volume of water to ensure transport to the soil. Alternatively, lightly irrigate after application.

<u>For hose-end sprayer</u>: put 1 teaspoon Agrispon in sprayer reservoir and fill reservoir with water to 10 oz. Set sprayer to deliver 10 tablespoons/gallon of water. This will treat 500 square feet.

Shake well before using. Agitate solutions either during or immediately after dilution. Apply solution within eight hours of mixing.

APPLICATION INSTRUCTIONS

ANNUALS

Short cycle (plants having a growth cycle of 60 days or less)

Make a single application at, or as soon after planting as possible. Maximum number of applications/area/year: 6

Mid season (plants having a growth cycle of 60 to 120 days)

Make the first application at planting or within two weeks of germination. Make subsequent applications every 45 to 60 days. Maximum number of applications/area/year: 6

Long-season (plants having a growth cycle longer than 120 days)

Make an initial application at planting or within two weeks of germination. Make subsequent applications at 45 to 60 day intervals during the period when the plant is growing vigorously. Maximum number of applications/area/year: 8

PERENNIALS

Trees:

Apply to soil surface under the tree extending out 3 to 4 times the distance of the trunk to drip line to ensure root coverage.

Evergreen: For newly planted trees, initial application should be made at, or as soon after planting as possible. For established trees, initial application should be made when soil temperature warms to 50°F. For climates where soil temperatures are ≥50°F year round, initial application may be made at any time. Application should be repeated every 60 days as weather permits. Maximum number of applications/area/year: 6

Deciduous: Make first application at bud break or leaf flush in the spring. Make subsequent application every 60 days until dormancy occurs. Maximum number of applications/area/year: 6

GREENHOUSE/GARDEN

Apply every 6 to 8 weeks during active growth. Number of applications will depend on the growth cycle length as explained above.

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Store in original container only.

CONTAINER DISPOSAL Securely wrap original container in several layers of

newspaper and discard in trash.