Antimica Jed Program Bearing Respect rate as Date Saion Com - Helicy

Thelosur

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

EPA Form 8570-6 (Rev. 5-76)

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.

Reg no 59370-1

ACCEPTED
with COMMENTS
in EPA Letter Dated:

AQUACIDE 601

SEP 23 1988

Under the Pederal Insecticide, Europeide una Redenticide Act as promited, for the pesticide registered under EPA Reg. No.

59371-

ACTIVE INGFIEDIENT:

This product contains 1.29 lb, of active ingredient per gallon and weighs 8.58 lb, per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARUS TO HUMANS AND DOMESTIC ANIMALS

...

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skipt pypt, or clothing.

FIRST AID: Inswallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of takes, streams, or ponds. Permits may be required for discharges containing this posticide into takes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

DIRECTIONS FO

It is a violation of Federal law to use this pro
ts labeling. AQUACIDE 601
and fung; in recirculating commercial and ind o its use, systems must be cleaned to rem
lime, and other deposits. An initial slug additional slug addi
of water to provide a concentration of 32 AQUACIDE 601 , t
n the system, is recommended. Repeat initia Subsequent slug additions of 1.0
AQUACIDE 601 per 100 80 parts per million of AQUACIDE
employed every 2 to 5 days, or as needed. T upon the relative amount of bleedoff and the s
em. Slug additions should be made in the su AQUACIDE 601 is us
air-washing systems that maintain effective notes use, systems should be cleaned to a deposits. An initial slug dose of14.8
mended. Repeat initial dosage until control tions of 10.1 to 24.7 fluid ounces of per 1000 gallons of water should be employed frequency of addition depends upon the relation the bacterial problem. Slug additions may be collection trays of the airwash system.

STORAGE AND D

Do not contaminate water, food, or feed by storage or disposal STORAGE: Do not stack more than four drums high, Leakin drums for disposal. Spills should be absorbed in sawdust or sall tainer closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of the proved waste disposal facility.

CONTAINER DISPOSAL:

- C PLASTIC Tiple imse for equivalent: Then offer for recyclin sanitary landfill, by incineration, or, if allowed by state and smoke
- strinke
 METAL Triple insertor equivalent) Then offer for recycling sanitary landfill, or by other procedures approved by state a

		Manufacture
	AQUARICH- W	VATER TREATME
	33672 DOREK	
-	FRASER, MI	48026
COA DCC	NO	