

PM 23 59639-3 10/5

SELECT 2 EC Herbicide
EPA Reg. No. 59639-3
July 16, 1992

The use of this product may pose a hazard to the federally designated endangered species of Solano Grass and Wild Rice. Use of this product is prohibited in the following areas where the species are known to exist:

Solano Grass: Solano County, California: the vernal lakes area bounded by the Union Pacific Railroad and Hastings Road to the north, Highway 113 to the east, Highway 12 to the south, and Travis Air Force Base to the west.

Wild Rice: Hays County, Texas.

ACCEPTED
13 AUG 1992
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 59639-3



SELECT[®]

2 EC HERBICIDE

Active Ingredient: Clethodim 25%
Inert Ingredients: 75%
Contains Petroleum Distillates
*E-2[1-[[[3-chloro-2-propenyl]oxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one
Contains 2.0 lbs. clethodim per gal.
U.S. Pat. No. 4,440,566

KEEP OUT OF REACH OF CHILDREN
WARNING—AVISO
SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS
PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING

Causes eye irritation. Avoid contact with skin. Do not get in eyes or on clothing. Wear goggles, chemical-resistant gloves, long-sleeved shirt, long-legged pants, shoes and socks. Harmful if swallowed or inhaled. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT
In case of eye contact, flush eyes immediately with fresh water for 15 minutes. If swallowed, give water or milk to drink and telephone for medical advice. DO NOT make person vomit unless directed to do so by medical personnel. If medical advice cannot be obtained, then take person and product container to the nearest medical emergency treatment center or hospital. If on skin, wash with plenty of soap and water. Get medical attention if irritation persists. If inhaled, remove person from exposure area.

Note to Physicians: Emergency Information—call 1-800-457-2022. This product contains a light petroleum solvent, and an aspiration hazard may exist.

ENVIRONMENTAL HAZARDS
Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply where weather conditions favor drift from areas treated.

Prohibited for use in these counties to avoid non-target injury to the endangered grass species of:
Sesuvia Grass—(California) Colusa, Contra Costa, Fresno, Glenn, Madera, Merced, San Joaquin, Solano, Stanislaus and Tehama counties.
Wild Rice—(Texas) Hays County.

PHYSICAL OR CHEMICAL HAZARDS
Do not use or store near heat or open flame.

DIRECTIONS FOR USE
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL AND PAMPHLET ACCOMPANYING THIS PRODUCT IN CONJUNCTION WITH PRECAUTIONARY STATEMENTS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

REENTRY STATEMENTS
Keep all unprotected persons out of treated areas until they have been notified there may be drift. Do not enter treated areas without proper protective clothing.

Certain states may require more restrictive labeling. Contact your State Department of Agriculture for further information. Written or oral warnings regarding user safety and drift exposure must be given to workers in treated areas or on areas about to be treated.

GENERAL INFORMATION

Do not apply this product through any irrigation system. SELECT[®] is a selective postemergence herbicide for control of annual and perennial grasses in soybeans and corn. It does not control broadleaf weeds.

Do not allow SELECT to come in contact with soybeans, corn, rice, sorghum, small grains, or other crops. Crops will be injured or killed.

Minor leaf spotting may occur on soybeans under certain environmental conditions. New foliage will be free of spotting.

Control Symptoms
Treated grass weeds show a yellowing, chlorosis/necrosis of younger plant tissue.

collapse of the remaining foliage. Symptoms will generally be observed in 7 to 14 days depending on grass species treated and environmental conditions.

APPLICATION INFORMATION

Timing of Applications
Apply SELECT postemergence to actively growing grasses according to rate table recommendations. Applications made to grass plants stressed by insufficient moisture or cold temperatures, or to grass plants exceeding recommended growth stages may result in unsatisfactory control. Do not apply under these conditions.

In arid regions where irrigation is used to supplement limited rainfall, SELECT should be applied as soon as possible after an irrigation (within 7 days). In arid regions, a second application of SELECT will generally provide more effective control of perennial grass weeds than a single application. Make second application to actively growing grass 2 to 3 weeks after emergence of new growth.

Cultivation of treated grasses 7 days prior to or within 7 days after application of SELECT may reduce weed control. DO NOT APPLY SELECT if rainfall is expected within one hour, since control may be reduced.

Ground Application
Use of sufficient spray volumes and pressure is essential to ensure complete coverage. Use a minimum of 10 gals. and a maximum of 40 gals. of spray solution per acre. Spray pressures should reflect a minimum of 20 psi and a maximum of 60 psi at the nozzle. Do not use flood nozzles.

Air Application
Use a minimum of 3 gals. of spray solution per acre. Increase spray volumes up to 10 gals. as grass or crop foliage becomes dense.

Spot Treatment
When using hand sprayers or high volume sprayers utilizing hand guns.

Mix 1/2% SELECT and 1% non-phytotoxic crop oil concentrate according to amounts shown in the table below. Treat to wet vegetation, while not allowing runoff of spray solution.

Desired Spray Vol. Gal.	SELECT/Oil To Be Added	
	SELECT Fl. Oz.	Oil Conc. Fl. Oz.
3	2	4
25	16	32
50	32	64
100	64	128

For ground or air application, add a crop oil concentrate containing at least 15% emulsifier at 1% v/v to the finished spray volume.

Annual and perennial grass weed control for cotton and soybeans

Region A
Washington, Oregon, Idaho, Montana, Wyoming, North Dakota, South Dakota, Nebraska, Kansas, Minnesota, Iowa, Missouri, Arkansas, Louisiana, Wisconsin, Illinois, Tennessee, Mississippi, Indiana, Kentucky, Ohio, Georgia, Florida, Alabama, South Carolina, North Carolina, Virginia, West Virginia, Maryland, Delaware, Pennsylvania, New Jersey, New York, Connecticut, Massachusetts, Vermont, New Hampshire, Maine, Rhode Island, Michigan, Washington D.C., Oklahoma and Texas East of Interstate 35

Region B
Arizona, California, Colorado, Nevada, New Mexico, Utah, Oklahoma and Texas West of Interstate 35.

RECOMMENDATIONS FOR ANNUAL AND PERENNIAL GRASS WEED CONTROL (EXCEPT CARNIVOROUS PLANTS)

- Region A & B**
- Apply only to actively growing grasses.
 - Apply when the first grass weed sprout reaches the recommended growth stage.
 - Always add a non-phytotoxic petroleum based crop oil concentrate to the spray tank at 1 qt./acre.
 - Do not exceed a total of 32 fl. oz./acre.

GRASS SPECIES

Seedling	Sorghum halepense
Johnsongrass	Sorghum bicolor
Shattercane	Zizania
Volunteer Corn	Avena fatua
Wild Oats	Panicum mitchellii
Wild Proso Millet	Panicum capillare
Witchgrass	Eriochloa villosa
Woolly Cupgrass	Echinochloa crus-galli
Barnyardgrass	Bracharia platyphylla
Broadleaf	Digitaria sanguinalis
Signalgrass	Digitaria ciliaris
Crabgrass	Digitaria pruriens
Large	Dactyloctenium aegyptium
Smooth	
Southern	
Crowfootgrass	

BEST AVAILABLE COPY

384

GRASS SPECIES (continued)		WEED* HEIGHT INCHES	1/RATE FL. OZ. ACRE	2/ HIGH RATE
Fall Panicum	Panicum dichotomiflorum	2-6	6	8
Field Sandbur roxtail	Cenchrus incartus	2-6	6	8
Joint Green Yellow	Setaria faberi Setaria viridis Setaria glauca	2-8 2-6 2-6	6 6 6	8 8 8
Goosegrass	Eleusine indica	2-6**	6	8
Itchgrass	Rottboellia exaltata	2-6	6	8
Junglerice	Echinochloa colonum	2-6	6	8
Red Sprangletop	Leptochloa filiformis	2-6	6	8
Red Rice	Oryza sativa	1-3	6	8
Southwestern Cupgrass	Eriochloa gracilis	2-6	6	8
Texas Panicum	Panicum polyanthemum	2-6	6	8
Volunteer Cereals				
Barley	Hordeum vulgare	2-6	6	8
Oats	Avena sativa	2-6	6	8
Rye	Secale cereale	2-6	6	8
Wheat	Triticum aestivum	2-6	6	8

* Generally occurs between 3-leaf stage and tillering

** Length of lateral growth.

1/ Apply under favorable soil moisture and humidity which exists within a few days after rainfall or within 7 days after irrigation. Apply at weed height as indicated on the label

2/ Use the high rate under heavy grass pressure and/or when grasses are at maximum height

7

RECOMMENDATIONS FOR PERENNIAL GRASSES IN SOYBEANS (EXCEPT CA) AND COTTON

Region A

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Always add a non-phytotoxic crop oil concentrate (petroleum based) to the spray tank at 1 qt./acre by ground equipment.
- Do not exceed a total of 32 fl. oz./acre (0.5 lb. ai) per season

GRASS SPECIES

GRASS SPECIES	WEED* HEIGHT INCHES	1/RATE FL. OZ. ACRE	2/ HIGH RATE
Rhizome Johnsongrass	Sorghum halepense		
First Application		12-24	8 10
Repeat Application(s) (if regrowth occurs)		6-10	6 8
Bermudagrass	Cynodon dactylon		
First Application		3 (or up to 6" runners)	8 16
Repeat Application(s) (if regrowth occurs)		3 (or up to 6" runners)	8 16
Quackgrass	Agropyron repens		
First Application		4-8	8 16
Repeat Application(s) (if regrowth occurs)		4-8	8 16

* Generally occurs between 3-leaf stage and tillering

1/ Apply under favorable soil moisture and humidity which exists within

8

a few days after rainfall or within 7 days after irrigation. Apply at weed height as indicated on the label.

2/ Use the high rate under heavy grass pressure and/or when grasses are at maximum height.

RECOMMENDATIONS FOR PERENNIAL GRASSES IN SOYBEANS (EXCEPT CA) AND COTTON

Region B

- Apply only to actively growing grasses at recommended weed heights
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Always add a non-phytotoxic crop oil concentrate (petroleum based) to the spray tank at 1 qt./acre by ground equipment.
- Do not exceed a total of 32 fl. oz./acre (0.5 lb. ai) per season.

GRASS SPECIES

GRASS SPECIES	WEED* HEIGHT INCHES	1/RATE FL. OZ. ACRE	2/ HIGH RATE
Rhizome Johnsongrass	Sorghum halepense		
First Application		6-10	8 16
Repeat Application(s) (if regrowth occurs)		4-8	6 8
Bermudagrass	Cynodon dactylon		
First Application		3 (or up to 6" runners)	8 16
Repeat Application(s) (if regrowth occurs)		3 (or up to 6" runners)	8 16

9

* Generally occurs between 3-leaf stage and tillering

1/ Apply under favorable soil moisture and humidity which exists within a few days after rainfall or within 7 days after irrigation. Apply at weed height as indicated on the label

2/ Use the high rate under heavy grass pressure and/or when grasses are at maximum height

TANK MIX WITH SELECT ZEC HERBICIDE & BROADLEAF HERBICIDES FOR CONTROL OF BROADLEAF AND GRASS WEEDS IN SOYBEANS (EXCEPT CA)

GENERAL INFORMATION REGIONS A AND B

Apply only to actively growing grass and broadleaf weeds at recommended heights listed on each label.

Apply when the first grass or broadleaf weed species in a mixed weed population reaches the recommended growth stage for treatment.

Apply under favorable soil moisture and humidity which exists within a few days after rainfall or within 7 days after irrigation.

Always add a crop oil concentrate (petroleum based) to the spray tank at the rate recommended for each tank mix using ground equipment.

Do not exceed a total of 32 fl. oz./acre (0.5 lb. ai) of SELECT ZEC Herbicide per season

Do not tank mix with SCEPTER® Herbicide.

10

BEST AVAILABLE COPY

Table 1H.
SELECT ZEC Herbicide & BASAGRAN® Tank Mix - Soybeans - Rate and Time of Application Table.

Product	Product Rate	Weeds Controlled/ Weed Size	Adjuvant
SELECT	8 to 10 fl. oz./A	Refer to Table - Regions A & B for Annual grasses controlled.	1 qt. Crop Oil Concentrate/A
	10 to 16 fl. oz./A	Refer to Tables - Region A/Region B for Perennial* grasses controlled.	
plus BASAGRAN			
BASAGRAN	1 to 2 pts./A	Refer to BASAGRAN label for weeds controlled.	

*If regrowth of Perennial grasses occurs, make a second application of SELECT ZEC Herbicide using 8 or 16 fl. oz./acre timed to grass weed and sizes noted for Perennial grasses in Region A and Region B.

SEQUENTIAL APPLICATION OF SELECT ZEC HERBICIDE AND BROADLEAF HERBICIDES

Some reduction in grass control is possible when SELECT ZEC Herbicide is sequentially applied immediately prior to or after an application of a broadleaf herbicide.

1. After an application of SELECT wait a minimum of 24 hours before applying a broadleaf herbicide.

2. In fields treated with a broadleaf herbicide, do not apply SELECT until grass plants are again developing new leaves (generally 7 days after the broadleaf herbicide application).

11

Table 2H.
SELECT ZEC Herbicide & BLAZER® Tank Mix - Soybeans - Rate and Time of Application Table.

Product	Product Rate	Weeds Controlled/ Weed Size	Adjuvant
SELECT	6 to 8 fl. oz./A	Refer to Table - Regions A & B for Annual grasses controlled.	1 pt. Crop Oil Concentrate/A
	8 to 16 fl. oz./A	Refer to Tables - Region A/Region B for Perennial* grasses controlled.	
plus BLAZER			
BLAZER	1 to 1.5 pts./A	Refer to BLAZER label for weeds controlled.	

*If regrowth of Perennial grasses occurs, make a second application of SELECT ZEC Herbicide using 8 or 16 fl. oz./acre timed to grass weed and sizes noted for Perennial grasses in Region A and B.

SEQUENTIAL APPLICATION OF SELECT ZEC HERBICIDE AND BROADLEAF HERBICIDES

Some reduction in grass control is possible when SELECT ZEC Herbicide is sequentially applied immediately prior to or after an application of a broadleaf herbicide.

1. After an application of SELECT wait a minimum of 24 hours before applying a broadleaf herbicide.

2. In fields treated with a broadleaf herbicide, do not apply SELECT until grass plants are again developing new leaves (generally 7 days after the broadleaf herbicide application).

12

Table 3M.
SELECT ZEC Herbicide & CLASSIC® Tank Mix—Soybeans—Rate and Time of Application Table.

Product	Product Rate	Weeds Controlled/ Wood Size	Adjuvant
SELECT	8 to 10 fl. oz./A	Refer to Table—Regions A & B for Annual grasses controlled.	Crop Oil Concentrate at 10% v/v
	10 to 16 fl. oz./A	Refer to Tables—Region A/Region B for Perennial* grasses controlled.	
_____ plus _____ CLASSIC			
CLASSIC	1/2 to 3/4 oz./A	Refer to CLASSIC label for weeds claimed.	

*If regrowth of Perennial grasses occurs, make a second application of SELECT 2 EC herbicide using 6 or 8 fl. oz./acre timed to grass weed and sizes noted for Perennial grasses in Region A and Region B.

SEQUENTIAL APPLICATION OF SELECT 2 EC HERBICIDE AND BROADLEAF HERBICIDES

Some reduction in grass control is possible when SELECT 2 EC Herbicide is sequentially applied immediately prior to or after an application of a broadleaf herbicide.

1. After an application of SELECT wait a minimum of 24 hours before applying a broadleaf herbicide.
2. In fields treated with a broadleaf herbicide, do not apply SELECT until grass plants are again developing new leaves (generally 7 days after the broadleaf herbicide application).

13

TANK MIX WITH SELECT 2 EC HERBICIDE & INSECTICIDES FOR CONTROL OF GRASS WEEDS AND INSECTS IN SOYBEANS (EXCEPT CA) AND COTTON

GENERAL INFORMATION REGIONS A & B

Applications of SELECT 2 EC Herbicide and ORTHENE® insecticide are compatible when tank mixed for immediate use. Under certain situations tank mixing may offer significant economic benefits. When mixing ORTHENE with SELECT 2 EC Herbicide, read the entire label of both pesticides to insure proper application timing. All application directions, restrictions, precautions and worker safety rules on the EPA registered labels are to be followed. Do not exceed a total of 32 fl. oz./acre (0.5 lb. ai) of SELECT 2 EC Herbicide per season.

Table 1I.
SELECT 2 EC Herbicide & ORTHENE Tank Mix—Cotton and Soybeans—Rate and Time of Application Table.

Product	Product Rate	Weeds and Insects Controlled	Adjuvant
SELECT	6 to 8 fl. oz./A	Refer to Table—Regions A & B for Annual grasses controlled.	1 qt. Crop Oil Concentrate/A
	8 to 16 fl. oz./A	Refer to Tables—Region A/Region B for Perennial* grasses controlled.	
_____ plus _____ ORTHENE			

16

Table 4M.
SELECT 2 EC Herbicide & COBRA® Tank Mix—Soybeans—Rate and Time of Application Table.

Product	Product Rate	Weeds Controlled/ Wood Size	Adjuvant
SELECT	6 to 8 fl. oz./A	Refer to Table—Regions A & B for Annual grasses controlled.	1 pt. Crop Oil Concentrate/A
	8 to 16 fl. oz./A	Refer to Tables—Region A/Region B for Perennial* grasses controlled.	
_____ plus _____ COBRA			
COBRA	12.5 fl. oz./A	Refer to COBRA label for weeds controlled.	

*If regrowth of Perennial grasses occurs, make a second application of SELECT 2 EC Herbicide using 6 or 8 fl. oz./acre timed to grass weed and sizes noted for Perennial grasses in Region A and Region B.

SEQUENTIAL APPLICATION OF SELECT 2 EC HERBICIDE AND BROADLEAF HERBICIDES

Some reduction in grass control is possible when SELECT 2 EC Herbicide is sequentially applied immediately prior to or after an application of a broadleaf herbicide.

1. After an application of SELECT wait a minimum of 24 hours before applying a broadleaf herbicide.
2. In fields treated with a broadleaf herbicide, do not apply SELECT until grass plants are again developing new leaves (generally 7 days after the broadleaf herbicide application).

14

Table 1I. (continued)
SELECT 2 EC Herbicide & ORTHENE Tank Mix—Cotton and Soybeans—Rate and Time of Application Table.

Product	Product Rate	Weeds and Insects Controlled	Adjuvant
ORTHENE 75 S	.33 to 1.33 lb./A	Refer to ORTHENE 75 S label for rate and insects controlled in cotton and soybeans.	

*If regrowth of Perennial grasses occurs, make a second application of SELECT 2 EC Herbicide using 6 or 8 fl. oz./acre timed to grass weed and sizes noted for Perennial grasses in Region A and Region B.

BEST AVAILABLE COPY

17

Table 5M.
SELECT 2 EC Herbicide & REFLEX® Tank Mix—Soybeans—Rate and Time of Application Table.

Product	Product Rate	Weeds Controlled/ Wood Size	Adjuvant
SELECT	6 to 8 fl. oz./A	Refer to Table—Regions A & B for Annual grasses controlled.	1 pt. Crop Oil Concentrate/A
	8 to 16 fl. oz./A	Refer to Tables—Region A/Region B for Perennial* grasses controlled.	
_____ plus _____ REFLEX			
REFLEX	.75 to 1.5 pts./A by region	Refer to REFLEX label for weeds controlled by region.	

*If regrowth of Perennial grasses occurs, make a second application of SELECT 2 EC Herbicide using 6 or 8 fl. oz./acre timed to grass weed and sizes noted for Perennial grasses in Region A and Region B.

SEQUENTIAL APPLICATION OF SELECT 2 EC HERBICIDE AND BROADLEAF HERBICIDES

Some reduction in grass control is possible when SELECT 2 EC Herbicide is sequentially applied immediately prior to or after an application of a broadleaf herbicide.

1. After an application of SELECT wait a minimum of 24 hours before applying a broadleaf herbicide.
2. In fields treated with a broadleaf herbicide, do not apply SELECT until grass plants are again developing new leaves (generally 7 days after the broadleaf herbicide application).

15

RESTRICTIONS AND LIMITATIONS

Do not graze treated fields or feed treated forage or hay to livestock. Do not apply if rain is expected within 1 hour of application as control may be unsatisfactory.

Do not apply a postemergence broadleaf herbicide within one day following application of SELECT or reduced grass control may result.

Do not apply SELECT within 60 days of harvest.

Do not apply more than 32.0 fl. oz. (0.5 lb. ai) per acre per season.

Do not apply under conditions of stress.

Grass crops such as corn, sorghum, wheat, rice, etc. are highly sensitive to SELECT.

Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotational and other crop restrictions.

Tank mixes of SELECT 2 EC Herbicide and broadleaf herbicides may result in reduced grass control. If grass regrowth occurs an additional application of SELECT 2 EC Herbicide may be necessary.

Do not tank mix with SCEPTER® Herbicide.

AVOID SPRAY DRIFT

Do not allow spray from ground or aerial equipment to drift onto adjacent land or crops. When drift may be a problem, do everything possible to reduce spray drift, including:

- Do not spray if wind speeds are or become excessive. Do not spray if wind speed is 10 MPH or greater. If sensitive crops or plants are downwind, extreme caution must be used under all conditions. Do not spray if winds are gusty.
- Use extreme caution when conditions are favorable for drift (high temperatures, drought, low relative humidity), especially when sensitive plants are located nearby.
- Do not apply when a temperature inversion exists. If inversion

18

5045

conditions are suspected, consult with local weather services before making an application.

- Further reductions in drift can be obtained by:

1. Using large droplet size sprays. Do not use nozzles that produce small droplets. Orient nozzles downward and slightly backward as needed to reduce drift for ground applications.
2. Orienting nozzles straight back with the windstream, using straight stream or cones for aerial applications. Use the lowest number of nozzles practical with the largest possible orifice size to obtain the minimum 3 GPA volume. Application height and boom length should be set according to manufacturer's instructions to minimize drift.
3. Increasing the volume of spray mixture (for example a minimum of 10 GPA for ground applications) by using higher flow rate nozzles. Using lower pressure with the appropriate nozzle to obtain higher volumes will also reduce drift.
4. Apply as close to target plants as practical while maintaining a good spray pattern for adequate coverage.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Open dumping is prohibited.

STORAGE

Keep pesticide in original container.

Do not put concentrate or dilute into food or drink containers.

Store in cool, dry place.

Do not store diluted spray.

Emergency Response: For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-457-2022, or Chemtrec 1-800-424-9300.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Do not reuse container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT INFORMATION. READ BEFORE USING PRODUCT.

LIMITED WARRANTY: Valent U.S.A. Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for use under average conditions when used strictly in accordance with the directions on the labeling. Valent does not make nor authorizes any agent or representative to make, any other warranty, guarantee or representation, express or implied, concerning this product. Specifically, **NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE IS MADE.**

2. Critical and unforeseeable factors beyond Valent's control prevent it from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants and crops to which product is applied, lack of complete control over the handling and application of this product, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit under average conditions for the uses stated on the labeling and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Valent under 1 above) resulting from handling, storage, and use of the product.

3. Precautions stated on the labeling should be followed to avoid hazardous exposure to the product.

BASAGRAN® - Reg. TM of BASF AG.

BLAZER® - Reg. TM of BASF AG.

CLASSIC® - Reg. TM of E.I. du Pont de Nemours & Co. Inc. for chlorimuron ethyl herbicide.

ORTHENE® - Reg. TM of Chevron Chemical Company for acephate insecticide.

REFLEX® - Reg. TM of ICI Americas, Inc. for fomesafen herbicide.

SCEPTER® - Reg. TM of American Cyanamid Co. for imazaquin herbicide.

Manufactured for

Valent U.S.A. Corporation © 1992

Walnut Creek, CA 94596-8025

R Form 383-B Made in U.S.A.

EPA Reg. No. 59639-3

BEST AVAILABLE COPY

