

Supplemental Label



Valent U.S.A. Corporation
1333 N. California Blvd., Ste. 600
Walnut Creek, CA 94596-8025

SELECT® 2 EC HERBICIDE

TOMATOES, ALFALFA, DRY BEANS AND PEANUTS
EPA Reg. No. 59639-03

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

ADDITION OF ADJUVANT OR CROP OIL CONCENTRATE

Tomatoes, Alfalfa, Dry Beans,
Peanuts

Always use a crop oil concentrate containing at least
15% emulsifier at 1% v/v in the finished spray
volume.

RESTRICTIONS AND LIMITATIONS

TOMATOES

Do not apply SELECT 2 EC Herbicide within 20 days of tomato harvest.

ALFALFA

SELECT 2 EC Herbicide may be applied to seedling or established alfalfa (including sainfoin, holy clover, and birdsfoot trefoil) grown for seed, hay, silage, green chop, or direct grazing.

Do not apply SELECT 2 EC Herbicide within 15 days of grazing, feeding, or harvesting (cutting) alfalfa for forage or hay.

Do not apply SELECT 2 EC Herbicide and 2,4-DB as a tank mix unless the 60 day feeding, grazing, and harvesting restriction on the 2,4-DB label can be observed.

Do not plant rotational crops until 30 days after application of SELECT 2 EC Herbicide.

DRY BEANS

Do not apply SELECT 2 EC Herbicide within 30 days of dry bean harvest (cutting or pulling plants from ground).


PEANUTS

Do not apply SELECT 2 EC Herbicide within 40 days of peanut harvest (cutting vines or digging peanuts from ground).

DIRECTIONS FOR USE IN TOMATOES, ALFALFA, DRY BEANS AND PEANUTS

RECOMMENDATIONS FOR ANNUAL GRASSES

- Apply only to actively growing grasses at recommended weed heights.
- Apply when the first grass weed species in a mixed grass weed population reaches the recommended growth stage for treatment.
- Use the high rate under heavy grass pressure and/or when grasses are at maximum height.
- For the control of annual grasses in seedling alfalfa use 6 to 8 fl. oz. per acre. For the control of annual grasses in established alfalfa use a minimum of 8 fl. oz. per acre.



GRASS SPECIES		WEED* HEIGHT INCHES	RATE FL. OZ./ ACRE	HIGH RATE
Barnyardgrass	Echinochloa crusgalli	2-8	6	8
Broadleaf Signalgrass	Brachiaria platyphylla	2-6	6	8
Brome				
California	Bromus carinatus	2-6	6	8
Cheatgrass	Bromus secalinus	2-6	6	8
Downy	Bromus tectorum	2-6	6	8
Ripgut	Bromus diandrus	2-6	6	8
Crabgrass				
Hairy	Digitaria adscendens	2-6**	6	8
Large	Digitaria sanguinalis	2-6**	6	8
Smooth	Digitaria ischaemum	2-6**	6	8
Southern	Digitaria ciliaris	2-6**	6	8
Crowfootgrass	Dactyloctenium aegyptium	2-6**	6	8
Fall Panicum	Panicum dichotomiflorum	2-8	6	8
Field Sandbur	Cenchrus incertus	2-6	6	8
Foxtail				
Giant	Setaria faberi	2-12	6	8
Green	Setaria viridis	2-8	6	8
Yellow	Setaria glauca	2-8	6	8
Foxtail Barley	Hordeum jubatum	2-6	6	8
Goosegrass	Eleusine indica	2-6**	6	8
Itchgrass	Rottboellia exaltata	2-6	6	8
Junglerice	Echinochloa colona	2-6	6	8
Lovegrass (Stinkgrass)	Eragrostis cilanensis	2-6	6	8
Red Rice	Oryza sativa	1-3	6	8
Rygrass				
Hardy	Lolium perenne	2-6	6	8
Italian	Lolium multiflorum	2-6	6	8
Seedling Johnsongrass	Sorghum halepense	4-10	6	8
Shattercane	Sorghum bicolor	6-18	6	8
Southwestern Cupgrass	Eriochloa gracilis	2-6	6	8
Sprangle top				
Amazon	Leptochloa panicoides	2-6	6	8
Bearded	Leptochloa fascicularis	2-6	6	8
Mexican	Leptochloa uninervia	2-6	6	8
Red	Leptochloa filiformis	2-6	6	8
Texas Panicum	Panicum texanum	2-6	6	8
Volunteer Cereals				
Barley	Hordeum vulgare	2-6	6	8
Oats	Avena sativa	2-6	6	8
Rye	Secale cereale	2-6	6	8
Wheat	Triticum aestivum	2-6	6	8
Volunteer Corn ²	Zea mays	4-12	4	6
Volunteer Corn (S.R.) ¹	Zea mays	4-12	8(suppression only)	
Volunteer Corn ²	Zea mays	12-24	6	8
Volunteer Grain Sorghum	Sorghum bicolor	8-12	6	8
Wild Oats	Avena fatua	2-6	6	8
Wild Proso Millet	Panicum miliaceum	2-10	6	8
Witchgrass	Panicum capillare	2-8	6	8
Woolly Cupgrass	Eriochloa villosa	2-8	6	8

* Generally occurs between 3-leaf stage and tillering.

** Length of lateral growth

1. Sethoxydim resistant volunteer corn

2. Includes Roundup Ready®, Liberty Link® and IMI-CORN™ volunteer corn

**RECOMMENDATIONS FOR ANNUAL BLUEGRASS CONTROL
WITH SELECT 2 EC HERBICIDE**

GRASS SPECIES	WEED STAGE	RATE FL. OZ./ ACRE	HIGH RATE
Annual Bluegrass (Poa annua)	to 4-Leaf	6*	16
<p>Apply under favorable soil moisture and humidity which exists within a few days after rainfall or within 7 days after irrigation. Grass needs to be actively growing at time of application(s).</p> <p>Apply at weed stage indicated on the label, as reduced control can be expected with more mature annual bluegrass.</p> <p>Use the high rate under heavy grass pressure and/or when annual bluegrass is more mature.</p> <p>Always add a crop oil concentrate containing at least 15% emulsifier at 1% v/v (but not less than 1 pint per acre) to the finished spray volume.</p>			

* Use a minimum of 8 fl. oz. per acre to control annual bluegrass in seedling and established alfalfa.

**TANK MIX APPLICATION OF SELECT 2 EC HERBICIDE AND
2,4-DB HERBICIDE FOR CONTROL OF GRASSES
AND BROADLEAF WEEDS IN ALFALFA**

A tank mix of SELECT 2 ECHerbicide plus 2,4-DB can be used to control grass and broadleaf weeds listed on the two product labels. Include a crop oil concentrate containing at least 15% emulsifier at 1% v/v in the finished spray. Follow rate and other recommendations on the individual herbicide labels when applying this tank mix.

NOTE: SELECT 2 EC Herbicide plus 2,4-DB may increase the severity of crop injury when tank mixed. Alfalfa plants will generally outgrow this temporary crop injury within a few weeks.

PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.
FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA-
REGISTERED LABEL.

Approved by EPA April _____ 1998
SU3100 A0.0 980403