


VALENT ORTHORIX Foliage Spray
EPA Reg. No. 59639-11-AA
Date Submitted: 11/30/89

Copy A

Wash contaminated clothing before reuse. Discard contaminated footwear.

Copy B

For skin contact, remove contaminated clothing and wash skin thoroughly with soap and water. See a doctor if irritation persists. If swallowed, give water or milk to drink and telephone for medical advice. DO NOT make person vomit unless directed to do so by medical personnel. If medical advice cannot be obtained, then take the person and product container to the nearest medical emergency treatment center or hospital. If inhaled, remove person from exposure area.





DANGER

STATEMENT OF PRACTICAL TREATMENT

Note to Physicians Emergency Information - call (415) 233 3127

ENVIRONMENTAL HAZARDS

PHYSICAL OR CHEMICAL HAZARDS

DIRECTIONS FOR USE

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH
PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH
APPLICABLE STATE AND FEDERAL REGULATIONS

REENTRY STATEMENT

Certain states may require more restrictive entry interviews.
Consult your State Department of Agriculture for further information.

Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas able to be treated.

This product contains the toxic inert ingredient methylene chloride.

KEEP OUT OF REACH OF CHILDREN

DANGER – PELIGRO

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

NET CONTENTS
28 GALLONS

Manufactured for
Valent U.S.A. Corporation • 1989
Walnut Creek CA 94596 8025
Product 1872 Made in U.S.A.
EPA Reg. No. 59639 11 AA
R Form 264 A EPA Est. 48975 CA

[illegible]

DIRECTIONS

D. Do not apply this product through any type of irrigation system.

CANE BERRIES Red Berry Mite, Leaf Spot, Cane Blight—3 gal's per 100 gal's water. Apply as a foliar dormant spray.

GRAPES Powdery Mildew—1 pt. OPTICOR® Foliar Spray plus 4 oz. Ventrol® 500 per 100 gal's water.

PEARS: California and Northwest only: **POST HARVEST AND LATE FALL TREATMENTS**—Pear Bud Mites, Blister Mites—Immediately after harvest: **APR. 15** and **5/10**—**CUTHORIX** Fungus Spray for control of **fruit rot** for 100% coverage apply after over 90% of fruit is **Fall Spray**: September-October, just prior

Leaf Roll—Apply 2½ gals. of ETHIONERB foliage spray per acre as a sufficient spray for thorough coverage application. **DE-LAYED DORMANT SEASON TREATMENT**—Pear Leaf Blister Mites, Pear Psylla, Scale Insects, Aphid and Mite Eggs—Apply 2½ gals. of ETHIONERB foliage spray per acre combined with 3 to 5 gals. of KEEN-EE Foliar Feeder Insecticide to 2 gals. of VITCK Supreme Spray in each 100 gal. of water as a sufficient spray for thorough coverage application.

RED CLOVER AND ALFALFA Powdery Mildew—1 gal. = 100 to 200 sq. ft. of alfalfa or 100 to 200 sq. ft. of red clover.

ROSES Nursery Stock and Field Grown Ornamentals. Powdery Mildew, Rust, Black Spot, Red Spider Mites - can't get past 10% control with fungicides, insecticides and miticides. 100% control with low cost biologicals severe. Pink open flowers before spring sale and disease control.

STONE FRUITS, CHERRIES, PEACHES, NECTARINES, PRUNES AND PLUMS—Brown Rot, Leaf Spot—(See Cherry, p. 166) *See also* Apples, p. 161. Water-borne bacteria—beginning of week, *See* p. 161. *See also* Apples, p. 161. **Powdery Mildew**—(See p. 161) *See also* Apples, p. 161.

PEACHES Poudery Mildew

Do not apply under conditions involving possible drift to food
crops or other plantings that might be damaged or the crops
thereof rendered unfit for sale, use or consumption.