



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

Corrosive to eyes. Concentrate causes eye damage and skin irritation. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid breathing spray mist.

STATEMENT OF PRACTICAL TREATMENT

In case of eye contact, immediately flush eyes with fresh water for 15 minutes and see a doctor. Wash skin thoroughly with soap and water. If swallowed, give large quantities of water, induce vomiting and see a doctor immediately.

Note to Physicians: Emergency Information - call (415) 233-3737.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment wastewater.

PHYSICAL OR CHEMICAL HAZARDS

MAY GIVE OFF HIGHLY TOXIC AND EXTREMELY FLAMMABLE HYDROGEN SULFIDE GAS IF MIXED WITH ACIDS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

REENTRY STATEMENT

Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift.

Do not enter treated areas without protective clothing until sprays have dried.

Certain states may require more restrictive reentry intervals; consult your State Department of Agriculture for further information.

Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated.

ORTHO ORTHORIX® Foliage Spray

(FUNGICIDE-MITICIDE)

Active Ingredient	By Wt.
Calcium Polyvalent Sulfide	27.5%
Inert Ingredients	72.5%
Density Base at 68° F. - 29.9. Contains calcium and sulfur expressed as gypsum—2.8 lbs./gal. Total sulfur 2.29 lbs./gal. Other combined sulfur 1.8 lbs./gal. 10.5 lbs. Per Gal.	

This product contains the toxic inert ingredient Methylene Chloride.

KEEP OUT OF REACH OF CHILDREN DANGER-PELIGRO

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.

PRECAUCION AL USUARIO: Si usted no lee
ingles, no use este producto hasta que la
etiqueta le haya sido traducida ampliamente.

NET CONTENTS 28 GALLONS

Chevron Chemical Company © 1988

Ortho Agricultural Chemicals Division
San Francisco CA 94120-7144
Product 1872 Made in U.S.A.

Form 2488-62

EPA Est. 48976-CA-01

EPA Reg. No. 239-709-AA

ORTHORIX Foliage Spray is a balanced fungicide-insecticide composition containing special effective wetting agents. It is for Growing Season Spraying. For best results spray to cover all exposed plant parts and repeat as necessary.

Do not use on crops as they are susceptible to sulfur injury. ORTHORIX Foliage Spray may burn foliage when temperature is high (85°F). Do not make applications at such times. When applied to crops with foliage, let 30 days elapse between treatments with ORTHORIX Foliage Spray and treatments with an oil spray. If crust occurs on surface of ORTHORIX Foliage Spray, break through. Crystallization does not impair product quality.

DIRECTIONS

Do not apply this product through any type of irrigation system.

APPLES AND PEARS: Scab and Mildew Control—Use 2 qts. in 100 gals. of water. Apply in Prepink, Pink and Calyx periods. Powdery Mildew: Continued spraying is required during growing season. Do not use ORTHORIX Foliage Spray on D'Anjou pears after the calyx period.

CANEBERRIES: Red Berry Mite, Leaf Spot, Cone Blight—3 gals. per 100 gals. water. Apply as a delayed dormant spray.

GRAPES: Powdery Mildew—1 pt. ORTHORIX Foliage Spray plus 4 lbs. Wettable Sulfur in 100 gals. water.

PEARS (California and Northwest only): POST HARVEST AND LATE FALL TREATMENTS—Pear Bud Mites, Blister Mites—Immediately after harvest: Apply 2½ gals. ORTHORIX Foliage Spray per acre in sufficient water for a thorough coverage application on average size trees. Fall Spray (September-October, just prior to Leaf Fall)—Apply 2½ gals. ORTHORIX Foliage Spray per acre. Use sufficient spray for a thorough coverage application. DELAYED DORMANT SEASON TREATMENT—Pear Leaf Blister Mites, Pear Psylla, Scale Insects, Aphid and Mite Eggs—Apply 2½ gals. ORTHORIX Foliage Spray per acre combined with 3 to 5 gals. KLEENUP Flowable Emulsion or 1½ to 2 gals. VOLCK Supreme Spray in each 100 gals. water. Use sufficient spray for thorough coverage application.

RED CLOVER AND ALFALFA: Powdery Mildew—1 gal. in 50 to 100 gals. water per acre at bud stage.

ROSES (Nursery, Stage and Field Grown Ornamentals): Powdery Mildew, Rust, Black Spot, Red Spider Mites—Use 1½ pts. per 100 gals. of water on tender new growth and 1½ qts. in 100 gals. of water if mildew becomes severe. Pick open flowers before spraying to avoid discoloration.

STONE FRUITS (CHERRIES, PEACHES, NECTARINES, PRUNES AND PLUMS): Brown Rot, Leaf Spot—Use 2 qts. per 100 gals. water. Apply 3 to 5 weeks before harvest, repeating at weekly intervals until just before picking. Powdery Mildew—2 qts. per 100 gals. water, when disease appears, or use ½ pt. with 4 lbs. wettable sulfur. PEACHES: Powdery Mildew—1 pt. plus 5 to 10 lbs. wettable sulfur per 100 gals. of water as a dilute application or 2 to 4 pts. plus 15 to 32 lbs. wettable sulfur per acre in 20 to 120 gals. per acre as a concentrate application. Do not permit workers to enter any part of such treated field to engage in any activity that may involve substantial body contact with the foliage within 1 day of application.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

PROHIBITIVE

Do not contaminate cleaning of equipment. Open dumping.

STORAGE

Keep pesticide. Do not put container in material, call d

PESTICIDE

This product is a pesticide. Use according to label. Environmental representative of t

CONTAINER

Completely empty. Do not use for other purposes. Do not dump. Do not use for other purposes. Do not use for other purposes.

CONDITIONS OF SALE: (a) warrants the description on the hereon Chevron representative to MERCHANTABILITY, concerning

2 Critical and not prevent the use of chemical damage to plant, lack of complete plants or crops, reasonably fit for directions are followed, all risks and liabilities resulting

CONFIDENTIAL
PROPERTY OF
CHEVRON CHEMICAL COMPANY
UNTIL ACCEPTED BY EPA