

PM


19

59639-20

page 19 10



**VOLCK® Supreme Spray
(INSECTICIDE-MITICIDE)
A Superior Oil Spray**



Active Ingredient	By Wt.
Petroleum Oil	97.4%
Inert Ingredients	2.6%
Petroleum Oil Min. Unsulfonated Residue - 99%	
Petroleum Oil Classification - Unclassified.	


For Western Agriculture

KEEP OUT OF REACH OF CHILDREN

CAUTION

**SEE BELOW FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

**NET CONTENTS____
GALLONS_____**



2910

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes, or clothing. If contact occurs, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush skin with plenty of water. Get medical attention if irritation persists.
If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If swallowed: **DO NOT INDUCE VOMITING.** Contact Physician immediately. Aspiration may be a hazard.

Note to Physicians: Emergency Information - call 1-800-892-0099.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long-sleeved shirt and long pants, and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY REQUIREMENTS

Users should:


- * Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- * Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS

Do not weld, heat or drill this container. Residues may ignite with explosive violence if heated sufficiently.



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, and shoes plus socks.

DIRECTIONS

VOLCK Supreme Spray is an especially refined Phytonomic Oil Contact Insecticide. It is for use on plants in foliage or the dormant stage.

When using this product in combination with other pesticides as directed on the label, observe appropriate cautions, limitations for the product involved.

For foliage applications, the spray tank must be free of any sulfur residue before using VOLCK Supreme Spray. Do not combine sulfur sprays with VOLCK Supreme Spray on foliage and do not follow a previous application of sulfur or apply sulfur following VOLCK Supreme Spray until sufficient time has elapsed for safety (at least 30 days). If in doubt, spray a portion of one tree previously treated with sulfur, using 1 qt. to 25 gals. water. If sufficient time has not elapsed,

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

Do not apply this product through any type of irrigation system.

Apply quantity of product shown for each listed crop for control of listed insects in sufficient water to make a minimum of 20 gals. dilute spray per acre by air application. Follow timing shown for each crop as shown on container label. Use only as emergency application when soil conditions do not permit regular ground application.

GENERAL NOTES: Observe on all crops and uses. - Oil effectiveness in dormant and delayed dormant applications is determined by thoroughness of coverage. Concentrate applications may reduce coverage and effectiveness. - Oil applications made to trees that are weak or suffering from moisture stress or when slow drying of spray occurs may cause bud, twig or leaf injury.

Check this product for proper emulsification by mixing a small quantity of it with water in a small container before filling the entire spray tank. Correct emulsification is indicated when a uniform mixture, with a blue color, results on shaking. If this mixture is not a milky emulsion after shaking, do not use the remaining product. With agitator running, start filling spray tank with water. When combining with wettable powder insecticide or fungicide products, add these first when tank is 1/3 filled. Add VOLCK Supreme Spray when tank is about 1/2 full and while there is sufficient agitation of the spray to ensure complete emulsification.

age on other plant consumption.

...to improve coverage for maximum ...
 ...if San Jose scale is moderate to severe problem use dilute ...
 ...application may delay bud development. The following ...
 ...recommendations are ...

FOUR APPLICATIONS, DILUTE SPRAYS

CROP	PESTS	VOLCK PER 100 GALS WATER	FURTHER USE INSTRUCTIONS
ALMONDS APRICOTS CHERRIES NECTARINES PEACHES PLUMS PRUNES	Brown Acanth Scale San Jose Scale	1 gal	For summer stages of Brown Acanth and San Jose Scale.
	Red Spider Mite	8 to 4 gal	For summer use on Red Spider Mite.
APPLES	Red Scale	1 to 1 1/2 gal	For summer use.
APPLES PEARS	Red Spider Mite	8 to 4 gal	For summer use.
	San Jose Scale Russet Pear Scale	1 to 1 1/2 gal	Apply VOLCK Supreme Spray to summer forms of San Jose and Russet Pear Scale.
GRAPES	Nursery	See NOTE	For (Post Harvest) Fall spray use only. NOTE: Use 4 to 6 gal. in 200 to 300 gal. of water per acre.
GRAPEFRUIT LEMONS	Black Scale Purple Scale Red Scale Yellow Scale Red Spider Mite	1 to 1 1/4 gal	For best results, apply VOLCK Supreme Spray during winter and spring months when Red Scale are more easily controlled and these generally are less reactive to oil sprays. Do not use within 8 weeks of harvest on lemons.
NAVEL ORANGES	Black Scale Purple Scale Red Scale Yellow Scale Red Spider Mite	5 gal	Do not use on Navel oranges after September 1.
OLIVES	Black Scale Olive Scale Olive Scale (Parasitic Scale) Spider Mite	1 1/2 gal	Apply VOLCK Supreme Spray in the prebloom or in postbloom up to budburst stages.
OLIVES (CALIFORNIA ONLY)	Black Scale	1 to 1 1/2 gal	Apply 400 to 800 gal. dilution per acre. Make applications between July 1 and August 31. Higher concentrations of VOLCK Supreme Spray are not recommended above that cause phytotoxicity.

2013年12月9日 星期日

SYMBOL	PLANT PARTS	PLANT SPECIES	ADDITIONAL INSTRUCTIONS
PLUM	PLUM PRUN	PLUM PRUN	PLUM PRUN
	PLUM PRUN	PLUM PRUN	PLUM PRUN
PLUM	PLUM PRUN	PLUM PRUN	PLUM PRUN
WALNUTS	WALNUT PRUN	WALNUT PRUN	WALNUT PRUN
ORNAMENTAL	ORNAMENTAL PRUN	ORNAMENTAL PRUN	ORNAMENTAL PRUN
SHADE TREES	SHADE TREES PRUN	SHADE TREES PRUN	SHADE TREES PRUN
CONIFERS	CONIFERS PRUN	CONIFERS PRUN	CONIFERS PRUN
FLOWERING PEACH	FLOWERING PEACH PRUN	FLOWERING PEACH PRUN	FLOWERING PEACH PRUN
FLOWERING CHERRY	FLOWERING CHERRY PRUN	FLOWERING CHERRY PRUN	FLOWERING CHERRY PRUN
APPLE	APPLE PRUN	APPLE PRUN	APPLE PRUN
MAGNOLIA	MAGNOLIA PRUN	MAGNOLIA PRUN	MAGNOLIA PRUN
MAPLE	MAPLE PRUN	MAPLE PRUN	MAPLE PRUN
DOGWOOD	DOGWOOD PRUN	DOGWOOD PRUN	DOGWOOD PRUN
OAK	OAK PRUN	OAK PRUN	OAK PRUN
PLUM	PLUM PRUN	PLUM PRUN	PLUM PRUN
BIRCH	BIRCH PRUN	BIRCH PRUN	BIRCH PRUN

BEST AVAILABLE COPY

[illegible]

DORMANT OR DELAYED DORMANT - DILUTE SPRAYS

CHOP	PESTS	VOLCK PER 100 GALS. WATER	FURTHER USE INSTRUCTIONS
ALMONDS APRICOTS CHERRIES NECTARINES PEACHES PLUMS PRUNES (see NOTE)	Brown Ackerhorm Scale Black Scale San Jose Scale Olive Scale (Palmolive Scale) Olivehorm Scale Fruitworm Scale Sawfly Pear Scale European Fruit Lethargy Peach Scale Soft Scale Scale Garden other Scale Insects Scale of Ackerhorm Red Spider Mite	1 1/2 to 2 gals.	For dormant or delayed dormant use only. NOTE: For plums and prunes, spray only during the delayed dormant period. Of sensitive varieties (such as, but not limited to, Moyer or Queen Ann) may develop dead limbs or cankers, which will reduce yield.
APPLES	Pink Mite	1 1/2 to 2 gals.	For delayed dormant use only.
APPLES PEARS	San Jose Scale Sawfly Pear Scale Scale of Ackerhorm Red Spider Mite	1 1/2 to 2 gals.	For dormant or delayed dormant use only.
PEARS	Pink Mite	2 gals.	For dormant or delayed dormant use only.
CHERRIES	Black Scale	See NOTE	For dormant use only. NOTE: Use 4 to 6 gals. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 gals. of water per acre.
PEARS	Pink Mite	1 to 1 1/2 gals.	For dormant or delayed dormant use only.
	Black Scale	1 to 1 1/2 gals.	For dormant or delayed dormant use only.

BEST AVAILABLE COPY

[illegible]

SEED CROP USES

CROP	PESTS	VOLCK PER ACRE	FURTHER USE INSTRUCTIONS
ALFALFA (SEED ALFALFA ONLY)	SPIDER MITE	1 qt.	For use on seed alfalfa only. Apply in 20 to 30 gals. of water by ground application or 10 to 15 gals. of water by air with recommended rate of Kathon-A EO.

BEST AVAILABLE COPY

10 of 10

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment.
Open dumping is prohibited.

STORAGE

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Protect from water contamination during storage and transfer. Contamination with water may make this product unfit for its intended use. Always check emulsification (see MIXING DIRECTIONS) before use. For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-892-0099 or Chemtrec 1-800-424-9300.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

IMPORTANT INFORMATION. READ BEFORE USING PRODUCT.

LIMITED WARRANTY: 1. Valent U.S.A. Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for use under average conditions when used strictly in accordance with the directions on the labeling. Valent does not make nor authorizes any agent or representative to make, any other warranty, guarantee or representation, express or implied, concerning this product.

Specifically, NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE IS MADE.

2. Critical and unforeseeable factors beyond Valent's control prevent it from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants and crops to which product is applied, lack of complete control over the handling and application of this product, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit under average conditions for the uses stated on the labeling and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Valent under 1 above) resulting from handling storage, and use of the product.

3. Precautions stated on the labeling should be followed to avoid hazardous exposure to the product.

Copyright© 1994 by Valent U.S.A. Corporation

VOLCK® Reg. TM of Monsanto Company.

KELTHANE® - Reg. TM of Rohm & Haas Company.

Manufactured for

Valent U.S.A. Corporation

Walnut Creek CA 94596-8025

R12100B1.1940523 Made in U.S.A.

EPA Reg. No. 59639-20 EPA Est. _____

