


Reg # 59639-26

PM-14

ORTHENE 75 S Soluble Powder
EPA Reg. No. 59639-26
Submitted: September 5, 1990

Copy A

For use on agricultural crops, commercially grown ornamentals, commercial sod farms, and in greenhouses, reentry into treated areas is prohibited for 24 hours after the end of application unless the protective clothing specified elsewhere in this label is worn. For other uses, including golf courses and other non-agricultural areas, do not enter treated areas without protective clothing until sprays have dried.



(INSECTICIDE)

CAUTION
SEE BELOW FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

1. The ingredient
"Acetate" is a long name
of a substance in the
ingredient
"with the Acetate"
is a 3588

8. 4-

Manufactured for
Valent U.S.A. Corporation
Agricultural Division
P.O. Box 1000
Vallejo, California 94588

ORTHENE KILLS FIRE ANTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

CAUTION: 1. Soap-Detergents cause eye irritation. Avoid contact with eyes. 2. Avoid breathing dust or spray. 3. Wash hands thoroughly after handling. 4. Do not use if contents have been in contact with treated areas with 20 days after use.

STATEMENT OF PRACTICAL TREATMENT

STATEMENT OF PHYSICAL TREATMENT:
 1. Wash eyes 1 or 2 times with water for 1 min. and induce vomiting by touching the back of the throat with a finger.
 2. If possible contact physician for advice. Wash eyes or emergency center before inducing vomiting.
 3. Induce vomiting by drinking 1 to 2 cups of water or 1 to 2 cups of 1% solution of table salt.
 4. If the nearest emergency treatment center is at least 15 min. away contact wash eyes with fresh water for 15 minutes.
 5. If vision continues poor a doctor for same contact wash skin with plenty of soap and water if inhaled remove person from exposure area.

toxic to Physicians (emergency information: call 800 451 7022). Aceprate is a cholinesterase inhibitor. If signs of cholinesterase inhibition appear, atropine is indicated. 2 PAM may also be used in conjunction with atropine but only in mild to moderate cases.

WOMEN'S SAFETY

APPLICATION

[illegible][illegible]

ENVIRONMENTAL HAZARDS

To prevent damage to crops, Apple Growers may adversely affect birds, a large group of farmstead species, by not applying products to water or wetlands. However, such an application is not appropriate.

To protect the environment, the label requires that the product be applied to crops or weeds if the crops or weeds are not being treated and the crops or weeds are not being treated.

DIRECTIONS FOR USE

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS

WARRANT STATEMENT

Keep all unprotected persons out of operating areas 31-61 to where there may be 31-7.
~~Do not allow unprotected persons to enter the area.~~
 Certain states may require more restrictive reentry intervals. Consult your State Department of Agriculture for further information.

Information
 Instruction or training regarding use of protective clothing and all dental exposure must be given to workers who are expected to be in treated areas or areas about to be treated

DIRECTIONS:

Apply to Cushman's Law: 1. Sprinkler irrigation systems. Do not apply to chemigation or any other crop or this crop using any other type of irrigation system.

[illegible]

COTTON TACKLING MIXES				
Crops	Insecticides	Average GUTHRIE 75 S + PYRINEX 2.4 EC Per Acre	Further Use Instructions	Days to Harvest
Cotton (CA AZ)	Armigera Bombyx Lobesia botrana cotton Aphid caterpillar leafhopper grasshoppers Pink Bollworm Pearl Borer cactus moth etc. Sawfly Colorado potato beetle Thrips Tobacco budworm stinkbug	3 to 1 1/2 lbs 1 to 2 pts	Do not feed gun trash or treated forage to livestock. Do not allow livestock to graze on treated areas.	
Cotton (Except CA AZ)	Tobacco Budworm	2 3/4 - 2 2/3 lb ac	Began laying eggs on cotton or musclic poplar and apply as needed to maintain control. Apply in 3 to 5 gals. of water with aerial treatment or in 10 to 25 gals. of water with ground treatment. Do not feed gun trash or livestock or allow animals to graze treated areas.	21

BEST AVAILABLE COPY

EDITOR BOB HOPPER ON TREATMENT
EXCERPT AT PA AND THE BLACK LAMBS OF TITAN

PELLE GRASS

1999年12月

SCIENTIFIC INVESTIGATION APPLICABLE TO CHARGE ONE

performance by removing the strongest test the tests of effectiveness

BEST AVAILABLE COPY

VEGETABLE COMPS

		Amount Required 75 lb Per Acre		Further Use Instructions		Days to Harvest	
Crops		Insects					
Beans and Peas	and Peas	Colorado potato beetles		Apply 1/2 water at minimum of 3 gal. spray per acre by air or 100 gal. spray per acre by ground		4-5 weeks	
		Colorado potato beetle		Apply when eggs or insects first appear. Repeat at 7-10 day intervals as necessary. Severe insect infestations use higher rates.		4-5 weeks	
Beans and Peas	and Peas	Colorado potato beetle		Do not feed treated plants to livestock			
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					
Beans and Peas	and Peas	Colorado potato beetle					

PASTURE, RD 1, BELAND WASTELAND

Crop	Insects	Amount OUTLINE 75 S Per Acre	Further Use Instructions	Days to Harvest
Pasture Rangeland Woods	Assheepers Midges Flies Black Grass Gnats	0.5 lb. 0.04 to 0.15 lb. 1 lb. 1 pint/l.	Apply at water at a maximum of - 1 qt. spray per acre or 10 to 20 gal. spray per acre by ground. Use the higher rate in thicker vegetation or on larger insects. An approved "hot target" additive may be added to aid in concentrating drift or reducing evaporation of spray gases. Do not make more than one application per season. Do not apply when lactating dairy cattle are present. Do not slaughter or feed treated hay to slaughtering dairy cattle within 21 days after application. Remove meat animals from treated areas or pastures as before a slaughter. They were present at application or grazed treated areas within 21 days after application.	14-28
-parted fire A-1's	Banded Thrips— Dry Blattellid Every distribute to 2 seasons over the mound Branch Blattellid Mix 1 oz. in 5 gals. of water. Apply 1 gal. of mix to each mound three by sprinkling the mound until it is wet and break it four or five seasons once around the mound.	For best results apply the mixture in the early morning or late afternoon when the ants are most active. Applications made under prolonged hot and dry conditions may be ineffective due to the drying of the ants deep within the nest. Grass in treated area may be ruined. Treat maximum of 13 mounds per acre. Do not treat mounds more than once per season. Do not apply when lactating dairy cattle are present. Do not pasture or feed hay to slaughtering dairy cattle within 21 days after application. Remove meat animals from treated areas at least 1 day before slaughter if they were present at application or grazed treated areas within 21 days after application.		

SPECIALTY USES

SPECIALTY CROPS				
Crops	Months	Average OFFSEASON PLS Per Acre	Further Use Instructions	
Tomatoes and peppers and cucumbers and cabbages and broccoli etc.	Greenhouse 7	10 75% above average	For early in the season application to greenhouses in red borders, terraces, roadside ditches, and between the dunes in 1 to 5 yard of water in or in 10 to 20 yard of water with ground elevation. Use the higher returns when spraying heavier crops. Do not graze or feed vegetation cut from treated areas.	
Crops	Months	Average OFFSEASON PLS Per Acre	Further Use Instructions	Best to Harvest
Tomatoes and peppers and cucumbers and cabbages and broccoli etc.	Greenhouse 7	10 75% above average	For best results apply the material to greenhouses in red borders, terraces, roadside ditches, and between the dunes in 1 to 5 yard of water in or in 10 to 20 yard of water with ground elevation. Use the higher returns when spraying heavier crops. Do not graze or feed vegetation cut from treated areas.	

Apply 1 gal before emergence or
they were present at application
or ground treated areas within 21
days after application

SPECIALTY USES

Crops	Insects	Amount DITHANE 75 S Per Acre	Further Use Instructions
Non-Crop Areas - tree borders - fence rows - orchards - ditches - and barrow - pits	Grasshoppers	1 lb 25-50 active ingredients	For early to mid season application to grasshoppers in tree borders, fence rows, - ditches, ditches and barrow pits. Apply 1 to 5 gal. of water by air or - 10 to 20 gal. of water with ground - equipment. Use the higher volumes when - spraying heavier foliage. Do not graze or - feed vegetation cut from treated areas.

Crops	Insects	Amount DITHANE 75 S Per Acre	Further Use Instructions	Days to Harvest
Non-Crop Areas	Imported Fire Ants	Amount Dithane 75 S Dry Material Every distribute 1 to 2 teaspoons over the mound Branch Material Use 1 qt. in 5 gal. of water Apply 1 gal. of mix to each mound area by sprinkling the mound until it is wet and wait a few 100 feet diameter circle around the mound	For best results apply the material in the early morning or late afternoon when the ants are most active. Applications made under prolonged hot and dry C. seasons may be ineffective due to the location of the ants deep within the nest. Grass in treated area may be injured. Do not treat mound more than once per season.	

Plants	Insects	Amount DITHANE 75 S Per 100 Gal.	Time of Application	Recommended Application
Container Grown Nursery Stock Arbor vitae Azalea Camelia Rhododen- dron Roses Viburnum Wein	Black Vine Weevil Sawflies Root Weevil	1 lb (16 oz.)	Application should be made by mid September for green house stock and by mid October for out door stock. Consult your local county extension agent for information on the dormant stage and control of root weevils on ornamentals.	Apply the specified amount of DITHANE 75 S Soluble Powder per 100 gal. of solution as to thoroughly drench the root system.

SMALL SPRAY EQUIPMENT

Preparation of DITHANE Spray Mix using a 16 oz. Measuring Cup

Label Recommendation	Equivalent DITHANE in 20 Gal. Water*	Equivalent DITHANE in 1 Gal. Water (per Gal.)
DITHANE 75 S Sol. Powder		
1.3 lb	5.2 oz	2
2.3 lb	9.2 oz	4
1.3 lb	0.2	

*Use 16 oz. measuring cup

TREES AND SHRUBS

Plants	Insects	Amount DITHANE 75 S Per 100 Gal.	Time of Application	Recommended Application
Trees and Shrubs - recept - flowering - crab - apple - tree - Drzewi	Aphids Bagworms Birch Leafminer tent Caterpillar Lace Bugs Leafrollers	1.3 lb (5.3 oz.)	As the insects begin to appear	Apply the specified amount of DITHANE 75 S Soluble Powder in 100 gal. water with a hydraulic sprayer. The addition of a suitable sticker improves control of Gypsy Moth larvae.
	Douglas Fir Tussock Moth Larvae Gypsy Moth Larvae Weevils	2.3 lb (18.5 oz.)	As the insects begin to appear	*Must be applied before bud break. Adjust rates to 1.3 lb. per 100 gal. water for Gypsy Moth control and 18 lb. per 100 gal. water for tent Caterpillar control.
	Scales (Crowners)	2.3 lb (18.5 oz.)	As crowners begin to appear. Repeat applications at 2 week intervals may be necessary where there is continuous crown production.	Do not apply to Hackberry, Elm, of Green, Camellia, Lombardy Poplar and Viburnum suspension.
Trees and Shrubs	Pondosa Pine Needle Miner	2.3 lb (18.5 oz.)	Time of application is important. Consult your Farm Advisor or County Extension Agent.	Nursery crop. Before treating large plantings spray only a few plants and observe the results for phyllosticta.
	Grasshoppers	2.3 lb (18.5 oz.)	As the grasshoppers begin to appear	Apply the specified amount of DITHANE 75 S Soluble Powder in 100 gal. water with a hydraulic sprayer as a full coverage spray.
	Cantharis Dormant Caterpillars (Spring and Fall)	1.3 lb 2.3 lb (5.3 lb 18.5 lb)	As the insects begin to appear. Use the higher amount when the larger larvae are present.	Apply the specified amount of DITHANE 75 S Soluble Powder in 100 gal. water with a hydraulic sprayer.
	Nannucal Pine Tip Moth Larvae	1.3 lb (5.3 oz.)	Time of application is important. Consult your Farm Advisor or County Extension Agent. Repeat applications can be required for control of some populations.	
	Red Weevil Adults	1.3 lb (5.3 oz.)	Apply when first feeding damage occurs. Repeat applications at four week intervals until the last weevil host may be necessary for complete damage protection.	
	Ban Elder Bugs Sawflies Barkbeetles Leafhoppers	1.3 lb (5.3 oz.)	As the insects begin to appear	Apply the specified amount of DITHANE 75 S Soluble Powder in 100 gal. water with a hydraulic sprayer as a full coverage spray.
	Japanese Beetle	1.3 lb (5.3 oz.)	As the Japanese Beetles begin to appear. Repeat applications at 2 week intervals may be necessary.	
	Em Leaf Beetle Larvae	1.3 lb (5.3 oz.)	As the larvae begin to appear. DITHANE will not provide Em Leaf Beetle eggs from hatching.	Apply the specified amount of DITHANE 75 S Soluble Powder in 100 gal. water with a hydraulic sprayer as a full coverage spray.
Douglas Fir Christmas Trees	Douglas Fir Needle Midge	1.3 lb (5.3 oz.)	Apply often should be made no more than 2 weeks prior to bud burst. For additional pest management in greenhouse, consult your County Extension Service.	Apply the specified amount of DITHANE 75 S Soluble Powder in 100 gal. water with a hydraulic sprayer as a full coverage spray.
Flowering Cereals	Aphids Tent Caterpillars Leafrollers	1.3 lb (5.3 oz.)	As the insects begin to appear	Apply the specified amount of DITHANE 75 S Soluble Powder in 100 gal. water with a hydraulic sprayer. Do not apply more than 3 times in a growing season at 4 week intervals. Caution: Phytotoxicity has been reported on the following Cereals - sorghum, maize, sorghum, - Amaranthus, Pigeon Pea, Pigeon Pea, - Red Wing and Grain Clover.

BEST AVAILABLE COPY

COMMERCIAL TURFGRASS, LAWN, AND OTHER RECREATIONAL TURFGRASS AREAS
DO NOT ALLOW LIVESTOCK TO GRAZE TREATED AREAS. DO NOT FEED TREATED GRASS TO LIVESTOCK

Plants	Insects	Amount ORTHERE 75 S Per 1000 Sq. Ft.	Time of Application	Recommended Application
Turfgrass Recreational & Commercial Lawns	For Armyworms reared on A. m. m. m. Armyworms	1 to 12 oz. 1 to 2 1/2 lbs. per acre	As the insects appear. A repeat application at 2 week intervals may be necessary.	Apply the specified amount of ORTHERE 75 S Soluble Powder per 1000 sq. ft. Use a minimum of 5 gals. water per 1000 sq. ft. to obtain good coverage.
	Chinch Bugs	12 to 24 oz. 3 to 6 lbs. per acre	Apply as needed for control. Repeat each week (10 to 14 days).	Apply the specified amount of ORTHERE 75 S Soluble Powder per 1000 sq. ft. Use 1 to 15 gals. water per 1000 sq. ft. to obtain good coverage.
	Grass	12 to 24 oz.	Apply as needed for control. Repeat (10 to 14 days) only if needed after treatment.	
	For Weevils (Grasshopper)	1 to 1 lb. 1 to 2 1/2 lbs. per acre	As the weevils begin to appear. Use the higher amounts when quick knock down is needed or when heavy infestations Repeat application may be necessary. Do not repeat at more than 1 week intervals.	Apply the specified amount of ORTHERE 75 S Soluble Powder per 1000 sq. ft. Use 1 to 15 gals. water per 1000 sq. ft. to obtain good coverage.
	Grasshopper	1 oz. (2 1/2 lbs.) per acre	As the grasshoppers begin to appear. A repeat application at 1 week intervals may be necessary.	Apply the specified amount of ORTHERE 75 S Soluble Powder per 1000 sq. ft. Use 1 to 15 gals. water per 1000 sq. ft. to obtain good coverage.
	More Crickets Except CA	10 to 15 oz. (2 1/2 to 3 lbs.) per acre	As these crickets begin to appear. For knockdown of heavy infestations, more than one application may be required. Throughout the grow- ing season, for heavy infestations, use the higher dosage rate.	Apply the specified amount of ORTHERE 75 S Soluble Powder per 1000 sq. ft. Use 1 to 15 gals. water per 1000 sq. ft. to obtain good coverage.
	Grasshopper (Grasshopper)	1 oz. (1 1/2 lbs.) per acre	Apply when insects of their damage first appear. Repeat as necessary. Insects do not to be re- peated at more than 1 week intervals.	Apply the specified amount of ORTHERE 75 S Soluble Powder per 1000 sq. ft. Use 1 to 15 gals. water per 1000 sq. ft. to obtain good coverage. Do not mow turfgrass for at least 24 hours after application.

Plants	Insects	Amount ORTHERE 75 S Per 1000 Sq. Ft.	Time of Application	Recommended Application
Turfgrass	For Ants	1 to 2 oz. 1 to 2 lbs. per acre	For best results, apply the material in the early morning or late afternoon when the ants are most active. Applications made under prolonged hot and dry conditions may be ineffective due to the action of the ants deep within the nest.	Apply the specified amount of ORTHERE 75 S Soluble Powder as directed. Grass in treated areas may be killed. Do not treat more than once per season.

Plants	Insects	Amount ORTHERE 75 S Per 1000 Sq. Ft.	Time of Application	Recommended Application
Ornamentals	For Beetles Southern Armyworm Grasshopper Armyworm	10 to 15 oz. (2 1/2 to 3 lbs.) per acre	As the insects appear. Repeat at 2 week intervals as necessary.	Apply the specified amount of ORTHERE 75 S per 1000 sq. ft. Use a minimum of 15 gals. of water per 1000 sq. ft. to obtain good coverage.

OUTDOOR WARP AND PERIMETER SPRAY

Location	Insects	Amount ORTHERE 75 S Per Gallon	Time of Application	Recommended Application
Outdoor Warp and Perimeter Area	For Beetles	16 oz. (1 lb.)	Treat early in the day, as insects are generally less active during these hours.	Apply specified amount of ORTHERE 75 S Soluble Powder per each gallon of water used. Apply as a spot treatment to the most hot entrance and surrounding areas where the insects enter.
	For Beetles Ants Grasshopper Grasshopper Grasshopper	16 oz. (1 lb.)	As the insects appear.	Apply specified amount of ORTHERE 75 S Soluble Powder per each gallon of water used. Apply to a band of soil 6 to 10 feet adjacent to the structure and to a height of 2 to 3 feet on the foundation where pests may be active or may find entrance. Also apply as a residual spray or wash a paint brush to sur- face of buildings, window frames, shutters, eaves, etc. Apply to porch, garage, etc. around garage area and other areas where the pests congregate.

OUTDOOR FLORAL CROPS AND GREENHOUSE CROPS

Plants	Insects	Amount ORTHERE 75 S Per 100 Gals.	Time of Application	Recommended Application
Flowers Dahlias Siberian Lilacs Grapefruit Peach Apple Roses Sage Stachys Sage Roses Zinnia	For Beetles	2.3 lb. (10.5 oz.)	As insects begin to appear. Repeat application may be necessary.	Apply the specified amount of ORTHERE 75 S in 100 gals. water with a hydraulic sprayer as a full coverage spray. Repeat application may cause slight tip burn or marginal leaf necrosis on some varieties. Test on a few plants to determine varietal susceptibility.
Roses Blossom	For Beetles	1.1 lb. (5.25 oz.)	As the Japanese beetles begin to appear. Repeat application at 2 week intervals may be necessary.	

GREENHOUSE FLORAL AND POTTED PLANT CROPS

Plants	Insects	Amount ORTHERE 75 S Per 100 Gals.	Time of Application	Recommended Application
Roses	For Beetles	2.3 lb. (10.5 oz.)	As beetles begin to appear. Use the higher amount when the large larvae are present.	Apply the specified amount of ORTHERE 75 S Soluble Powder in 100 gals. water with a hydraulic sprayer as a full coverage spray.
Lilacs Pines Orchids Anemones Cacti Poinsettia	For Beetles	2.3 lb. (10.5 oz.)	As the insects begin to appear. Repeat application at 2 week intervals may be necessary for control of mealybugs and scales.	Apply the specified amount of ORTHERE 75 S Soluble Powder in 100 gals. water with a hydraulic sprayer as a full coverage spray. The addition of a wetting agent may be required on difficult to wet foliage.
Stachys (Cranberry)	For Beetles	2.3 lb. (10.5 oz.)	As the insects begin to appear. Repeat	Caution: Phytotoxicity has occurred on the following foliage plants: Begonia, Hibiscus, Cactus

BEST AVAILABLE COPY

				the 2 to 3 oz. 1/2 lb. (5.3 oz.) higher amount when the large larvae are present.	Apply the specified amount of ORTHENE 75 S Soluble Powder in 100 gals. water with a hydraulic sprayer as a full coverage spray.
Foliage Plants Orchids Anthuriums Cacti Poinsettia	Aphids	1/2 lb. (5.3 oz.)	As aphids begin to appear.	Apply the specified amount of ORTHENE 75 S Soluble Powder in 100 gals. water with a hydraulic sprayer as a full coverage spray. The addition of a wetting agent may be required on difficult to wet foliage.	
	Mealybugs Thrips Whiteflies	2/3 lb. (10.5 oz.)	As the insects begin to appear. A repeat application, at a 2 week interval, may be necessary for control of mealybugs and whiteflies.	Caution: Phytotoxicity has occurred on the following foliage plants: <i>Blechnum spaldingii</i> , <i>Cissus antarctica</i> , <i>Ficus triangularis</i> , <i>Fittelia verschaffelii</i> , <i>Maranta leuconeura kerchoveana</i> , <i>Pachystachya lutea</i> , <i>Plectranthus australis</i> , <i>Polypodium aureum</i> , <i>Polystichum</i> , <i>Psidium guajava</i> , <i>Sinningia speciosa</i> . Before treating large plantings spray only a few plants and observe 2 weeks for varietal phytotoxicity.	
	Scales (Crawlers)	2/3 lb. (10.5 oz.)	As crawlers begin to appear. Repeat applications, at a 2 week or more interval, may be necessary where there is continuous crawler production.		
Roses Carnations Chrysanthemums	Aphids Thrips	2/3 lb. (10.5 oz.)	As aphids begin to appear. As thrips begin to appear or at the tight flower bud stage. Repeat applications may be necessary.	Apply the specified amount of ORTHENE 75 S Soluble Powder in 100 gals. water with a hydraulic sprayer as a full coverage spray. Do not apply more often than once every 28 days to Carnations and Chrysanthemums. Caution: Phytotoxicity has occurred on the following Chrysanthemum varieties: Abnissa, Bonnie Jean, Dixie, Garland, Gem, Iceberg, Pride, Showoff, Statesman, Tally Ho, Westward Ho, and Wild Honey. Before treating large Chrysanthemum plantings, spray only a few plants and observe two weeks for varietal phytotoxicity.	

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

NOTE: This product is sold by weight and package is full when packed, but due to fluffy nature this material is inclined to settle.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.
Open dumping is prohibited.

STORAGE

Keep pesticide in original container.
Do not put concentrate or dilute into food or drink containers.
Store in cool, dry place. Protect from excessive heat.
Do not contaminate food or foodstuffs.
Do not store or transport near food or feed.
For help with any spill, leak, fire or exposure involving this material, call day or night (800) 457-2022.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty container into application equipment. Do not reuse container. Dispose of empty container in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT INFORMATION, READ BEFORE USING PRODUCT.

LIMITED WARRANTY: Valent U.S.A. Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for use under average conditions when used strictly in accordance with the directions on the labeling. Valent does not make nor authorizes any agent or representative to make any other warranty, guarantee or representation, express or implied, concerning this product. Specifically, NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE IS MADE.

2. Critical and unforeseeable factors beyond Valent's control prevent it from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants and crops to which product is applied, lack of complete control over the handling and application of this product, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit under average conditions for the uses stated on the labeling and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Valent under 1 above) resulting from handling, storage and use of the product.

3. Precautions stated on the labeling should be followed to avoid hazardous exposure to the product.

LORSBAN® — Reg. TM of Dow Chemical Company for chlorpyrifos insecticide.
PYDRIN® — Reg. TM of E.I. du Pont de Nemours & Co., Inc. for fenvalerate insecticide.
ORTHENE® — Reg. TM of Chevron Chemical Company for acephate insecticide.

BEST AVAILABLE COPY