Propert 57637

Please read instructions on reverse before completing form. Form Approved OMB No 2070-0060 Approval expires 11-30 93 United States Environmental Protection Agency OPP Identifier Number Registration Office of Pesticide Programs (H7505C) **Amendment** Washington, DC 20460 167248 Application for Pesticide: Other Section I 2. EPA Product Manager 3 Proposed Classification Company/Product Number 59639-77 George T. LaRocca None Restricted 4. Company/Product (Name) TAME 2.4 EC Spray 15 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) 5. Name and Address of Applicant (Include ZIP Code) (b)(i), my product is similar or identical in composition and labeling Valent U.S.A. Corporation 1333 N California Blvd. 59639-77 Walnut Creek, CA 94596 EPA Reg. No. Check if this is a new address TAME 2.4 EC Spray Product Name\_ Section I i Final printed labels in response to Amendment - Explain below Agency letter dated\_ Resubmission in response to Agency letter dated "Me Too" Application Notification - Explain below. Other - explain below Explanation: Use additional page(s) if necessary. (For section I and Section II ) We are submitting five copies of finished labeling complying with PR Notice 93-3 inserting "For terrestrial uses", under the Environmental Hazards the label Section III 1. Material This Product WIN Be Packaged in: Child-Resistant Packaging **Unit Packaging** Water Soluble Packaging Type of Container Yes Metal Yes\* Yes Plastic No Glass Nο Nο Paper If "Yes." If "Yes," No. per No. per Other (Specify) Package wgt. Unit Package wgt container container ". Certification must be submitted. 3 Location of Net Contents Information Size(s) of Retail Container 5. Location of Label Directions On Label Container On Labeling accompanying product 6 Manner In Which Label Is Affixed To Product Lithograph Other ( Paper glued Stenailed Section IV 1 Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Name Telephone No. (Include Area Code) (510) 256-2745 Mark Herbst <u>Registration Assistant</u> Certification Date Application Received I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete Lacknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or (Stimped) both under applicable law Signature Registration Assistant Mark Herbst June 15, 1993



# TAME 2.4 EC Spray

# (INSECTICIDE-MITICIDE)

#### For Commercial Greenhouse Use Only.

Active Ingredient	By Wt.
*Fenpropathrin	
Inert Ingredients	
*(alpha-Cyano-3-phenoxybenzyl 2,2,3,3-tetramethyl cyclopropanecarboxylate)	
Contains 2.4 pounds a.i. per gallon.	
This product contains Xylene Range Aromatic Solvent	

# DANGER - PELIGRO

SEESIDEPANELFORSTATEMENTOFPRACTICAL TREATMENT AND ADDITIONAL PRECAUTIONARY STATEMENTS.

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Manufactured for VALENT U.S.A. Corporation Walnut Creek, CA 94596-8025 Made in U.S.A. Form R019A-06/15/93 EPA Reg. No. 59639-77

### **NET CONTENTS 1 QUART**

NGTEFICATION
LABEL NOT REVIEWED
FER PRINCTICE 64-6

#### **PRECAUTIONARY STATEMENTS**

## HAZARDS TO HUMANS & DOMESTIC ANIMALS DANGER

Corrosive, causes potentially irreversible eye damage. Do not get in eyes, on skin, or clothing. Contact with skin can result in a temporary burning, tingling, or itching sensation. Weer goggles or face shields and rubber gloves when handling. May be fatal if swellowed. Harmful if inheled or absorbed through skin. Avoid breething spray mist. Wash thoroughly with plenty of soap and water after handling and before eating, drinking, or using tobacco. Remove contaminated clothing and wash before reuse.

#### STATEMENT OF PRACTICAL TREATMENT

If in eyes: Immediately hold eyelids open and flush with a gentle

steady stream of water.

If on skin: Remove contaminated clothing and wash affected areas

with soap and water. Get medical attention,

If Inhaled: Remove victim to fresh air. If not breathing, give artificial

respiration, preferably mouth to mouth. Get medical

attention.

If swallowed: Call a physician or Poison Control Center immediately. If

medical advice cannot be obtained, drink promptly a large quantity of milk, egg white or gelatin moture, or if these are not available a large amount of water and take person and product container to the nearest emergency treatment facility. DO NOT INDUCE VOMITING. Petroleum distillates in product may cause aspiration pneumonia.

Note to Physicians: Possible nucceal damages may contraindicate the use of gastric larage. For Emergency Information Call 1-800-892-0099.

#### **ENVIRONMENTAL HAZARDS**

This product is extremely toxic to fish and squatic organisms and is toxic to wildlife. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

This product is highly took to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

#### PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat of open flame.

#### DIRECTIONS FOR USE

It is a violations of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

#### REENTRY STATEMENT

Do not reenter treated areas until sprays have dried.

#### STORAGE AND DISPOSAL

#### **PROHIBITIONS**

Do not contaminate water, food or feed by storage or cleaning of equipment

Open dumping is prohibited.

#### STORAGE

Keep pesticide in original container.

Do not put concentrate or dilute into food or drink containers

Not for use or storage in or around the home.

Do not store next to feed or food, or transport in or on vehicles obstaining foodstuffs or feeds.

Store in a cool, dry place.

Protect from heet

For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-892-0099, or Chemtrec 1-800-424-9300

#### PESTICIDE DISPOSAL

Pesticide westes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these weets carried be disposed of by use according to tabel instructions, confest year State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL

Triple rinse (or equivalent). Do not reuse container. Dispose of in a sanitary landfill or by other procedures allowed by State and local authorities.

#### DIRECTIONS

Do not apply this product through any type of irrigation system.

FOR GREENHOUSE USE AND FOR LATH HOUSE AND SHADE HOUSE USE (WHERE PLANTS ARE GROWN IN CONTAINERS) ONLY.

Application instructions: Apply specified desage in sufficient water for complete coverage. Spray all foliage surfaces including underside of leaves. Repeat as needed to maintain control. Under severe pest pressure, use the higher recommended rates and/or shorter spray intervals. TAME may be tank mixed and/or alternated with many commonly used insecticides and miticides. See recommendations for rotating sprays for SweetPotato Whitelly.

Note: Since ornamental varieties are numerous, constantly changing, and may react differently to TAME, test the production a small scale before making large-scale applications. High gallonage applications to certain chrysanthemum varieties may result in crop injury. Do not apply to chrysanthemums and roses with open flowers. Application of the TAME plus ORTHENE® tank mix on poincettles after bract formation may result in phytotoxicity on certain varieties. Certain strains of SweetPotato Whitefly may be more susceptible to a TAME plus ORTHENE® tank mix.

#### **PLANTS AND SHRUBS**

Crops	Pests	Greenhouse, Shade and Lath House Dosage PL. Oz./100 Gals.	Special instructions
Crossandra Columbine Foliage Plants Geranum Gladiolis Impetens Linope, Lify Mangold Pometha Snaporagon  Aphide  SweetPoteto Whitely, Adult, hymph cludes all inte and transition  Adult, hymph	Greenhouse	51/3 to 102/3	May betanit mead with ORTH-ENE Turl, Tree & Ornamental Spra- 1/3 le /1/10 gais. NOTE: Do no use the TAME plus ORTH-ENE tank me on the following plants Chamorries, Costandra, Colum tene. Geranum, Impatiene Linope, Lify, Mangold, and Snag dragon. These plants are no lateled on the Orthern Turl, Tree / Omarmental Spray tabel.
	Two-Spotted Spider Mile, Southern Red Mile (low to moderate pres- sure):	8 to 16	
	Best Armyworm	102/3	
	stages of citrus	16	
	Aphida	102/3	Tank mer with ORTHENE Turf Tree & Ornamental Spray 1/3 to 100 gate. (See NOTE above)
	SweetPoteo Wheely, Adult and Smell Hymphs (derly indexs)	16	Apply when adults are first de- tected. Apply up to 3 successive applications, then rotate to a non- pyrethrous product to avoid po- tented for resistance.
	Adult, Nymph (in- cludes all inster and transitional stepse (jrupee)).	102/3 Plus ORTHENE Turf, Tree & Ome when edules and rymphe appear. If a p one application of the landware and folio at 10 ft. or_1/100 gale. Apply up to 3 succ non-pyrethroid/organephosphale insections. See instructions above in the N SweetPosto Whitely is present, tree; then TAME plus ORTHENE. (See NOT	oputation is well established, make in 5 to 7 days later with TAME alone easing applications, then rotate to a acide to excell potential for reseables Section. It a tolerant etrain of with EPA labeled inschedular other

Crops	Pests	Greenhouse, Shade and Lath House Dosage FL. Oz/100 Gals.	Special Instructions
SHRUBS Azalea Bonwoods Carrelia Cotonester	European Red Mile Lacebuge Leefnoppers Spot- ted Tentform Leefnmer Green- house Whitefly	51/3 to 102/3	Maybetank mead with ORTHENE Turf, Tree & Omemental Spray 1/3 to /100 gate
Groton Elaegnus Euonymus Gardenia Hibiscus	Two-Spotted Spider Mile Southern Red Mile (low to moderate mile pres- sure)	8 to 16	
Hollies Honeysuckie	Best Amnyworm Japanese Beste	102/3	
Hydrangea Junipër Lantana	A hids	102/3	Tank met with ORTHENE Turf, Tree & Omemorital Sersy 1/3 lb / 100 gate.
Mahonia Oleander Podocareus Pyracaneus	SweetPotato Whitefly Adult and Smell Nymphs (early instant)	16	Apply when adults are first de- tected. Apply up to 3 successive applications then rotate to a non- pyrethroid product to aroust po- terfall for resistance.
Rhododendron Rose Vibumum	Adult Nymph (m- chique all mater all "Michanorei sta, es (mase)	102/3 Plus ORTHENE Turf, Tree & Omes when adults and nymphs sepaci. If a pi one septication of the tentoms and folio- at 18 ff or /100 gats. Apply up to 3 succ- ron-pyrethrostorpenephraphate insec- tance. See instructions above in the N SwissPusits Whitely is present, treat in then TAME also ORTHEIN.	opulation is well established melté w 5 to 7 days later volt TAME alone assens applications from rotate to a facile to avoid potential for rese- lates Section of a tolerant strain of

Crops	Pests	Dosage FL. Oz / 100 Gafs.	Special Instructions	
TREFS Conters On avous	Apple Aphill European Red MAs San Jose Scale (crtwikts) Suptled Teneform Leatminer Things Wildly Apple Aphill	51/3 102/3	May be rank mirred with ORTHENE Turk Tree & Ome- mental Sersy 2/3 to /100 gate	
• •	"inc spotted Serdermite. Southern Red Mits. (low-to- mit). It im is pressure).	8 - 16		
Non-Rearing s Acete Pety 9	Coting with Leafrollers: European Red Mite McDaniel Mite Orange Torthy Pendamis Mish Tar nishes, Plant Tug	513	Apply pagening after paral	
	No sector Severmine Southern Red Mile (low-to- moderate mile problems)	5 - 15	er to ge 'manged	

TREES

#### MIXING INSTRUCTIONS

TAME mixes readily with water. When tank mixing with ORTHENE Turf, Tree & Ornamental Spray, first mix the recommended amount of ORTHENE Turf, Tree & Ornamental Spray in the appropriate amount of water, then add the TAME last. Agitate well. All applicable directions, restrictions and precautions on the ORTHENE EPA registered label are to be followed.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

# IMPORTANT INFORMATION. READ BEFORE USING PRODUCT.

LIMITED WARRANTY 1 Valent U.S.A. Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for use under average conditions when used strictly in accordance with the directions on the labeling. Valent does not make nor authorizes any agent or representative to make, any other warranty, guarantee or representation, express or implied, concerning this product. Specifically, NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE IS MADE.

- 2. Critical and unforeseeable factors beyond Valent's control prevent it from eliminating all risks in connections with the use of this product. Such risks include, but are not limited to, damage to plants and crope to which product is applied, lack of complete control over the handling and application of this product, and damage caused by drift to other plants or crope. Such risks occur even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Valent under 1 above) resulting from handling, storage, and use of product.
- 3 Precautions stated on the labeling should be followed to avoid hazardous engueure to the product.

Copyright e1993 by Valent U.S.A. Corporation

ORTHENEs- Reg. TM of Chevron Chemical Company