

59639-99

09/28/2012

1/84



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D C 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

Robert L Hamilton, Ph D  
Valent U S A Corporation  
1600 Riviera Ave , Suite 200  
Walnut Creek, CA 94596-8025

SEP 28 2012

Subject Label Amendment to add new uses of Petition No 1F7886 (crop subgroup 6C, dried shelled pea and bean (except soybean), crop subgroup 20A, rapeseed, crop subgroup 20B, sunflower, and wheat) to the herbicide product labels listed below, containing the active ingredient, Flumioxazin  
EPA Reg Nos 59639-97, 59639-99, and 59639-119  
Decision Numbers 450810, 450814, 450815, 450817

Dear Dr Hamilton

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable, provided the following label amendment is made

- 1 Add the following statement to the PPE section of the end-use labels (59639-99 and 59639-119)  
"For aerial application to Field Peas, Flax, Lentils, Sunflower, Safflower, and Wheat, mixer/loaders must also wear filtering face piece respirator (N95, R95, or P95) "

One copy of labeling for these products, stamped "Accepted," is enclosed for your records Products released for shipment after 18 months from the date on this notice or the next printing of the label, whichever occurs first, must bear the new revised label Amended labeling will supersede all previously accepted ones Per 40 CFR 156 10(6), submit one copy of your final printed labeling before you release the product for shipment If you have questions or concerns regarding this letter, please contact Beth Benbow at (703) 347-8072 or email at [benbow.bethany@epa.gov](mailto:benbow.bethany@epa.gov)

Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn V. Montague", is written over the typed name.

Kathryn V Montague  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)



GROUP	14	HERBICIDE
-------	----	-----------

ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
SEP 28 2012

Note ***Bold italicized text*** is information for the reader and is not part of the label

[Bracketed information is optional text]

Under the Federal Insecticide  
Fungicide and Rodenticide Act  
as amended for the pesticide  
registered under EPA Reg. No.

59639-99

# VALOR<sup>®</sup> Herbicide

FOR CONTROL AND/OR SUPPRESSION OF CERTAIN WEEDS IN ALFALFA,  
ASPARAGUS, BUSHBERRIES, CELERY, COTTON, CUCURBIT VEGETABLES, DRY  
BEANS, FIELD CORN, FIELD PEAS, FLAX FRUITING VEGETABLES (INCLUDING  
OKRA), GARLIC, GRAPE, HOPS, LENTILS, MINT, NUT TREES (INCLUDING  
PISTACHIO), ONION (DRY BULB), PEANUT, POME FRUIT, POTATO, SOYBEAN,  
STONE FRUIT, STRAWBERRY, SUGARCANE, SUNFLOWER AND SAFFLOWER,  
SWEET POTATO, WHEAT, NON-BEARING FRUIT TREES, FALLOW LAND AND TO  
MAINTAIN BARE GROUND ON NON-CROP AREAS OF FARMS, ORCHARDS AND  
VINEYARDS

Active Ingredient	By Wt
Flumioxazin*	51%
Other Ingredients	49%
Total	100%

\*2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione

VALOR<sup>®</sup> Herbicide is a water dispersible granule containing 51% active ingredient

KEEP OUT OF REACH OF CHILDREN

## CAUTION

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT \_\_\_\_ POUNDS

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

### CAUTION

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing dust and spray mist. Avoid contact with skin, eyes or clothing.

FIRST AID	
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration preferably by mouth-to-mouth if possible</li> <li>• Call a poison control center or doctor for further treatment advice</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice</li> <li>• Have person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact <b>1 800 892-0099</b> for emergency medical treatment information.	

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**Applicators and other handlers must wear** long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes and socks.

**For aerial application to sugarcane, mixer/loaders must also wear** coveralls, chemical resistant apron and chemical resistant boots.

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Users should

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing

## ENVIRONMENTAL HAZARDS

This product is toxic to non target plants and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff may be hazardous to non target plants and aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

This pesticide is toxic to plants and should be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

Under some conditions this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing. These methods also reduce pesticide run-off. Use of vegetation filter strips along rivers, creeks, streams, wetlands or on the downhill side of fields where run-off could occur will minimize water run off and is recommended.

***Note to EPA reviewer: if this product is shipped in containers greater than 50 lbs, the following environmental hazard statement will be added to the label***

[Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.]

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical resistant gloves made of waterproof material, shoes plus socks.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries or greenhouses.

Keep all unprotected persons out of operating areas or vicinity where there may be drift.  
Do not enter or allow others to enter treated areas until sprays have dried.

**DISCLAIMER, RISKS OF USING THIS PRODUCT  
LIMITED WARRANTY  
AND LIMITATION OF LIABILITY**

**IMPORTANT** Read the entire Label including this Disclaimer, Risks of Using this Product Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT rather return the unopened product within 15 days of purchase for a refund of the purchase price

**RISKS OF USING THIS PRODUCT**

The Buyer and User (referred to collectively herein as Buyer) of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include but are not limited to injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including but not limited to loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

**LIMITED WARRANTY**

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label under average use conditions when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

**LIMITATION OF LIABILITY**

To the fullest extent allowed by law Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes but is not limited to loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY LAW THE EXCLUSIVE REMEDY OF THE BUYER AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

**PROMPT NOTICE OF CLAIM**

To the extent consistent with applicable law allowing such requirements Valent must be provided notice as soon as Buyer has reason to believe it may have a claim but in no event later than twenty one days from date of planting or twenty one days from the date of application whichever is latter so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law if Buyer does not notify Valent of any claims in such period it shall be barred from obtaining any remedy.

**NO AMENDMENTS**

Valent and Seller offer this product and Buyer accepts it subject to the foregoing Disclaimer, Risks of Using This Product Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

**TANK MIXES**

**NOTICE** Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor to the extent allowed by applicable law.

Read and follow the entire label of each product to be used in the tank mix with this product.

## RESISTANCE MANAGEMENT RECOMMENDATIONS

VALOR is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to VALOR and other Group 14 herbicides. Weed species with acquired resistance to Group 14 herbicides may eventually dominate the weed population if Group 14 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by VALOR or other Group 14 herbicides.

To delay herbicide resistance consider

- Avoiding the consecutive use of VALOR or other target site of action Group 14 herbicides that might have a similar target site of action on the same weed species
- Using tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern
- Basing herbicide use on a comprehensive Integrated Pest Management (IPM) program
- Monitoring treated weed populations for loss of field efficacy
- Contacting your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes

For further information or to report suspected resistance, you may contact Valent U.S.A. Corporation at the following toll-free number: 800 682-5368.

## TABLE OF CONTENTS

### USE INFORMATION

- Restrictions and Limitations
- Environmental Conditions and Biological Performance
  - Preemergence Application
  - Burndown Application
  - Postemergence Application
- Rainfastness
- Soil Characteristics
- Herbicide Rate
  - Residual Weed Control
- Carrier Volume and Spray Pressure
  - Preemergence Application
  - Burndown Application
  - Postemergence Application
- Additives
  - Burndown Application
- Jar Test to Determine Compatibility of Adjuvants and VALOR
- Sprayer Preparation
- Mixing Instructions
- Sprayer Cleanup
- Application Equipment
- Broadcast Application
- Band Application
- Aerial Application
- Chemigation
- Application with Dry Bulk Fertilizers

Rotational Restrictions

Broadleaf Weeds Controlled by Residual Activity of VALOR Table 1

Weeds Suppressed by Residual Activity of VALOR Table 2

### **DIRECTIONS FOR USE IN FALL AND SPRING PREPLANT BURNDOWN AND FALLOW SEEDBED PROGRAMS IN FIELD CORN, PEANUT AND SOYBEAN**

- Restrictions and Limitations
- Fall Burndown and Fallow Seedbed Programs
- Weeds Controlled by Fall and Spring Preplant Burndown Programs Table 3
- Spring Burndown Programs

### **DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN PROGRAMS IN COTTON AND SUGARCANE**

- Restrictions and Limitations
- Fall Burndown Programs
- Spring Burndown Programs

### **DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN PROGRAMS IN RICE, SORGHUM, SUNFLOWERS, TOBACCO AND WHEAT (Preplant to Crop)**

- Restrictions and Limitations
- Fall Burndown Programs
- Spring Burndown Programs



**DIRECTIONS FOR USE IN FALL BURNDOWN PROGRAMS IN FIELDS TO BE PLANTED TO BARLEY, FIELD PEA, FLAX, LENTIL, SAFFLOWER, SUNFLOWER AND SPRING WHEAT (Preplant to Crop)**

**DIRECTIONS FOR USE IN FALLOW LAND**

**DIRECTIONS FOR USE IN ESTABLISHED ALFALFA**

Restrictions and Limitations  
Timing to Alfalfa  
Timing to Weeds

**DIRECTIONS FOR USE IN ESTABLISHED ASPARAGUS**

Restrictions and Limitations  
Timing to Asparagus  
Timing to Weeds

**DIRECTIONS FOR USE IN CELERY**

Restrictions and Limitations  
Timing to Celery  
Timing to Weeds

**DIRECTIONS FOR USE IN COTTON**

Restrictions and Limitations  
Environmental Conditions and Biological Performance  
Herbicide Rate  
Emerged Broadleaf Weeds Controlled by Hooded Shielded and Layby Application of VALOR  
Tank Mixes with Glyphosate or MSMA in Cotton Table 4  
Carrier Volume and Spray Pressure  
Additives  
Application Equipment  
Timing to Cotton  
Timing to Weeds  
Tank Mixes  
Tank Mixes with VALOR for Hooded Shielded and/or Layby Use in Cotton Table 5

**DIRECTIONS FOR USE IN CUCURBIT VEGETABLES**

Restrictions and Limitations  
Timing to Cucurbit Vegetables  
Timing to Weeds

**DIRECTIONS FOR USE IN DRY BEANS**

Weed Suppression  
Restrictions and Limitations  
Timing to Dry Beans  
Timing to Weeds  
Additional Residual Grass Control  
Harvest aid  
Restrictions and Limitations  
Timing to Dry Beans

**DIRECTION FOR USE IN FIELD CORN**

Restrictions and Limitations  
Timing to Field Corn  
Burndown Use Directions – For Preplant Application in Field Corn  
Increasing Speed of Glyphosate Burndown Activity  
Tank Mixes  
Tank Mix Partners for Burndown and/or Residual Control of Weeds in Field Corn Table 6  
Tank Mix Restrictions

**DIRECTIONS FOR USE IN FIELD PEAS**

- Weed Control
  - Restrictions and Limitations
  - Timing to Field Peas
  - Timing to Weeds
  - Additional Residual Grass Control
- Harvest Aid
  - Restrictions and Limitations
  - Timing to Field Peas

**DIRECTIONS FOR USE IN FLAX**

- Harvest Aid
  - Restrictions and Limitations
  - Timing to Flax

**DIRECTIONS FOR USE IN FRUITING VEGETABLES (INCLUDING OKRA) ROW MIDDLES**

- Restrictions and Limitations
- Timing to Fruiting Vegetables
- Timing to Weeds

**DIRECTIONS FOR USE IN GARLIC**

- Restrictions and Limitations
- Timing to Garlic
- Timing to Weeds

**DIRECTIONS FOR USE IN HOPS**

- Restrictions and Limitations
- Timing to Hops for Sucker Control
- Timing to Hops for Preemergence Weed Control
- Timing to Weeds

**DIRECTIONS FOR USE IN LENTILS**

- Harvest Aid
  - Restrictions and Limitations
  - Timing to Lentils

**DIRECTIONS FOR USE IN MINT (Peppermint and Spearmint)**

- Restrictions and Limitations
- Timing to Mint
- Timing to Weeds

Weeds Controlled by Residual Activity of VALOR

Table 7

**DIRECTIONS FOR USE IN ONION (DRY BULB)**

- Restrictions and Limitations
- Timing to Onion (dry bulb)
- Timing to Weeds

**DIRECTIONS FOR USE IN PEANUT**

- Restrictions and Limitations
- Wind Management
- Timing to Peanuts
- Timing to Weeds
- Additional Residual Grass Control Sequential
- Additional Residual Grass Control Tank Mixed

## **DIRECTIONS FOR USE IN POTATO**

Restrictions and Limitations

Timing to Potatoes

Timing to Weeds

Weeds Suppressed by Residual Activity of VALOR at 1.5 oz/A

Table 8

## **DIRECTIONS FOR USE IN SOYBEAN**

Restrictions and Limitations

Timing to Soybeans

Timing to Weeds

Tank Mixes

Tank Mix Partners for Control of Emerged Weeds in Reduced Tillage Soybeans

Table 9

Additional Residual Broadleaf Control

Additional Residual Grass Control

ROUNDUP READY® Program

## **DIRECTIONS FOR USE IN STRAWBERRY**

Restrictions and Limitations

Weeds Controlled by Preemergence Application of VALOR

Table 10

## **DIRECTIONS FOR USE IN SUGARCANE**

Restrictions and Limitations

Timing to Sugarcane

Timing to Weeds

Broadleaf Weeds Controlled by Post-Directed or Layby Application of VALOR  
in Sugarcane

Table 11

Tank Mixes

Tank Mixes with VALOR for Post-Directed or Layby Use in Sugarcane

Table 12

Additional Preemergence Broadleaf Control

Additional Preemergence Grass Control

## **DIRECTIONS FOR USE IN SUNFLOWER AND SAFFLOWER**

Harvest Aid

Restrictions and Limitations

Timing to Sunflower and Safflower

## **DIRECTIONS FOR USE IN SWEET POTATO**

Restrictions and Limitations

Timing to Sweet Potatoes

Timing to Weeds

## **DIRECTIONS FOR USE IN WHEAT**

Restrictions and Limitations

Pre-plant applications Pre-emergence Weed Control

Restrictions and Limitations

Burndown Use Directions – For Preplant Applications in Wheat

Post-Plant Pre emergence Weed Control

Restrictions and Limitations

Use Directions Post-plant Pre emergence Weed Control

Harvest Aid

Restrictions and Limitations

Use Directions – Harvest Aid

Timing to Wheat

**DIRECTIONS FOR USE IN BUSHBERRIES, GRAPES, NUT TREES (INCLUDING PISTACHIO),  
POME FRUIT, STONE FRUIT AND NON-BEARING FRUIT TREES**

Restrictions and Limitations

Preemergence Application

Postemergence Application

Carrier Volume and Spray Pressure

Banded Application

Use Precautions for Bushberries

Use Precautions for Grapes

Use Precautions for Nut Trees (Including Pistachio) Pome Fruit and Stone Fruit

Use Precautions for Non Bearing Fruit Trees

Weeds Controlled by Postemergence Activity of VALOR Tank Mixes

Table 13

Additional Residual Weed Control

**DIRECTIONS FOR USE TO MAINTAIN BARE GROUND ON NON CROP AREAS OF FARMS,  
ORCHARDS OR VINEYARDS**

Restrictions and Limitations

Preemergence Application

Postemergence Application

Tank Mix Combinations to Maintain Bare Ground Non-Crop Areas

Table 14

**STORAGE AND DISPOSAL**

## USE INFORMATION

### VALOR uses

- VALOR provides residual control of susceptible weeds in alfalfa asparagus bushberries celery cotton cucurbit vegetables dry bean field corn garlic grape hops mint nut trees (including pistachio) onion (dry bulb) non bearing fruit trees peanut pome fruit potato soybean stone fruit strawberry sugarcane and sweet potato
- VALOR provides additional burndown activity when used as part of a burndown program in alfalfa asparagus celery cotton cucurbit vegetables dry bean field corn fruiting vegetables (including okra) row middles grape hops nut trees (including pistachio) non-bearing fruit trees peanut soybean and sugarcane  
VALOR can be applied as part of a fall burndown program for control of susceptible winter annuals
- VALOR can be applied with a hooded or shielded sprayer as well as part of a layby application in cotton and sugarcane for postemergence weed control as well as residual control of susceptible weeds
- VALOR can be used on farms orchards and vineyards for non-selective vegetation control to maintain bare ground non-crop areas that must be kept weed free
- **Read tank mix product label for rates and weeds controlled Always read and follow label directions for all tank mix products before using The most restrictive labeling of any tank mix product must be followed VALOR, when applied according to label use directions, will control the weeds claimed in crop specific use directions This label makes no claims concerning control of other weed species**

### AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR

The interaction of many equipment and weather related factors determine the potential for spray drift The applicator is responsible for considering all of these factors when making decisions Where states have more stringent regulations they should be observed

### RESTRICTIONS AND LIMITATIONS

Do not apply this product when weather conditions favor spray drift from treated areas

Do not apply during low-level inversion conditions including fog

When applying by air observe drift management restrictions and precautions listed under "AERIAL APPLICATION"

Do not apply to frozen or snow covered soil

Mechanical incorporation into the soil will reduce residual weed control

Post directed and layby applications of VALOR should be applied only to healthy growing crops

Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation

Do not apply within 300 yards of non dormant pears

Do not apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be applied immediately after application

**Spray equipment used to apply VALOR should not be used to apply other materials to any crop foliage, unless the proper cleanout procedures are followed See ' SPRAYER CLEANUP' for more information**

### ENVIRONMENTAL CONDITIONS AND BIOLOGICAL PERFORMANCE

#### Preemergence Application (Conventional Tillage)

**Important** Crop injury may occur from applications made to poorly drained soils and/or applications made under cool wet conditions Risk of crop injury can be minimized by using on well drained soils planting at least 1 5 inches deep using high quality seed and completely covering seeds with soil prior to preemergence applications Treated soil that is splashed onto newly emerged crops may result in temporary crop injury

Moisture is necessary to activate VALOR in soil for residual weed control Dry weather following applications of VALOR may reduce effectiveness However when adequate moisture is received after dry conditions VALOR will control susceptible germinating weeds VALOR may not control weeds that germinate after application but before an activating rainfall/irrigation or weeds that germinate through cracks resulting from dry soil

When adequate moisture is not received after a VALOR application weed control may be improved by irrigation with at least 1/4 inch of water. If emerged weeds are controlled by cultivation residual weed control will be reduced.

#### **Burndown Application**

For best results VALOR should be applied as part of a burndown program to actively growing weeds. Applying VALOR under conditions that do not promote active weed growth will reduce herbicide effectiveness. Do not apply VALOR when weeds are under stress due to drought, excessive water, extremes in temperature, disease or low humidity. Weeds under stress tend to become less susceptible to herbicidal action. VALOR is most effective when applied under warm, sunny conditions.

Reduced residual weed control may occur when burndown applications are made to fields where heavy crop and/or weed residue exist.

#### **Postemergence Application**

VALOR should only be applied to healthy crops labeled for postemergence use. Do not apply VALOR to crops that have been weakened by disease, drought, flooding, excessive fertilization, soil salts, previously applied pesticides, nematodes, insects or winter injury.

#### **Rainfastness**

VALOR is rainfast one hour after application. Applications should not be made if rain is expected within one hour of application or postemergence efficacy may be reduced.

#### **Soil Characteristics**

Application of VALOR to soils with high organic matter and/or high clay content may require higher dosages than soils with low organic matter and/or low clay content. Application to cloddy seedbeds can result in reduced weed control.

### **HERBICIDE RATE**

#### **Residual Weed Control (Including Preemergence Applications or Applications as Part of a Fall or Spring Burndown and Fallow Seedbed Program)**

Based upon soil characteristics (organic matter content and texture), the most difficult to control weed species being targeted, and the crop being grown, select the proper VALOR dosage from the rate range tables contained in this label.

**CARRIER VOLUME AND SPRAY PRESSURE** (Ground Equipment only. See Information for Aerial Equipment under AERIAL APPLICATION.)

#### **Preemergence Application (Conventional Tillage)**

To ensure uniform coverage, use 10 to 30 gals. of spray solution per acre for conventional tillage applications. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for preemergence herbicide application.

#### **Burndown Application (Prior to Crop Emergence)**

To ensure thorough coverage in burndown applications, use 15 to 60 gals. spray solution per acre. Use 20 to 60 gals. per acre if dense vegetation or heavy crop residue is present. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for postemergence herbicide application. Do not use flood jet nozzles.

**Postemergence Application (Emerged Crop)** Check use directions for specific crops in which VALOR can be applied postemergence. To ensure thorough coverage in burndown applications, use a minimum of 15 gallons spray solution per acre. Use a minimum of 20 gallons per acre if dense vegetation or heavy crop residue is present. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for postemergence herbicide application.

## ADDITIVES

### Burndown Application (Prior to Crop Emergence)

Postemergence control of weeds from VALOR tank mixes will require the addition of an agronomically approved adjuvant to the spray mixture. When an adjuvant is to be used with VALOR, Valent recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Either a crop oil concentrate or methylated seed oil which contains at least 15% emulsifiers and 80% oil or a non-ionic surfactant at 0.25% v/v may be used when applying VALOR as part of a burndown program. Some tank mix partners, such as Roundup Power Max<sup>®</sup>, are formulated with sufficient adjuvants and do not require the addition of a crop oil concentrate, methylated seed oil or non-ionic surfactant when tank mixed with VALOR. The addition of a crop oil concentrate or methylated seed oil may increase the burndown activity on certain weeds such as cutleaf eveningprimrose and Carolina geranium. Mixing compatibility qualities should be verified by a jar test.

A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with either a crop oil concentrate, methylated seed oil or non-ionic surfactant to enhance weed control. The addition of a nitrogen source does not replace the need for a crop oil concentrate, a methylated seed oil or a non-ionic surfactant.

### JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND VALOR

When using VALOR and an adjuvant, such as in stale seed bed, layby, hooded/shielded or reduced tillage situations, a jar test should be performed before mixing commercial quantities of VALOR when using VALOR for the first time, when using new adjuvants or when a new water source is being used.

- 1 Add 1 pt. of the water to a quart jar. The water should be from the same source and temperature as which will be used in the spray tank mixing operation.
- 2 Add 1 g of VALOR to the quart jar for every 3 oz. of VALOR per acre being applied (4 g if 12 oz/A is the desired VALOR rate). Gently mix until product goes into suspension.
- 3 Add 60 ml (4 Tbsps. or 2 fl. oz.) of the crop oil or methylated seed oil to the quart jar or 1 ml of non-ionic surfactant if it is being used in place of oil. Gently mix.
- 4 If nitrogen is being used, add 16 ml (1 Tbsp. or 0.5 oz.) of the 28 to 32% nitrogen source to the quart jar. If ammonium sulfate is being used, add 19 g AMS to the quart jar in place of the 28 to 32% nitrogen.
- 5 Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
- 6 An ideal tank mix combination will be uniform and free of suspended particles. If any of the following conditions are observed, the choice of adjuvant should be questioned:
  - a) Layer of oil or globules on the mixture's surface
  - b) Flocculation: fine particles in suspension or as a layer on the bottom of the jar
  - c) Clabbering: thickening texture (coagulated) like gelatin

### SPRAYER PREPARATION

Before applying VALOR, start with clean, well-maintained application equipment. The spray tank, as well as all hoses and booms, must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. Some pesticides, including but not limited to the sulfonylurea and phenoxy herbicides (i.e. Classic<sup>®</sup> and 2,4-D respectively), are active at very small amounts and can cause crop injury when applied to susceptible crops. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply VALOR. If two or more products were tank mixed prior to VALOR application, the most restrictive cleanup procedure should be followed.

### MIXING INSTRUCTIONS

- 1 Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
- 2 If a drift retardant is to be used, add 10 lbs of spray grade ammonium sulfate per 100 gals. of spray solution.
- 3 To ensure a uniform spray mixture, pre-slurry the required amount of VALOR with water prior to addition to the spray tank. Use a minimum of 1 gal. of water per 10 oz. of VALOR.
- 4 While agitating, slowly add the pre-slurried VALOR to the spray tank. Agitation should create a rippling or rolling action on the water surface.
- 5 If tank mixing VALOR with other labeled herbicides, add water-soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
- 6 Add any required adjuvants.
- 7 Fill spray tank to desired level with water. **Agitation should continue until all spray solution has been applied.**
- 8 Mix only the amount of spray solution that can be applied the day of mixing. VALOR should be applied within 6 hours of mixing.

### SPRAYER CLEANUP

Spray equipment including mixing vessels and nurse tanks must be cleaned each day following VALOR application. After VALOR is applied, the following steps must be used to clean the spray equipment:

- 1 Completely drain the spray tank, rinse the sprayer thoroughly including the inside and outside of the tank and all in-line screens.
- 2 Fill the spray tank with clean water and flush all hoses, booms, screens, and nozzles.
- 3 Top off tank, add 1 gal of 3% household ammonia (or equivalent) for every 100 gals of water, circulate through sprayer for 5 minutes, and then flush all hoses, booms, screens, and nozzles for a minimum of 15 minutes. If diaphragms are being used on the spray boom, loosen diaphragms before flushing the spray system, allowing cleaning solution to spray through the open diaphragm. If spray lines have any end caps, they must be loosened before flushing the system, allowing cleaning solution to spray through the loosened caps. To enhance removal of VALOR from the spray system, add a tank cleaner such as 'Valent Tank Cleaner' from Valent U.S.A. Corporation in place of ammonia and allow the cleaning solution to remain in the pressurized spray system (spray tank, hoses, and boom) overnight before flushing the system for a minimum of 15 minutes.
- 4 Drain tank completely.
- 5 Add enough clean water to the spray tank to allow all hoses, booms, screens, and nozzles to be flushed for 2 minutes.
- 6 Remove all nozzles and screens and rinse them in clean water.

Spray equipment including all tanks, hoses, booms, screens, and nozzles should be thoroughly cleaned before it is used to apply postemergence pesticides. Equipment with VALOR residue remaining in the system may result in crop injury to the subsequently treated crop.

### APPLICATION EQUIPMENT

Application equipment should be clean and in good repair. Nozzles should be uniformly spaced on boom and frequently checked for accuracy.

### BROADCAST APPLICATION

Apply VALOR and VALOR tank mixes with ground equipment using standard commercial sprayers equipped with flat fan or flood nozzles (preemergence applications only) designed to deliver the desired spray pressure and spray volume.

### BAND APPLICATION

When banding, use proportionately less water and VALOR per acre. The rate of VALOR required per acre when applied as a banded application can be calculated with the following formula:

Amount Needed per Acre for Banded Application	=	$\frac{\text{Band Width in Inches}}{\text{Row Width in Inches}}$	X	Rate per Broadcast Acre
--------------------------------------------------	---	------------------------------------------------------------------	---	-------------------------

### AERIAL APPLICATION

Spray drift away from the site of application may cause damage to non-target vegetation. To minimize drift, apply the largest droplet size consistent with uniform coverage and satisfactory weed control. To obtain satisfactory application and avoid drift, the following directions must be observed:

- Do not apply during low-level inversion conditions (including fog) when winds are gusty or under other conditions that favor drift. Do not spray when wind velocity is less than 2 mph or more than 10 mph.
- Do not apply this product by air within 40 ft. of non-target plants including non-target crops.
- Do not apply this product by air within 100 ft. of emerged cotton crops.
- Do not apply this product by air within 40 ft. of streams, wetlands, marshes, ponds, lakes, and reservoirs.
- **Carrier Volume and Spray Pressure** When used as part of a burndown weed control program, apply VALOR in 7 to 10 gals. of water per acre. Application at less than 7 gals. per acre may provide inadequate control. When used for preemergence weed control, apply VALOR in 5 to 10 gals. of water per acre. The higher gallonage applications generally afford more consistent weed control. Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.



- **Nozzle Selection and Orientation** Formation of very small drops may be minimized by appropriate nozzle selection by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray pressure. Use nozzles that produce flat or hollow cone spray patterns. Use non-drip type nozzles such as diaphragm type nozzles to avoid unwanted discharge of spray solution. The nozzles must be directed toward the rear of the aircraft at an angle between 0 and 15° downward. Do not place nozzles on the outer 25% of the wings or rotors.
- **Adjuvants and Drift Control Additives** Refer to tank mix partner's label for adjuvant recommendation. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

### CHEMIGATION

Follow all label recommendations for crops regarding rates, timing of application, special instructions and precautions.

[For [onion (dry bulb)] [and] [potatoes] follow supplemental labeling provided by Valent U.S.A. Corporation.]

Apply this product only through center pivot systems. End guns must be turned off due to uneven application. Do not apply this product through any other type of irrigation system.

Crop injury, lack of efficacy or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

The system must be properly calibrated (with water only) to ensure that the amount of VALOR applied corresponds to the recommended rate.

Apply VALOR in 1/2 to 3/4 inches of water during the first sprinkler set. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide, a dye indicator may be injected into the lines to mark the end of the application period. Once chemigation has begun, the run must be completed to ensure no product is left in the system.

If you have any questions about calibration, you should contact your State Extension Service Specialist, equipment manufacturers or other experts.

### Special Precautions for Chemigation

- 1 Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- 2 A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.
- 3 The system must be free of leaks and clogged nozzles.
- 4 The pesticide must be supplied continuously for the duration of the aqueous application. An uneven application may cause injury to the crop or poor weed control.
- 5 Agitation must be maintained in the nurse tank.
- 6 The sprinkler chemigation system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- 7 The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump.
- 8 The pesticide injection pipeline must contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 9 The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in the case where there is no water pump when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 10 The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 11 Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with the pesticides and capable of being fitted with a system interlock.
- 12 Do not apply when wind speed favors drift beyond the area intended for treatment.

### Chemigation Systems Connected to Public Water Systems

- 1 Public water system means a system for the provision to the public of piped water for human consumption if such a system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year
- 2 Chemigation systems connected to the public water system must contain a functional reduced pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction As an option to the RPZ the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe
- 3 All chemigation systems connected to the public water system must also follow restrictions listed in the preceding section titled **Special Precautions for Chemigation**

### APPLICATION WITH DRY BULK FERTILIZERS

Dry bulk fertilizer may be impregnated or coated with VALOR Application of dry bulk fertilizer with VALOR provides weed control equal to or slightly below the same rate of VALOR applied in liquid carriers due to better coverage with application via spray equipment Follow label recommendations for VALOR regarding rates special instructions cautions and special precautions Apply 400 to 700 lbs of the fertilizer/herbicide mixture per acre to obtain adequate soil coverage Apply the mixture to the soil with properly calibrated equipment immediately after blending Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury and to obtain uniform weed control

Ammonium nitrate and/or limestone should not be used as the sole source of fertilizer as the VALOR may not adhere to these materials

Compliance with all Federal and State regulations relating to blending pesticide mixtures with dry bulk fertilizer registrations labeling and application are the responsibility of the individual and/or company offering the fertilizer and VALOR mixture for sale

VALOR must be premixed with water to form a slurry prior to impregnation on dry bulk fertilizer For best results use a minimum of 1 pt of water for each 2 oz of VALOR A minimum of 6 pts of the VALOR slurry should be used to impregnate 2000 lbs of the fertilizer for uniform coverage of the fertilizer Closed drum belt ribbon or other commonly used dry bulk blenders may be used

The amount of VALOR required can be calculated with the following formula

$\frac{\text{ounces of VALOR per ton of fertilizer}}{1} = \frac{\text{ounces of VALOR per acre}}{1} \times 2000 - \frac{\text{pounds of fertilizer per acre}}{1}$
-------------------------------------------------------------------------------------------------------------------------------------------------------------------

Thoroughly clean dry fertilizer blending equipment after VALOR has been placed in the system to avoid injury to sensitive crops that may be treated with fertilizers blended after the equipment has been used for VALOR Rinse the sides of the blender and the herbicide tank with water Then impregnate the rinsate onto a load of dry fertilizer intended for an approved crop Use a maximum rate of 1 gal of rinsate per ton of fertilizer Follow with 1 to 2 loads of unimpregnated fertilizer in the blender before switching herbicides

## ROTATIONAL RESTRICTIONS

The following rotational crops may be planted after applying VALOR at the listed rate. Planting earlier than the recommended rotational interval may result in crop injury.

- Do not plant any crop, except corn (field), cotton, peanut, soybean, sugarcane and sweet potato earlier than 30 days after applying VALOR.

VALOR RATES	CROPS	ROTATION INTERVALS
1 oz /A	Cotton (no till or strip till only)	14 days <sup>1</sup>
1.5 to 2 oz /A	Cotton (no till or strip till only)	21 days <sup>1</sup>
2 oz /A or less	Peanut Soybean Sugarcane and Sweet Potato	immediately
	Field Corn (minimum and no till)	7 days
	Cotton and Field Corn (conventional tillage) Rice Sorghum Sunflower Tobacco and Wheat	30 days <sup>1</sup>
	Barley Dry and Snap Beans Flax Peas Rye Safflower and Sweet Corn	3 months
	Alfalfa Canola Clover Oats Potato Sugar Beet and all other crops not listed <sup>2</sup>	4 months if soil is tilled prior to planting 8 months if no tillage is performed
	Lentil	6 months
	Peanut Soybean Sugarcane and Sweet Potato	immediately
Up to 3 oz /A	Field Corn (minimum and no till)	14 days
	Field Corn (conventional tillage) and Sorghum	30 days <sup>1</sup>
	Cotton Rice Sunflower Tobacco and Wheat	2 months <sup>1</sup>
	Barley Dry and Snap Beans Flax Pea Rye Safflower and Sweet Corn	4 months
	Alfalfa Clover Oats Potato Sugar Beet	5 months if soil is tilled prior to planting 10 months if no tillage is performed
	Canola and all other crops not listed <sup>2</sup>	6 months if soil is tilled prior to planting 12 months if no tillage is performed
	Lentil	7 months
Up to 4 oz /A	Sugarcane	Immediately
	Alfalfa Canola Potato Sugar Beet and all other crops not listed <sup>2</sup>	6 months if soil is tilled prior to planting 12 months if no tillage is performed
	Cotton Field Corn Peanut Rice Sorghum Soybean Sunflower Tobacco and Wheat	4 months
	Transplanted on raised beds only melon pepper and tomato <sup>3</sup>	2 months (if the top 4 inches of the beds have been removed)
6 to 12 oz /A	Cotton Field Corn Peanut Rice Sorghum Soybean Sunflower Tobacco and Wheat	9 months
	Alfalfa Canola Sugar Beet and all other crops not listed <sup>2</sup> Trees can be transplanted 2 months after an application of VALOR <sup>4</sup>	12 months if soil is tilled prior to planting 18 months if no tillage is performed

<sup>1</sup> At least one inch of rainfall/irrigation must occur between application and planting or crop injury may occur.

<sup>2</sup> Successful soil bioassay must be performed prior to planting these crops.

<sup>3</sup> **Arizona, California and Hawaii only** For fallowbed application on transplanted melon, pepper and tomato beds follow supplemental labeling provided by Valent U.S.A. Corporation.

<sup>4</sup> Transplanted apple, apricot, avocado, bushberries (including blueberry), cherry, fig, grape, grapefruit, lemon, nectarine, nut trees (including pistachio), olive, orange, peach, pear, plum (including dried plum) and tangerine can be planted 2 months after a VALOR application of 2 to 12 oz /A.

Table 1 Broadleaf Weeds Controlled by Residual Activity of Valor Herbicide

BROADLEAF WEED SPECIES				
SECTION A				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	VALOR HERBICIDE RATE
Carpetweed	<i>Mollugo verticillata</i>	Up to 5%	All Soil Types	2 oz /A
Chickweeds				
Common	<i>Stellaria media</i>			
Mouseeear	<i>Cerastium vulgatum</i>			
Dandelion	<i>Taraxacum officinale</i>			
Eclipta	<i>Eclipta prostrata</i>			
Eveningprimrose Cutleaf	<i>Oenothera laciniata</i>			
Field Pennycress	<i>Thlaspi arvense</i>			
Florida Pusley	<i>Richardia scabra</i>			
Henbit	<i>Lamium amplexicaule</i>			
Lambsquarters Common	<i>Chenopodium album</i>			
Little Mallow	<i>Malva parviflora</i>			
Marestail/Horseweed	<i>Conyza canadensis</i>			
Mayweed/False Chamomile	<i>Matricaria maritima</i>			
Nightshades				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Hairy	<i>Solanum sarrachoides</i>			
Pigweeds				
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Lettuce	<i>Lactuca serriola</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane Common	<i>Portulaca oleracea</i>			
Radish Wild	<i>Raphanus raphanistrum</i>			
Redmaids	<i>Calandrinia ciliata var menziessii</i>			
Shepherd s purse	<i>Capsella bursa-pastoris</i>			
Smallflower Morningglory	<i>Jacquemontia tamnifolia</i>			
Sowthistle Prickly	<i>Sonchus asper</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Venice Mallow	<i>Hibiscus trionum</i>			

continued

**Table 1 Broadleaf Weeds Controlled by Residual Activity of Valor Herbicide (continued)**

<b>SECTION B</b>				
All weeds listed in Section A plus				
<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>	<b>ORGANIC MATTER</b>	<b>SOIL TYPE</b>	<b>VALOR HERBICIDE RATE<sup>2</sup></b>
Coffee Senna	<i>Cassia occidentalis</i>	Up to 3%	All Soil Types	2 oz /A Cotton and Dry Bean 2.5 oz /A Field Corn and Soybean 3 oz /A Peanut and all other labeled crops
Common Ragweed <sup>1</sup>	<i>Ambrosia artemisiifolia</i>			
False Chamomile	<i>Tripleurospermum maritima</i>			
Florida Beggarweed	<i>Desmodium tortuosum</i>			
Golden Crownbeard	<i>Verbesina encelioides</i>			
Hairy Indigo	<i>Indigofera hirsuta</i>			
Hemp Sesbania	<i>Sesbania exaltata</i>	3 to 5%	Coarse and Medium Soils (sandy loam loamy sand loamy silt loam silt sandy clay sandy clay loam)	2 oz /A Cotton and Dry Bean 2.5 oz /A Field Corn and Soybean 3 oz /A Peanut and all other labeled crops
Jimsonweed	<i>Datura stramonium</i>			
Kochia	<i>Kochia scoparia</i>			
London Rocket	<i>Sisymbrium irio</i>			
Morningglories <sup>3</sup>				
Entireleaf	<i>Ipomoea hederacea</i> var <i>integruscula</i>			
Ivyleaf	<i>Ipomoea hederacea</i>			
Red/Scarlet	<i>Ipomoea coccinea</i>			
Tall	<i>Ipomoea purpurea</i>			
Mustard Wild	<i>Brassica kaber</i>			
Palmer Amaranth	<i>Amaranthus palmeri</i>			
Spurred Anoda	<i>Anoda cristata</i>		Fine Soils (silty clay silty clay loam clay clay loam)	2 oz /A Cotton and Dry Bean 3 oz /A Field Corn Peanut Soybean and all other labeled crops
Tropic Croton	<i>Croton glandulosus</i>			
Waterhemp <sup>1</sup>				
Common	<i>Amaranthus rudis</i>			
Tall	<i>Amaranthus tuberculatus</i>			
Wild Poinsettia	<i>Euphorbia heterophylla</i>			
Yellow Rocket	<i>Barbarea vulgaris</i>			

<sup>1</sup>A postemergence herbicide such as COBRA® PHOENIX™ or glyphosate (ROUNDUP READY® soybeans only) may be needed following a preemergence application of Valor Herbicide to adequately control common ragweed or waterhemp in soybean fields with heavy pressure

<sup>2</sup>Due to differences in crop canopy timing between peanuts and soybeans 3 oz /A of Valor Herbicide should be used in peanuts regardless of soil type and organic matter content except in the states of North Carolina Oklahoma and Virginia where a maximum of 2 oz /A can be applied in peanuts unless supplemental labeling provided by Valent U S A Corporation is followed Valor Herbicide will provide residual control of these weeds at 2 oz /A when applied under a cotton canopy

<sup>3</sup>Morningglory species are not adequately controlled on fine soils or soils with greater than 3% organic matter

**Table 2 Weeds Suppressed by Residual Activity of Valor Herbicide**

BROADLEAF WEED SPECIES		ORGANIC MATTER	OUNCES PER ACRE
COMMON NAME	SCIENTIFIC NAME		
Bristly Starbur	<i>Acanthospermum hispidum</i>	Up to 5%	2 to 3
Copperleaf Hophornbeam	<i>Acalypha ostryifolia</i>		
Ragweed Giant	<i>Ambrosia trifida</i>		
Russian Thistle	<i>Salsola iberica</i>		
Smartweeds			
Ladysthumb	<i>Polygonum persicaria</i>		
Pennsylvania	<i>Polygonum pensylvanicum</i>		
Smellmelon	<i>Cucumis melo</i>		
Velvetleaf	<i>Abutilon theophrasti</i>		
Wild Buckwheat	<i>Polygonum convolvulus</i>		
Wormwood Biennial	<i>Artemisia biennis</i>		
GRASS WEED SPECIES			
Barnyardgrass	<i>Echinochloa crus galli</i>	Up to 5%	1 5 to 3
Bluegrass Annual	<i>Poa annua</i>		
Crabgrass Large	<i>Digitaria sanguinalis</i>		
Foxtail Giant	<i>Setaria faber</i>		
Goosegrass	<i>Eleusine indica</i>		
Lovegrass California	<i>Eragrostis diffusa</i>		
Panicums			
Fall	<i>Panicum dichotomiflorum</i>		
Texas	<i>Panicum texanum</i>		
Ryegrass Italian	<i>Lolium multiflorum</i>		
Signalgrass Broadleaf	<i>Brachiana platyphylla</i>		
Cheat	<i>Bromus secalinus</i>		
Downy Brome	<i>Bromus tectorum</i>		

**DIRECTIONS FOR USE IN FALL AND SPRING PREPLANT BURNDOWN AND  
FALLOW SEEDBED PROGRAMS IN FIELD CORN, PEANUT AND SOYBEAN  
(Preemergence to Crop)**

[For Use in the States of Arizona, California and Hawaii Only]

**RESTRICTIONS AND LIMITATIONS**

- Do not apply to frozen or snow covered soil
- Do not perform any tillage operation after application or residual weed control will be reduced
- [Observe all rotational intervals prior to planting as listed in the ROTATIONAL RESTRICTIONS table ]

**FALL BURNDOWN AND FALLOW SEEDBED PROGRAMS**

VALOR[ at 2 to 4 oz/A] can be used in the fall to provide residual weed control in fields that will be planted the following spring with field corn peanut or soybean [(refer to Rotational Restrictions table for rates and rotational intervals prior to planting)] Weeds controlled by residual activity are listed in Table 1 (sections A and B) Broadleaf Weeds Controlled by Residual Activity of VALOR Table 3 Weeds Controlled by Fall and Spring Preplant Burndown Programs and Table 7 Weeds Controlled by Residual Activity of VALOR If weeds have emerged at the time of application use VALOR in combination with a labeled burndown herbicide [Application must be made no earlier than October 15 in Region 2 or November 15 in Region 1 or when soil temperature falls below 50°F at a 2 inch depth to maintain residual weed control into the spring (April 1 in Region 1 and May 1 in Region 2) or up until planting whichever comes first ] VALOR can be used in a fall burndown or fallow seedbed program [outside of Regions 1 and 2] however the length of residual control may be variable

Abnormally warm or wet winters will reduce the length of weed control observed in the spring

[Fall Application Regions]

Region 1 Alabama Arkansas Georgia Kentucky Mississippi Oklahoma Tennessee and Virginia

Region 2 Delaware Kansas Illinois Indiana Iowa Maryland Michigan Minnesota Missouri Nebraska North Dakota Ohio Pennsylvania South Dakota West Virginia and Wisconsin]

Weeds controlled by postemergence or residual activity are listed in Table 3. Preplant burndown treatment tank mixes and rates are

Herbicide	Rate
<b>Program 1<sup>1</sup></b>	
VALOR Plus	2 to 3 oz/A
Glyphosate Plus	0.5 to 1.0 lb ai/A (equivalent to 1 to 2 pt/A of ROUNDUP Original®)
2,4-D LVE (2,4-D for use on preplant soybeans only) Plus	0.5 to 1.0 lb ai/A (equivalent to 1 to 2 pt/A of 2,4-D 4 LVE)
NIS + AMS	0.5% v/v + 17 lbs/100 gals of water

or

<b>Program 2<sup>1</sup></b>	
VALOR Plus	2 to 3 oz/A
Glyphosate Plus	0.5 to 1.0 lb ai/A (equivalent to 1 to 2 pt/A of ROUNDUP Original)
COC <sup>2</sup> or NIS + AMS	1 pt/A or 0.5% v/v + 17 lbs/100 gals of water

or

<b>Program 3<sup>1</sup></b>	
VALOR Plus	2 to 3 oz/A
2,4-D LVE (2,4-D for use on preplant soybeans only) Plus	0.5 to 1.0 lb ai/A (equivalent to 1 to 2 pt/A of 2,4-D 4 LVE)
COC	1 pt/A

<sup>1</sup>Dicamba (BANVEL®) at 0.188 lb ai/A (6 fl. oz./A of BANVEL 4) can be added to Programs 1, 2 & 3 to assist in the control of emerged broadleaves. Refer to dicamba label for rotational restrictions.

<sup>2</sup>Crop oil concentrate has been found to increase glyphosate burndown of emerged cutleaf evening primrose and Carolina geranium.



**Table 3 Weeds Controlled by Fall and Spring Preplant Burndown Programs**

WEEDS CONTROLLED <sup>1</sup>		POSTEMERGENCE			RESIDUAL
COMMON NAME	SCIENTIFIC NAME	Program 1	Program 2	Program 3	
		Weeds 3 inches or less			
Chamomile False	<i>Matricaria maritima</i>	Yes	Yes	No	Yes
Cheatgrass	<i>Bromus tectorum</i>	Yes	Yes	No	Yes
Chickweed Common	<i>Stellaria media</i>	Yes	Yes	No	Yes
Chickweed Mouseear	<i>Cerastium vulgatum</i>	Yes	Yes	No	Yes
Cockle White	<i>Silene latifolia</i>	No	Yes	Yes	Yes
Dandelion	<i>Taraxacum officinale</i>	Yes	No	Yes <sup>2</sup>	Yes
Deadnettle Purple	<i>Lamium purpureum</i>	Yes	Yes	Yes	Yes
Groundsel Cressleaf	<i>Senecio glabellus</i>	Yes	Yes		Yes
Henbit	<i>Lamium amplexicaule</i>	Yes	Yes	Yes	Yes
Kochia	<i>Kochia scoparia</i>	Yes	Yes	Yes	Yes
Marestail/Horseweed	<i>Conyza canadensis</i>	Yes	Yes <sup>3</sup>	Yes	Yes
Mallow Common	<i>Malva Neglecta</i>	Yes	Yes	No	Yes
Prickly Lettuce	<i>Lactuca serriola</i>	Yes	Yes	Yes	Yes
Wormwood Biennial	<i>Artemisia biennis</i>	Yes	Yes	Yes	Yes
<b>Weeds 12 inches or less</b>					
Canola Volunteer	<i>Brassica napus</i>	Yes	Yes	Yes	Yes
Carolina Geranium	<i>Geranium carolinianum</i>	Yes	Yes	Yes	-
Eveningprimrose Cutleaf <sup>4</sup>	<i>Oenothera laciniata</i>	Yes	Yes	Yes	Yes
Flixweed	<i>Descurainia sophia</i>	Yes	Yes	Yes	Yes
Mustard Tansy	<i>Descurainia pinnata</i>	Yes	Yes	Yes	Yes
Mustard Wild	<i>Brassica kaber</i>	Yes	Yes	Yes	Yes
Shepherd s purse	<i>Capsella bursa-pastors</i>	Yes	Yes	Yes	Yes

<sup>1</sup> Refer to glyphosate and/or 2,4-D labels for additional weeds controlled and rotational restrictions

<sup>2</sup> 1 lb ai/A of 2,4-D LVE (equivalent to 2 pt /A of 2,4-D 4 LVE) should be used for control of emerged dandelion

<sup>3</sup> Program 2 will not control emerged glyphosate resistant marestail/horseweed

<sup>4</sup> Program 1 should be used to control cutleaf eveningprimrose that are nearing 12 inches in height or are past the rosette stage

Programs 2 or 3 should be used to control cutleaf evening primrose that are 12 inches or less and in the rosette stage

### SPRING BURNDOWN PROGRAMS

VALOR can be used in combination with labeled preplant burndown herbicides to assist in the postemergence burndown of emerged weeds and provide residual weed control prior to crop emergence. Weeds controlled by residual activity are listed in Table 1.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row. Apply VALOR after planting peanuts and soybeans when these types of planters are used (within 3 days after planting soybeans, within 2 days after planting peanuts and before the crop emerges). VALOR cannot be applied after planting field corn.

VALOR can be used [at 1 to 3 oz/A] with labeled preplant burndown herbicides to enhance the speed of burndown and increase weed spectrum.

VALOR can be used [at 1 to 3 oz/A [1 to 2 oz/A]] in field corn, peanut and soybean burndown programs. See DIRECTIONS FOR USE IN FIELD CORN, DIRECTIONS FOR USE IN PEANUT, DIRECTIONS FOR USE IN SOYBEAN for more information.

## **DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN PROGRAMS IN COTTON AND SUGARCANE**

[For Use in the States of Arizona, California and Hawaii Only]

### **RESTRICTIONS AND LIMITATIONS**

- Do not apply to frozen or snow covered soil
- Do not perform any tillage operation after application or residual weed control will be reduced
- VALOR can be used [at 1 to 2 oz/A] with labeled burndown herbicides to enhance the speed of burndown and increase weed spectrum
- A minimum of 30 days must pass and 1 inch of rainfall/irrigation must occur between VALOR application and planting of conventionally tilled cotton
- A minimum of 14 days must pass and 1 inch of rainfall/irrigation must occur between VALOR application and planting of no-till or strip till cotton when a VALOR rate of 1 oz/A is used and 21 days when a VALOR rate of 1.5 to 2 oz/A is used. The field must contain the stubble from the previous crop
- VALOR can be applied as part of a burndown application to sugarcane until cane emergence
- Observe all rotational intervals prior to planting as listed in the **ROTATIONAL RESTRICTIONS** table
- Refer to most restrictive label for minimum interval between application and planting

### **FALL BURNDOWN PROGRAMS**

VALOR [at 2 to 4 oz/A] can be used in the fall to provide residual weed control in fields that will be planted the following spring with cotton or sugarcane [(refer to Rotational Restrictions table for rates and rotational intervals prior to planting)] Weeds controlled by residual activity are listed in Table 1 and Table 7. If weeds have emerged at the time of application use VALOR in combination with a labeled burndown herbicide. [Application must be made no earlier than October 15 in Region 2 or November 15 in Region 1 or when soil temperature falls below 50°F at a 2 inch depth to maintain residual weed control into the spring (April 1 in Region 1 and May 1 in Region 2) or up until planting whichever comes first.] [VALOR can be used in a fall burndown or fallow seedbed program outside of Regions 1 and 2.]

Abnormally warm or wet winters will reduce the length of weed control observed in the spring

### **SPRING BURNDOWN PROGRAMS**

VALOR [at 1 to 2 oz/A] can be used in combination with labeled preplant burndown herbicides to assist in the postemergence burndown of emerged weeds and provide residual weed control prior to crop emergence in fields that will be planted with cotton or sugarcane. Weeds controlled by residual activity are listed in Table 1.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row

## **DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN PROGRAMS IN RICE, SORGHUM, SUNFLOWER, TOBACCO AND WHEAT (Preplant to Crop)**

[For Use in the States of Arizona, California and Hawaii Only]

### **RESTRICTIONS AND LIMITATIONS**

- Do not apply to frozen or snow covered soil
- Do not perform any tillage operation after application or residual weed control will be reduced
- VALOR can be used [at 1 to 2 oz/A] with labeled burndown herbicides to enhance the speed of burndown and increase weed spectrum. A minimum of 30 days must pass and 1 inch of rainfall/irrigation must occur between VALOR application and planting of rice, sorghum, sugarcane, sunflowers, tobacco or wheat. Refer to most restrictive label for minimum interval between application and planting
- [Observe all rotational intervals prior to planting as listed in the **ROTATIONAL RESTRICTIONS** table.]

### **FALL BURNDOWN PROGRAMS**

VALOR can be used in combination with labeled burndown programs to control emerged weeds and provide residual weed control in fields that will be planted the following spring [(refer to Rotational Restrictions table for rates and rotational intervals prior to planting)] [Application must be made no earlier than October 15 in Region 2 or November 15 in region 1 or when soil temperature falls below 50°F at a two inch depth to maintain residual weed control into the spring.]

Abnormally warm winters may reduce the length of weed control observed in the spring

#### **SPRING BURNDOWN PROGRAMS**

VALOR can be used in combination with labeled burndown programs to control emerged weeds and provide residual weed control prior to crop emergence. Weeds controlled by residual activity are listed in Table 1 Section A. Crops that will be planted following application must be in compliance with the rotational interval listed in the Rotational Restriction table above.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row.

### **DIRECTIONS FOR USE IN FALL BURNDOWN PROGRAMS IN FIELDS TO BE PLANTED TO BARLEY, FIELD PEA, FLAX, LENTIL, SAFFLOWER, SUNFLOWER AND SPRING WHEAT (Preplant to Crop)**

[For Use in the States of Arizona, California and Hawaii Only]

#### **RESTRICTIONS AND LIMITATIONS**

- Do not apply to frozen or snow covered soil
- Do not perform any tillage operation after application or residual weed control will be reduced
- *Valor* Herbicide can be mixed with 2,4-D and/or glyphosate formulations labeled for burndown programs (preplant to crop) in accordance with the most restrictive label limitations and precautions. Labeled application rates can not be exceeded. Do not mix *Valor* Herbicide with any product containing a label prohibition against such mixing.
- [Observe all rotational intervals prior to planting as listed in the ROTATIONAL RESTRICTIONS table.]

#### **FALL BURNDOWN PROGRAMS**

VALOR can be used [at 2 to 4 oz/A] with labeled burndown herbicides to enhance the speed of burndown, increase weed spectrum and provide residual weed control of the weeds listed in Table 3 until the following spring. Rotational intervals must be followed for crop to be planted in the spring following the fall VALOR application. Refer to most restrictive label for minimum interval between application and planting.

### **DIRECTIONS FOR USE IN FALLOW LAND**

[For Use in the States of Arizona, California and Hawaii Only]

VALOR may be used as a preemergence fallow treatment. Weeds controlled by residual activity are listed in Table 1.

VALOR [at 2 to 4 oz/A] can be used in the fall to provide residual weed control in fallow fields [(refer to Rotational Restrictions table for rates and rotational intervals prior to planting)]. If weeds have emerged at the time of application, use VALOR in combination with a labeled fallow herbicide. [Application must be made no earlier than October 15 in Region 2 or November 15 in Region 1 or when soil temperature falls below 50°F at a 2 inch depth to maintain residual weed control into the spring (April 1 in Region 1 and May 1 in Region 2)]. Abnormally warm or wet winters will reduce the length of weed control observed in the spring.

VALOR [at 1 to 4 oz/A] can be used in spring in combination with labeled burndown herbicides to control emerged weeds and provide residual weed control.

## DIRECTIONS FOR USE IN ESTABLISHED ALFALFA

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 4 oz of VALOR per acre during a single application
- Do not apply more than 8 oz of VALOR per acre during a single growing season
- Do not make a sequential VALOR application within 60 days of the first VALOR application
- Do not apply to alfalfa with greater than 6 inches of growth. Application will result in burning of treated leaves and stems. **Users should understand and accept this risk before using VALOR on alfalfa**
- Do not apply within 25 days of harvest or grazing
- Do not use on alfalfa grown for seed unless approved by a State authority to support a Special Local Need (SLN) under FIFRA section 24(c)
- Only apply with an adjuvant or tank mix with products formulated as an emulsifiable concentrate (EC) when targeting control of emerged weeds (crop burn and/or stunting should be expected and accepted if Valor Herbicide is used with an adjuvant, a tank mix partner formulated as an emulsifiable concentrate (EC) or a tank mix partner formulated with an adjuvant)
- Application with paraquat can be used to burndown winter annuals prior to winter dormant period
- Do not use on intended mixed alfalfa-grass stands

### TIMING TO ALFALFA

VALOR may be applied to established alfalfa with a maximum amount of growth of 6 inches or less for the preemergence control of the weeds listed in Table 7 Weeds Controlled by Residual Activity of VALOR. Established alfalfa is defined as alfalfa planted in the fall or spring which has gone through a first cutting/mowing. Application to alfalfa with greater than 6 inches of growth may result in unacceptable crop injury.

For control of winter annual weeds, the best timing for preemergence control is in the fall immediately after the last cutting or shearing off has occurred.

For control of summer annual weeds, the best timing for preemergence control is in the spring prior to alfalfa growth and before 6 inches of growth.

### TIMING TO WEEDS

#### Preemergence – Preemergence To Weeds

Apply VALOR before alfalfa growth exceeds 6 inches in height for the preemergence control of weeds listed in Table 7 Weeds Controlled by Residual Activity of VALOR. Applications should be made as soon as possible after cutting and removing alfalfa to minimize injury to alfalfa growth.

#### Postemergence Dodder Suppression

Apply Valor Herbicide at 4 oz per acre with an adjuvant for postemergence suppression of dodder. Tank mixes with Pursuit® Herbicide or Raptor® Herbicide will increase control.

## DIRECTIONS FOR USE IN ESTABLISHED ASPARAGUS

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 6 oz of VALOR per acre during a single application
- Do not apply more than 6 oz of VALOR per acre during a single growing season
- Apply only to dormant asparagus no less than 14 days before spears emerge Application to non-dormant asparagus may result in unacceptable crop injury
- [Do not work soil within 60 days prior to application in the spring Soil can be worked after spear harvest in preparation for Valor Herbicide application prior to fern emergence Treated soil that is splashed onto the ferns may result in spotting ]

### TIMING TO ASPARAGUS - Dormant

VALOR may be applied to dormant asparagus for preemergence control of the weeds listed in Table 10 Weeds Controlled by Preemergence Application of VALOR Application to non dormant asparagus will result in unacceptable crop injury Applications should be made no less than two weeks prior to spear emergence and must be sprinkler or rainfall incorporated with 0.5 to 0.75 inches of water or some scoring may result

### TIMING TO ASPARAGUS – Post Harvest

Apply *Valor* Herbicide after the final harvest of the season but prior to fern emergence for preemergence control of the weeds listed in Table 10 Weeds Controlled by Preemergence Application of *Valor* Herbicide Application after fern emergence will result in unacceptable crop injury Apply no less than two weeks prior to fern emergence and must be sprinkler or rainfall incorporated with 0.5 to 0.75 inches of water Add a burndown tank mix partner for the control of emerged weeds labeled for asparagus in accordance with the most restrictive labeled limitations and precautions

### TIMING TO WEEDS

#### Burndown – Dormant Asparagus, Postemergence to Weeds

VALOR may be used for residual weed control as well as to assist in postemergence burndown of many annual and perennial weeds where asparagus is dormant For control of emerged weeds tank mix VALOR with paraquat Refer to paraquat label for recommended rate and application parameters To ensure thorough coverage use a minimum of 15 gals of spray solution per acre VALOR tank mixes applied to assist in the control of emerged weeds must be applied with a non ionic surfactant at 0.25% v/v A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to increase herbicidal activity

#### Burndown – After Last Harvest of Season, Postemergence to Weeds

Use *Valor* Herbicide for residual weed control and to assist in postemergence burndown for many annual and perennial weeds where asparagus harvest has been completed for the year For control of emerged weeds use a labeled tank mix partner with activity on the emerged weeds

#### Preemergence – Dormant Asparagus or After Last Harvest of Season, Preemergence to Weeds

Apply VALOR to dormant asparagus for the preemergence control of weeds listed in Table 10 Weeds Controlled by Preemergence Application of VALOR

## DIRECTIONS FOR USE IN CELERY

[For Use in the States of [California], Michigan and Wisconsin Only]

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 3 oz of *Valor* Herbicide per acre during a pre transplant application
- [In the state of California use as pre-transplant application only ]
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a post-transplant application
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season
- Do not use with an adjuvant
- Post transplant applications must be made between 3 to 7 days following transplanting
- Do not apply as part of a tank mix

### TIMING TO CELERY

Apply *Valor* Herbicide at 3 oz/A prior to transplanting or between 3 and 7 days following transplanting for preemergence control of the weeds listed in Table 1 Broadleaf Weeds Controlled by Residual Activity of *Valor* Herbicide

### TIMING TO WEEDS

Use *Valor* Herbicide prior to weed emergence for residual control

**Read tank mix product label for rates and weeds controlled Always read and follow label directions for all tank mix products before using The most restrictive labeling of any tank mix product must be followed *Valor* Herbicide, when applied according to label use directions, will control the weeds listed in Table 1, Broadleaf Weeds Controlled by Residual Activity of *Valor* Herbicide This label makes no claims concerning control of other weed species**

## DIRECTIONS FOR USE IN COTTON

[For Use in the States of Arizona, California and Hawaii Only]

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 2 oz of VALOR per acre during a single application
- Do not apply more than 4 oz of VALOR per acre during a single growing season
- Do not make a sequential VALOR application within 30 days of the first VALOR application
- Do not apply within 60 days of harvest

### ENVIRONMENTAL CONDITIONS AND BIOLOGICAL PERFORMANCE

#### Hooded, Shielded and Layby Application

For best results VALOR should be applied to actively growing weeds within the growth stages indicated in this label Applying VALOR under conditions that do not promote active weed growth will reduce herbicide effectiveness Do not apply VALOR when the crop or weeds are under stress due to drought excessive water extremes in temperature disease or low humidity Weeds under stress tend to become less susceptible to herbicidal action VALOR is most effective when applied under sunny conditions at temperatures above 65°F

VALOR is rainfast one hour after application Applications should not be made if rain is expected within one hour of application or postemergence efficacy may be reduced Rainfall within one hour of application will not adversely affect residual activity

**HERBICIDE RATE****Hooded, Shielded and Layby Application**

For postemergence weed control VALOR should be applied through a hooded or shielded sprayer or at layby at 2 oz/A in combinations with MSMA or at 1 to 2 oz /A in combination with glyphosate to assist in the control of weeds listed in Table 4. Residual weed control can also be obtained through hooded, shielded and layby application of VALOR. Weeds that are controlled through residual activity of VALOR are listed in Table 1. Weeds that are suppressed by residual activity of VALOR are listed in Table 2.

**Table 4 Emerged Broadleaf Weeds Controlled by Hooded, Shielded and Layby Application of VALOR Tank Mixes With Glyphosate or MSMA in Cotton**

BROADLEAF WEED SPECIES		WEED HEIGHT (inches) 2 oz /A
COMMON NAME	SCIENTIFIC NAME	
Bindweed Field <sup>1</sup>	<i>Convolvulus arvensis</i>	4
Carpetweed	<i>Mollugo verticillata</i>	4
Chickweed Common	<i>Stellaria media</i>	4
Cocklebur Common	<i>Xanthium strumarium</i>	4
Florida Beggarweed	<i>Desmodium tortuosum</i>	2
Hemp Sesbania	<i>Sesbania exaltata</i>	6
Jimsonweed	<i>Datura stramonium</i>	4
Lambsquarters Common	<i>Chenopodium album</i>	4
Morningglories		
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integruscula</i>	4
Ivyleaf	<i>Ipomoea hederacea</i>	4
Pitted	<i>Ipomoea lacunose</i>	4
Red	<i>Ipomoea coccinea</i>	4
Tall	<i>Ipomoea purpurea</i>	2
Mustard Wild	<i>Brassica kaber</i>	6
Nightshades		
Black	<i>Solanum nigrum</i>	4
Eastern Black	<i>Solanum ptycanthum</i>	4
Hairy	<i>Solanum sarrachoides</i>	4
Pigweeds		
Palmer Amaranth	<i>Amaranthus palmeri</i>	4
Redroot	<i>Amaranthus retroflexus</i>	4
Smooth	<i>Amaranthus hybridus</i>	4
Plantain Broadleaf	<i>Plantago major</i>	6
Prickly Sida (Teaweed)	<i>Sida spinosa</i>	4
Purslane Common	<i>Portulaca oleracea</i>	2
Ragweeds		
Common	<i>Ambrosia artemisiifolia</i>	2
Giant	<i>Ambrosia trifida</i>	4
Rice Flatsedge	<i>Cyperus iria</i>	2
Sicklepod	<i>Senna obtusifolia</i>	4
Smartweeds		
Ladysthumb	<i>Polygonum persicaria</i>	4
Pale	<i>Polygonum lapathifolium</i>	4
Pennsylvania	<i>Polygonum pennsylvanicum</i>	4
Spotted Spurge	<i>Euphorbia maculata</i>	4
Velvetleaf	<i>Abutilon theophrasti</i>	4
Venice Mallow	<i>Hibiscus trionum</i>	2
Waterhemp		
Common	<i>Amaranthus rudis</i>	2
Tall	<i>Amaranthus tuberculatus</i>	2

<sup>1</sup> VALOR tank mixes will control the above ground portion of field bindweed. Repeated applications will be needed to control regrowth.

## CARRIER VOLUME AND SPRAY PRESSURE

### Hooded, Shielded and Layby Application

To ensure thorough coverage in hooded shielded and layby applications use 15 to 30 gals spray solution per treated acre Use 20 to 30 gals per treated acre under heavy weed pressure Nozzle selection should meet manufacturer s gallonage and pressure recommendations for application method being used Do not use Flood Jet nozzles as they tend to increase the chance of crop injury

## ADDITIVES

### Hooded, Shielded and Layby Application

Weed control from hooded shielded or layby application of Valor in cotton requires the addition of an agronomically approved non-ionic surfactant to the spray mixture Non ionic surfactant must contain at least 80% active ingredient Mixing compatibility qualities should be verified by a jar test **The use of crop oil concentrates, methylated seed oils, organo silicant surfactants or products containing these ingredients, may result in severe crop injury and should not be used**

## APPLICATION EQUIPMENT

Apply *Valor* tank mixes with ground equipment using standard commercial sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume Application equipment should be clean and in good repair Nozzles should meet manufacturer s recommendations for spray pattern and placement on spray boom and should be checked frequently for accuracy

## TIMING TO COTTON

### Hooded and Shielded Application

*Valor* tank mixes may be applied with a hooded or shielded sprayer after cotton has reached a minimum of 6 inches in height All nozzles must be under the hood or behind the shield to ensure no spray solution comes in contact with the cotton **Care must be taken to ensure the spray solution or drift does not come in contact with the cotton or severe crop injury can occur**

### Layby Application

Layby application of VALOR tank mixes may be made once cotton has reached a minimum of 16 inches in height Cotton that is smaller than 16 inches in height may be injured by VALOR applications VALOR application must be directed to the lower 2 inches of the cotton stem to avoid crop injury

## TIMING TO WEEDS

VALOR tank mix applications must be made to weeds within the height range given in Table 4

## TANK MIXES

VALOR must be tank mixed with one of the herbicides listed in Table 5 for postemergence control of the weeds listed in Table 4

**Table 5 Tank Mixes with VALOR for Hooded, Shielded and/or Layby Use in Cotton**

TANK MIX PARTNER	TARGET WEEDS	HOODED AND SHIELDED	LAYBY
glyphosate	Perennial Grasses and Broadleaves	X	X <sup>1</sup>
MSMA	Annual Grasses Yellow Nutsedge	X	X

<sup>1</sup> For use only in cotton with the ROUNDUP READY gene



## DIRECTIONS FOR USE IN CUCURBIT VEGETABLES (ROW MIDDLES)

**Cucurbit Vegetables (Crop Group 9) including** chayote (fruit) Chinese Waxgourd (Chinese preserving melon) citron melon cucumber gherkin gourd edible (includes hyotan cucuzza hechima Chinese okra) *Momordica* spp (includes balsam apple balsam pear bittermelon Chinese cucumber) muskmelon (includes cantaloupe) pumpkin squash summer squash winter (includes butternut squash calabaza hubbard squash acorn squash spaghetti squash) watermelon

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 4 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 8 oz of *Valor* Herbicide per acre during a single growing season
- Do not use with an adjuvant
- Plants should be grown on raised plastic mulched beds that are higher than the treated row middle
- [Arizona California and Hawaii only** For fallowbed application on transplanted melon beds follow supplemental labeling provided by Valent U S A Corporation ]
- Spray must be directed to the row middle away from the crop bed and with minimal contact with plastic including the sides of the bed If top of mulch beds (where plants are to be transplanted) is contacted severe injury can occur due to foliage contact with treated plastic In this scenario a rainfall event of 1/2 inch (natural or irrigation) must occur prior to transplanting to reduce *Valor* Herbicide residues
- Drift of treated soil particles onto plants may cause contact injury
- Irrigate treated field after application and prior to transplanting with minimum of 1/4 inch of water if rainfall does not occur between application and transplanting
- All applications must be made with hooded or shielded equipment

### TIMING TO CUCURBIT VEGETABLES

Apply *Valor* Herbicide at 4 oz per acre as a hooded or shielded application to row middles up to 14 days prior to transplanting or seeding for preemergence control of the weeds listed in Table 7 Weeds Controlled by Residual Activity of *Valor* Herbicide as well as to assist in the postemergence control of emerged weeds A second application of *Valor* Herbicide at 4 oz per acre may be applied up to 21 days after transplanting or emergence if needed Do not apply during or after bloom

### TIMING TO WEEDS

*Valor* Herbicide may be used for residual weed control as well as to assist in postemergence burndown of many annual and perennial weeds in row middles A registered preemergence grass herbicide may be added for control of additional grassy weeds For assisting in the control of emerged weeds tank mix *Valor* Herbicide with paraquat Aim™ or other registered burndown herbicide Do not tank mix with glyphosate after transplanting Refer to tank mix partner's label for recommended rate and application parameters

**Read tank mix product label for rate and weeds controlled Always read and follow label directions for all tank mix products before using The most restrictive labeling of any tank mix product must be followed *Valor* Herbicide when applied according to label use directions will control the weeds listed in Table 7 Weeds Controlled by Residual Activity of *Valor* Herbicide This label makes no claims concerning control of other weed species**

## DIRECTIONS FOR USE IN DRY BEANS

Dried cultivars of bean (*Lupinus*) bean (*Phaseolus*) (includes field bean kidney bean lima bean (dry) navy bean pinto bean tepary bean) bean (*Vigna*) (includes adzuki bean blackeyed pea catjang cowpea crowder pea moth bean mung bean rice bean southern pea urd bean) broad bean (dry) chickpea guar lablab bean and lentil

**WEED SUPPRESSION** *[Weed Suppression section not to be shown on production label]*

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 2 oz of VALOR per acre during a single application
- Do not apply more than 2 oz of VALOR per acre during a single growing season

**[Arizona California Colorado Hawaii Idaho Nebraska Oregon and Washington only]** For weed suppression in dry beans follow supplemental labeling provided by Valent U S A Corporation ]

**[Arizona California Hawaii Idaho Oregon and Washington only]** For weed suppression in garbanzo beans follow supplemental labeling provided by Valent U S A Corporation ]

**Many weather related factors including high wind splashing or heavy rains or cool conditions at or near crop emergence may result in dry bean injury in fields treated with VALOR. On occasion this has resulted in a delay in maturity. User should assume these risks before using VALOR.**

### TIMING TO DRY BEAN

VALOR may be applied to dry beans within 2 days after planting for the preemergence suppression of the weeds listed in Table 1 Broadleaf Weeds Controlled by Residual Activity of *Valor* Herbicide or Table 8 Weeds Suppressed by Residual Activity of VALOR. VALOR should be tank mixed with other labeled herbicides for broad spectrum weed control.

### TIMING TO WEEDS

VALOR may be applied to dry beans prior to planting or preemergence (after planting). Preemergence application of VALOR must be made within 2 days after planting and prior to dry bean emergence. Application after the dry beans have begun to crack or are emerged will result in severe crop injury. To avoid severe crop injury, do not apply to dry beans after beans begin to crack or have emerged. Preplant incorporation (PPI) applications may result in reduced weed control.

### ADDITIONAL RESIDUAL GRASS CONTROL

VALOR can be tank mixed with pendimethalin for additional grass control.

### HARVEST AID

#### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 3 oz of VALOR per acre during a single application
- Do not apply more than 3 oz of VALOR per acre during a single growing season
- Do not harvest within 5 days of application

Desiccation from VALOR requires the addition of an agronomically approved adjuvant to the spray mixture. A methylated seed oil which contains at least 15% emulsifiers and 80% oil at 2% v/v should be used. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with either a crop oil concentrate or methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for a crop oil concentrate or a methylated seed oil. Tank mixing VALOR with glyphosate or paraquat will increase control of emerged weeds and aid in harvest. Add a burndown tank mix partner for the control of emerged weeds labeled for dry bean in accordance with the most restrictive labeled limitations and precautions.

### TIMING TO DRY BEANS

Apply when crop is mature and at least 80% of the pods are yellowing and mostly ripe with no more than 40% (bush type beans) or 30% (vine type beans) of the leaves still green in color. Dry beans can be harvested 5 days after application. To ensure thorough coverage use 15 to 30 gallons spray solution per acre. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for postemergence application.

## DIRECTIONS FOR USE IN FIELD CORN

[For Use in the States of Arizona California and Hawaii Only]

### RESTRICTIONS AND LIMITATIONS

- Use only on no till or minimum tillage fields where last years crop residue has not been incorporated into the soil
- Corn must be planted between 14 and 30 days after application unless the application is made as part of a Fall burndown program
- Corn can be planted 7 days after an application of 2 oz/A if a minimum of 25% of the soil surface is covered with the residue of the preceding crop and a minimum of 1/4 inch of rainfall has occurred between application and planting
- Do not apply more than [2 oz] 3 oz of *Valor* per acre during a single growing season
- Do not irrigate between emergence and 2 leaf corn
- Do not use on popcorn sweet corn or corn grown for seed

### TIMING TO FIELD CORN

- Apply *Valor* Herbicide at 2 to 3 oz/A between 7 and 30 days prior to planting field corn for the preemergence control of the weeds listed in Table 1 Broadleaf Weeds Controlled by Residual Activity of *Valor* Herbicide
- Apply *Valor* Herbicide at 2 oz/A between 7 and 30 days prior to planting field corn if a minimum of 25% of the soil surface is covered with the residue of the preceding crop and a minimum of 1/4 inch of rainfall has occurred between application and planting
- Apply *Valor* Herbicide at 3 oz/A between 14 and 30 days prior to planting field corn

### Burndown Use Directions – For Preplant Applications in Field Corn

*Valor* Herbicide applied as part of a burndown program may be used for residual weed control as well as to assist in postemergence burndown of many weeds where field corn will be planted directly into the residue of the previous year See Directions for Use in Fall and Spring Preplant Burndown and Fallow Seedbed Programs in Field Corn Peanut and Soybean for rates and timing of applications For control of emerged weeds *Valor* Herbicide must be applied with an appropriate burndown tank mix partner listed in Table 6 To ensure thorough coverage use a minimum of 15 gallons of spray solution per acre Refer to tank mix partner's label for recommended application pressure and recommended adjuvant systems

#### INCREASING SPEED OF GLYPHOSATE BURNDOWN ACTIVITY

*Valor* Herbicide at 1 oz/A may be tank mixed with glyphosate (Roundup®) to increase the speed of burndown activity compared to glyphosate applied alone Residual weed control will not be provided at rates lower than 2 oz/A however suppression of the weeds in Table 2 may occur at *Valor* Herbicide rates as low as 1 oz/A Applications of *Valor* Herbicide at 1 oz/A must be made a minimum of 14 days prior to planting field corn

### TANK MIXES

*Valor* Herbicide may be tank mixed with the herbicides listed in Table 6 for pre plant burndown applications Refer to tank mix partner's label for adjuvant recommendations

**Table 6 Tank Mix Partners for Burndown and/or Residual Control of Weeds in Field Corn**

TANK MIX PARTNERS <sup>1</sup>	
2 4 D LVE	metribuzin
atrazine	paraquat
Basis®	Python®
dicamba	Resolve®
Express®	simazine
glyphosate	Weedmaster®
Hornet®	

<sup>1</sup>Refer to tank mix product labels for specific recommendations

**TANK MIX RESTRICTIONS**

Tank mixes with flufenacet (Axiom or Domain) metolachlor or s metolachlor (Dual Magnum or Dual II Magnum) dimethenamid or dimethenamid p (Frontier or Outlook) alachlor (Lasso) or acetochlor (Surpass or Harness) may result in injury to field corn when application is followed by prolonged periods of cool wet weather and should not be used with *Valor* Herbicide unless supplemental labeling provided by Valent U S A Corporation is followed

**DIRECTIONS FOR USE IN FIELD PEAS****WEED CONTROL****RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 2 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 2 oz of *Valor* Herbicide per acre during a single growing season
- [For use in Idaho Montana Oregon and Washington only ]

**Many weather related factors including high wind splashing or heavy rains or cool conditions at or near crop emergence may result in pea injury in fields treated with *Valor* Herbicide On occasion this has resulted in a delay in maturity User should assume these risks before using *Valor* Herbicide**

**TIMING TO FIELD PEAS**

*Valor* Herbicide may be applied to field peas within 2 days after planting for the preemergence control of the weeds listed in Table 1 Broadleaf Weeds Controlled by Residual Activity of VALOR or Table 8 Weeds Suppressed by Residual Activity of VALOR Tank mix *Valor* Herbicide with other labeled herbicides for broad spectrum weed control

**TIMING TO WEEDS**

*Valor* Herbicide may be applied to field peas prior to planting or preemergence (after planting) Preemergence application of *Valor* Herbicide must be made within 2 days after planting and prior to field pea emergence To avoid severe crop injury do not apply to field peas after peas begin to crack or have emerged

Preplant incorporation (PPI) applications may result in reduced weed control

**ADDITIONAL RESIDUAL GRASS CONTROL**

*Valor* Herbicide can be tank mixed with pendimethalin for additional grass control

**HARVEST AID****RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season
- Do not harvest within 5 days of application

Desiccation from *Valor* Herbicide requires the addition of an agronomically approved adjuvant to the spray mixture A methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A should be used A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation The addition of a nitrogen source does not replace the need for methylated seed oil Tank mixing *Valor* Herbicide with glyphosate will increase control of emerged weeds and aid in harvest

**TIMING TO FIELD PEAS**

Apply *Valor* Herbicide at 1.5 to 2 oz/A when crop is physiologically mature and a minimum of 80% of the pods are yellow to tan in color and 20% are yellow in color If field peas are treated too early a reduction in seed quality may occur Do not spray *Valor* Herbicide on any area of the field with a significant amount of plants with green color Peas can be harvested 5 days after application

To ensure thorough coverage use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure recommendations for postemergence application

## DIRECTIONS FOR USE IN FLAX

### **HARVEST AID**

#### **RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season
- Do not harvest within 5 days of application

Desiccation from *Valor* Herbicide requires the addition of an agronomically approved adjuvant to the spray mixture. A methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A should be used. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil.

### **TIMING TO FLAX**

Apply *Valor* Herbicide at 1.5 to 2 oz/A when crop is physiologically mature and at least 75% of the bolls are brown in color. Flax can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure recommendations for postemergence application.

## DIRECTIONS FOR USE IN FRUITING VEGETABLES (INCLUDING OKRA) ROW MIDDLES

Eggplant, Groundcherry (*Physalis* spp.), Okra, Pepino, Peppers (including Bell Pepper, Chili Pepper, Cooking Pepper, Pimento, Sweet Pepper), Tomatillo and Tomato.

### **RESTRICTIONS AND LIMITATIONS**

Do not apply more than 4 oz of VALOR per acre during a single application.

Do not apply more than 8 oz of VALOR per acre during a single growing season.

Plants should be grown on raised or plastic mulched beds that are higher than the treated row middle.

[**Arizona, California and Hawaii only** For fallowbed application on transplanted peppers and tomato beds, follow supplemental labeling provided by Valent U.S.A. Corporation.]

Spray must be directed to the row middle, away from the crop bed and with minimal contact with plastic, including the sides of the bed. If top of mulch beds (where plants are to be transplanted) is contacted, severe injury can occur due to foliage contact with treated plastic. In this scenario, a rainfall event of 1/2 inch (natural or irrigation) must occur prior to transplanting to reduce VALOR residues.

– Drift of treated soil particles onto plants may cause contact injury.

Irrigate treated field after application and prior to transplanting with minimum of 1/4 inch of water if rainfall does not occur between application and transplanting.

All applications must be made with hooded or shielded equipment.

### **TIMING TO FRUITING VEGETABLES**

Apply VALOR at 4 oz per acre as a hooded or shielded application to row middles up to 14 days prior to transplanting or seeding for preemergence control of the weeds listed in Table 7. Weeds Controlled by Residual Activity of *Valor* Herbicide, as well as to assist in the postemergence control of emerged weeds. A second application of VALOR at 4 oz per acre may be applied up to 21 days after transplanting or emergence if needed. Do not apply during or after bloom.

### **TIMING TO WEEDS**

VALOR may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds in row middles. A registered preemergence grass herbicide may be added for control of additional grassy weeds. For assisting in the control of emerged weeds, tank mix VALOR with paraquat, Aim™, or other registered burndown herbicide. Do not tank mix with glyphosate after transplanting or crop emergence. Refer to tank mix partner's label for recommended rate and application parameters.

## DIRECTIONS FOR USE IN GARLIC

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 6 oz of VALOR per acre during a single application
- Do not apply more than 6 oz of VALOR per acre during a single growing season

### TIMING TO GARLIC

VALOR may be applied at 6 oz /A to garlic prior to garlic emergence. Application should be made within 3 days after planting garlic.

### TIMING TO WEEDS

#### Preemergence – Preemergence To Weeds

Apply VALOR to weed free garlic for preemergence control of the weeds listed in Table 10 Weeds Controlled by Preemergence Application of VALOR

## DIRECTIONS FOR USE IN HOPS

[Not For Use in California or New York]

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 6 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 6 oz of *Valor* Herbicide per acre during a single growing season
- Do not allow spray to contact green stem (Unless used for sucker control) foliage flowers or cones or unacceptable injury may occur
- Do not apply within 30 days of harvest
- Do not use with an adjuvant

*Valor* Herbicide can be used in hops for preemergence weed control as well as sucker control

### TIMING TO HOPS FOR SUCKER CONTROL

Apply *Valor* Herbicide at 6 oz/A as a directed application after hops have reached a minimum of 6 feet in height for sucker control. Application should be directed to the lower 2 feet of the hops.

### TIMING TO HOPS FOR PREEMERGENCE WEED CONTROL

Apply *Valor* Herbicide at 6 oz/A as a 1 to 1.5 foot band to each side of the hop row to dormant hops January thru March to ensure time for rain incorporation and activation. If weeds are emerged at the time of application, tank mix *Valor* Herbicide with a labeled burndown herbicide such as paraquat or glyphosate to assist with control of emerged weeds. Do not mow or rake over treated areas as dust created by mowing may drift onto sensitive crops or vegetation resulting in injury.

### TIMING TO WEEDS

*Valor* Herbicide applications must be made prior to weed emergence for control of weeds listed in Table 10 Weeds Controlled by Preemergence Application of *Valor* Herbicide

**Read tank mix product label for rates and weeds controlled. Always read and follow label directions for all tank mix products before using. The most restrictive labeling of any tank mix product must be followed. *Valor* Herbicide when applied according to label use directions will control the weeds listed in Table 10 Weeds Controlled by Preemergence Application of *Valor* Herbicide. This label makes no claims concerning control of other weed species.**

## DIRECTIONS FOR USE IN LENTILS

### **HARVEST AID**

#### **RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season
- Do not harvest within 5 days of application

Desiccation from *Valor* Herbicide requires the addition of an agronomically approved adjuvant to the spray mixture. A methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A should be used. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* Herbicide with glyphosate or paraquat will increase control of emerged weeds and aid in harvest.

### **TIMING TO LENTILS**

Apply *Valor* Herbicide at 1.5 to 2 oz/A when crop is physiologically mature and a minimum of 80% of the pods are yellow to tan in color and 20% are yellow in color. If lentils are treated too early, a reduction in seed quality may occur. Do not spray *Valor* Herbicide on any area of the field with a significant amount of plants with green color. Lentils can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure recommendations for postemergence application.

## DIRECTIONS FOR USE IN MINT (Peppermint and Spearmint)

### **RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 4 oz of VALOR per acre during a single application
- Do not apply more than 8 oz of VALOR per acre during a single growing season
- Do not make a sequential VALOR application within 60 days of the first VALOR application
- Apply only to dormant mint. Application to non-dormant mint may result in unacceptable crop injury.
- Do not apply within 80 days of harvest
- Do not apply to row or baby mint. Use only on established meadow mint.
- Do not apply to mint that has been weakened by diseases, insects (example: mint root borer), nematodes, drought, soil salts, high soil pH, previous pesticides, winter injury, or double cutting, as severe injury may occur. Apply only to healthy, vigorous mint with undamaged rhizomes.
- Do not apply before November 25 or after March 1
- Do not apply a Fall application if roots and rhizomes are weak, thin, or damaged
- Do not apply to stands established longer than 3 years
- Do not apply VALOR on mint in Southern Union County (south of Ladd Canyon) or Baker County in Oregon

**Many weather-related factors, including high wind, splashing, or heavy rains, or cool conditions at or near mint emergence, may result in mint injury in fields treated with VALOR. User should assume these risks before using VALOR.**

Tank mixes with labeled rates of paraquat are recommended to control emerged weeds and increase crop safety.

### **TIMING TO MINT**

As a spray, VALOR may be applied only to established dormant mint for preemergence control of the weeds listed in Table 7, as well as to assist in the postemergence control of emerged weeds. Application to non-dormant mint or to baby (row) mint (time from planting of mint roots through the first cutting) may result in unacceptable crop injury. As a bulk fertilizer application, VALOR may be applied at least 80 days prior to harvest. Leaves must be dry at the time of applications or severe injury may occur.

## **TIMING TO WEEDS**

### **Burndown – Dormant Mint Postemergence To Weeds**

VALOR may be used for residual weed control as well as to assist in postemergence burndown of many annual and perennial weeds where established mint is dormant. For control of emerged weeds, tank mix VALOR with paraquat. Refer to paraquat label for recommended rate and application parameters. To ensure thorough coverage, use a minimum of 15 gals. of spray solution per acre. VALOR tank mixes applied to assist in the control of emerged weeds must be applied with a non ionic surfactant at 0.25% v/v. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs /A or 28 to 32% nitrogen solution at 1 to 2 qts /A) may be added to increase herbicidal activity.

### **Preemergence – Dormant Mint Preemergence To Weeds**

Apply VALOR to dormant mint for the preemergence control of weeds listed in Table 7. Fall applications of VALOR followed by a sequential application in the Spring have resulted in better Summer annual weed control than a single Fall or single Spring application.

Fall application is most effective for Fall germinating weeds such as groundsel. Fields plowed or harrowed after a VALOR application will result in less effective preemergence activity. In furrow irrigated fields, corrugating that is done after a VALOR application will expose untreated soil and break the herbicide barrier resulting in poor weed control.



Table 7 Weeds Controlled by Residual Activity of VALOR

BROADLEAF WEED SPECIES				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	VALOR RATE
Bristly Starbur	<i>Acanthospermum hispidum</i>	Up to 5%	All Soil Types	4 oz /A
Carpetweed	<i>Mollugo verticillata</i>			
Chickweeds				
Common	<i>Stellaria media</i>			
Mouseear	<i>Cerastium vulgatum</i>			
Coffee Senna	<i>Cassia occidentalis</i>			
Copperleaf Hophornbeam	<i>Acalypha ostryifolia</i>			
Dandelion	<i>Taraxacum officinale</i>			
Dodder (suppression only) <sup>1</sup>	<i>Cuscuta</i> spp			
Eclipta	<i>Eclipta prostrata</i>			
Evening Primrose Cutleaf	<i>Oenothera laciniata</i>			
False Chamomile	<i>Tripleurospermum maritima</i>			
Fiddleneck Coast	<i>Amsinckia menziesii</i>			
Field Pennycress	<i>Thlaspi arvense</i>			
Fleabane Hairy	<i>Conyza bonariensis</i>			
Flixweed	<i>Descurainia spophia</i>			
Florida Beggarweed	<i>Desmodium tortuosum</i>			
Florida Pusley	<i>Richardia scabra</i>			
Golden Crownbeard	<i>Verbesina encelioides</i>			
Groundsel Common	<i>Senecio vulgaris</i>			
Hairy Indigo	<i>Indigofera hirsuta</i>			
Hemp Sesbania	<i>Sesbania exaltata</i>			
Henbit	<i>Lamium amplexicaule</i>			
Jimsonweed	<i>Datura stramonium</i>			
Kochia	<i>Kochia scoparia</i>			
Lambsquarters Common	<i>Chenopodium album</i>			
Little Mallow	<i>Malva parviflora</i>			
London Rocket	<i>Sisymbrium irio</i>			
Marestail/Horseweed	<i>Conyza canadensis</i>			
Mayweed/False Chamomile	<i>Matricaria inodora</i>			

continued

<sup>1</sup>Chateau WDG Herbicide at 4 oz/A will provide postemergence dodder suppression when applied in combination with Pursuit Herbicide or Raptor Herbicide at labeled rates. The use of Pursuit Herbicide and Raptor Herbicide require the use of a NIS which will result in burn and stunting of alfalfa. Growers should expect and accept this prior to using this tank mix.

Table 7 Weeds Controlled by Residual Activity of VALOR (continued)

BROADLEAF WEED SPECIES				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	VALOR RATE
Morningglories		Up to 5%	All Soil Types	4 oz /A
Entireleaf	<i>Ipomoea hederacea</i> var <i>integruscula</i>			
Ivyleaf	<i>Ipomoea hederacea</i>			
Red/Scarlet	<i>Ipomoea coccinea</i>			
Smallflower	<i>Jacquemontia tamnifolia</i>			
Tall	<i>Ipomoea purpurea</i>			
Mustard				
Tansy	<i>Descurainia pinnata</i>			
Tumble	<i>Sisymbrium altissimum</i>			
Wild	<i>Brassica kaber</i>			
Nettle Burning	<i>Urtica urens</i>			
Nightshades				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Hairy	<i>Solanum sarrachoides</i>			
Pigweeds				
Palmer Amaranth	<i>Amaranthus palmeri</i>			
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Lettuce (China Lettuce)	<i>Lactuca scariola</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Sowthistle Prickly	<i>Sonchus asper</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane				
Common	<i>Portulaca oleracea</i>			
Horse	<i>Trianthema portulacastrum</i>			
Radish Wild	<i>Raphanus raphanistrum</i>			
Ragweed Common	<i>Ambrosia artemisiifolia</i>			
Redmaids	<i>Calandrinia ciliata</i> var <i>menziesii</i>			
Russian Thistle	<i>Salsola iberica</i>			
Shepherd s purse	<i>Capsella bursa pastoris</i>			
Smartweeds				
Ladysthumb	<i>Polygonum persicaria</i>			
Pennsylvania	<i>Polygonum pennsylvanicum</i>			
Smellmelon	<i>Cucumis melo</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Spurred Anoda	<i>Anoda cristata</i>			
Tropic Croton	<i>Croton glandulosus</i>			
Velvetleaf	<i>Abutilon theophrasti</i>			
Venice Mallow	<i>Hibiscus trionum</i>			
Waterhemp				
Common	<i>Amaranthus rudis</i>			
Tall	<i>Amaranthus tuberculatus</i>			
White Cockle	<i>Silene latifolia</i>			
Wild Poinsettia	<i>Euphorbia heterophylla</i>			
Wormwood Biennial	<i>Artemisia biennis</i>			
Yellow Rocket	<i>Barbarea vulgaris</i>			

continued

**Table 7 Weeds Controlled by Residual Activity of VALOR (continued)**

<b>BROADLEAF WEED SPECIES</b>				
<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>	<b>ORGANIC MATTER</b>	<b>SOIL TYPE</b>	<b>VALOR RATE</b>
<b>GRASS WEED SPECIES</b>		Up to 5%	All Soil Types	4 oz /A
Barnyardgrass	<i>Echinochloa crus galli</i>			
Bluegrass Annual	<i>Poa annua</i>			
Crabgrass Large	<i>Digitaria sanguinalis</i>			
Foxtail Giant	<i>Setaria faberi</i>			
Goosegrass	<i>Eleusine indica</i>			
Lovegrass California	<i>Eragrostis diffusa</i>			
Panicums				
Fall	<i>Panicum dichotomiflorum</i>			
Texas	<i>Panicum texanum</i>			
Ryegrass Italian	<i>Lolium multiflorum</i>			
Signalgrass Broadleaf	<i>Bracharia platyphylla</i>			

### **DIRECTIONS FOR USE IN ONION (DRY BULB)**

[For Use in the States of Michigan New York and North Dakota Only]

[For chemigation application on onion (dry bulb) follow supplemental labeling provided by Valent U S A Corporation ]

### **RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 2 oz of VALOR per acre during a single application
- Do not apply more than 3 oz of VALOR per acre during a single growing season
- Do not make sequential application within 14 days of the first application
- Do not apply more than 1 oz of VALOR per season on soils that contain greater than 90% sand plus gravel
- Do not apply as part of a tank mix other than with Prowl® H<sub>2</sub>O or unacceptable injury may result Other formulations of pendimethalin should not be tank mixed with VALOR for use in onions
- Do not apply with any type of adjuvant
- Do not apply within 45 days of harvest

**Use of VALOR may result in necrotic spotting of onion leaves that come in contact with the spray User should assume this potential crop response before using VALOR**

### **[Microrate Application]**

Sequential applications of VALOR may be applied to onions (dry bulb) between the 2 leaf and 6 leaf stage at rates of 0.5 to 1 oz /A on a 7 day interval ]

### **TIMING TO ONIONS (dry bulb)**

Apply VALOR to transplanted onions (dry bulb) between the 2 leaf and 6 leaf stage and on direct seed onions (dry bulb) between the 3 leaf and 6 leaf stage

### **TIMING TO WEEDS**

#### **Preemergence – Emerged Onions (dry bulb) Preemergence To Weeds**

Apply VALOR to weed free onions (dry bulb) for preemergence control of the weeds listed in Table 1 Section A

## DIRECTIONS FOR USE IN PEANUT

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 3 oz of VALOR per acre during a single growing season
- Do not apply more than 2 oz /A in the states of North Carolina Oklahoma or Virginia where climatic conditions may result in unacceptable injury to peanuts unless supplemental labeling provided by Valent U S A Corporation is followed
- Do not irrigate when peanuts are cracking
- Do not graze treated fields or feed treated hay to livestock

**Many weather related factors including high wind splashing or heavy rains or cool conditions at or near peanut emergence may result in peanut injury in fields treated with VALOR On occasion this has resulted in a delay in maturity or even a slight decrease in yield**

### WIND MANAGEMENT

In areas where shallow cultivation is used between rows to reduce wind borne sand damage to peanuts weed control from VALOR may be reduced

### TIMING TO PEANUTS

**VALOR may be applied to peanuts prior to planting or preemergence (after planting) Preemergence applications of VALOR must be made within 2 days after planting and prior to peanut emergence Application after the peanuts have begun to crack or are emerged will result in severe crop injury Application should not be made when peanuts have begun to crack Select VALOR rate from Table 1 according to anticipated weed spectrum**

### TIMING TO WEEDS

#### Burndown – Preemergence to Peanuts Postemergence to Weeds

VALOR applied as part of a burndown program may be used for residual weed control as well as to assist in postemergence burndown of many annual and perennial weeds where peanuts will be planted directly into a stale seedbed cover crop or in previous crop residues Apply VALOR before planting during planting or after planting but before the crop emerges For control of emerged weeds tank mix VALOR with glyphosate Refer to glyphosate label for recommended rate and application pressure To ensure thorough coverage use a minimum of 15 gals of spray solution per acre VALOR tank mixes applied to assist in the control of emerged weeds must be applied with an adjuvant such as a non ionic surfactant at 0.25% v/v or a crop oil concentrate or a methylated seed oil at 1 to 2 pt /A A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs /A or 28 to 32% nitrogen solution at 1 to 2 qts /A) may be added to increase herbicidal activity

Preemergence (conventional tillage) applications of VALOR must be applied prior to weed emergence

### ADDITIONAL RESIDUAL GRASS CONTROL SEQUENTIAL

VALOR may be applied sequentially following a preplant incorporated application of trifluralin (states of New Mexico Oklahoma and Texas only) SONALAN® DUAL® (metolachlor) pendimethalin or FRONTIER®

### ADDITIONAL RESIDUAL GRASS CONTROL TANK MIXED

VALOR can be tank mixed with alachlor metolachlor or FRONTIER for additional grass and broadleaf weed control VALOR can also be tank mixed with pendimethalin or SONALAN in states where they are labeled provided overhead irrigation guidelines on the pendimethalin and/or SONALAN labels are followed

## DIRECTIONS FOR USE IN POTATO

[Arizona California Colorado Delaware Florida Hawaii Idaho Maryland Minnesota Montana Nebraska Nevada New Jersey New Mexico North Carolina North Dakota Oregon South Dakota Texas Utah Virginia Washington Washington DC and Wyoming only ] [For chemigation application on potato follow supplemental labeling provided by Valent U S A Corporation ]

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 1.5 oz of VALOR per acre during a single application
- Do not apply more than 1.5 oz of VALOR per acre during a single growing season
- Do not apply to Rill (Furrow) irrigated potatoes

**Many weather related factors including high wind, splashing or heavy rains or cool conditions at or near potato emergence may result in potato injury in fields treated with VALOR. On occasion this has resulted in a delay in maturity. User should assume these risks before using VALOR.**

### TIMING TO POTATOES

VALOR may be applied to potatoes after hilling for the preemergence suppression of the weeds listed in Table 8. VALOR should be tank mixed with other labeled herbicides for broad spectrum weed control. A minimum of 2 inches of settled soil must cover the vegetative portion of the potato plant at the time of VALOR application. Application to potatoes with less than 2 inches of soil covering the vegetative portion of the potato may result in crop injury. In areas with historically higher amounts of rainfall during the time of preemergence herbicide applications, such as the Red River Valley, Minnesota and North Dakota, the requirement for 2 inches of settled soil is critical to avoid crop injury. Mechanical incorporation of VALOR will result in decreased weed control and should be avoided. In areas with sprinkler irrigation, VALOR should be incorporated with 0.5 to 0.75 inches of irrigation after application and before any sprouts are within 2 inches of the settled soil surface if a rainfall event has not yet occurred.

### TIMING TO WEEDS

#### Preemergence – Soil Covered Potatoes Preemergence To Weeds

Apply VALOR to soil covered potatoes for the preemergence suppression of the weeds listed in Table 8. Harrowing, cultivation or corrugating after VALOR application will reduce weed control.

**Table 8 Weeds Suppressed by Residual Activity of VALOR at 1.5 oz/A**

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	VALOR RATE
Lambsquarters Common	<i>Chenopodium album</i>	Up to 5%	1.5 oz /A
Mustard Wild	<i>Brassica kaber</i>		
Nightshades			
Black	<i>Solanum nigrum</i>		
Eastern Black	<i>Solanum ptycanthum</i>		
Hairy	<i>Solanum sarrachoides</i>		
Pigweeds			
Palmer Amaranth	<i>Amaranthus palmeri</i>		
Redroot	<i>Amaranthus retroflexus</i>		
Smooth	<i>Amaranthus hybridus</i>		
Spiny Amaranth	<i>Amaranthus spinosus</i>		
Tumble	<i>Amaranthus albus</i>		
Prickly Lettuce (China Lettuce)	<i>Lactuca sermola</i>		
Radish Wild	<i>Raphanus raphanistrum</i>		

## DIRECTIONS FOR USE IN SOYBEAN

### RESTRICTIONS AND LIMITATIONS

Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season  
 Do not tank mix *Valor* Herbicide with acetochlor (Warrant®) alachlor (Micro Tech®) flufenacet (Axiom® Domain®) metolachlor (Dual® Magnum Dual® II Magnum Boundary®) or dimethenamid (Frontier® or Outlook®) within 14 days of planting soybeans unless soybeans are planted under no till or minimum tillage conditions on wheat stubble or no till field corn stubble Do not irrigate when soybeans are cracking  
 Do not graze treated fields or feed treated hay to livestock

### TIMING TO SOYBEANS

**VALOR** may be applied to soybeans prior to planting or preemergence (after planting) Preemergence application of **VALOR** must be made within 3 days after planting and prior to soybean emergence Application after the soybeans have begun to crack or are emerged will result in severe crop injury Application should not be made when soybeans have begun to crack Select **VALOR** rate from Table 1 according to anticipated weed spectrum

### TIMING TO WEEDS

#### Burndown – Preemergence to Soybeans Postemergence to Weeds

**VALOR** applied as part of a burndown program may be used for residual weed control as well as to assist in postemergence burndown of many annual and perennial weeds where soybeans will be planted directly into a stale seedbed cover crop or in previous crop residues For control of emerged weeds choose the most appropriate tank mix partner from Table 9 Apply **VALOR** with ground equipment before planting during planting or within 3 days after planting **but before the crop emerges** To ensure thorough coverage use a minimum of 15 gals of spray solution per acre Refer to tank mix partner's label for recommended application pressure All **VALOR** tank mixes applied to assist in the control of emerged weeds must be applied with crop oil concentrate or methylated seed oil at 1 to 2 pt /A or a non ionic surfactant at 0.25% v/v

#### INCREASING SPEED OF GLYPHOSATE BURNDOWN ACTIVITY

**VALOR** at rates as low as 1 oz /A may be tank mixed with glyphosate (ROUNDUP®) to increase the speed of burndown activity compared to glyphosate applied alone Residual weed control will not be provided at rates lower than 2 oz /A however suppression of the weeds in Table 2 may occur at **VALOR** rates as low as 1 oz /A

### TANK MIXES

**VALOR** may be tank mixed with the herbicides listed in Table 9 for increased burndown activity additional residual broadleaf and/or additional grass control Refer to tank mix partner's label for adjuvant recommendations

**Table 9 Tank Mix Partners for Control of Emerged Weeds in Reduced Tillage Soybeans**

TANK MIX PARTNER	TARGET WEEDS <sup>1</sup>
2,4-D LVE	Marestail Giant Ragweed Dandelion
paraquat	Annual Grasses Henbit
glyphosate	General Burndown
Select Max®	Annual Grasses
SCEPTER® 70 DG	Cocklebur Common Sunflower
Weedmaster®	Marestail Giant Ragweed Dandelion

<sup>1</sup>Refer to tank mix product labels for specific recommendations for control of emerged weeds present

**ADDITIONAL RESIDUAL BROADLEAF CONTROL**

VALOR can be tank mixed with metribuzin FIRSTRATE® LOROX® PURSUIT PLUS® PYTHON® SQUADRON® SCEPTER or STEEL® for additional broadleaf control

**ADDITIONAL RESIDUAL GRASS CONTROL**

VALOR can be tank mixed with pendimethalin or COMMAND® for additional grass control Tank mixes with flufenacet (AXIOM or DOMAIN) metolachlor (DUAL products or BOUNDARY) dimethenamid (FRONTIER or OUTLOOK) or alachlor (MICRO TECH or IntRRo®) may result in severe injury to soybeans when application is followed by prolonged periods of cool wet weather and should not be used with VALOR unless supplemental labeling provided by Valent U S A Corporation is followed

**ROUNDUP READY PROGRAM**

VALOR may be applied as part of a burndown program or preemergence in conventional tillage programs at 2 to 3 oz /A to reduce early season weed competition from waterhemp velvetleaf nightshade and morningglories as well as other weeds listed in Tables 2 and 3 in ROUNDUP READY programs A sequential post emergence application of glyphosate will be required to control weeds not controlled by VALOR

**DIRECTIONS FOR USE IN STRAWBERRY****RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 3 oz of VALOR per acre per application
- Do not apply more than 3 oz of VALOR per acre during a single growing season
- VALOR at 3 oz per acre can be applied to the soil a minimum of 30 days prior to transplanting strawberries provided the strawberries will be transplanted through a plastic mulch
- VALOR at 3 oz per acre can be applied to dormant (established or newly planted) strawberries for the preemergence control of the weeds listed in Table 1 Broadleaf Weeds Controlled by Residual Activity of VALOR
- VALOR at 3 oz per acre can be applied in strawberry row middles with a shielded or hooded sprayer for the preemergence control of the weeds listed in Table 1 Broadleaf Weeds Controlled by Residual Activity of VALOR

Application Method	Minimum Time From Application to Harvest (PHI)	Use Rate Per Acre Per Application (oz)	Use Rate Per Acre Per Year (oz)	Special Use Instructions
Pre transplant	Not applicable	3	3	Apply a minimum of 30 days prior to transplanting and prior to plastic mulch being laid  Apply as part of a tank mix to control emerged weeds
Preemergence to dormant strawberries	Not applicable	3	3	Crop oil concentrate at 1% v/v or non ionic surfactant at 0.25% v/v may be added to help control emerged broadleaf weeds
Hooded or shielded sprayer application to row middles	Do not apply after fruit set	3	3	<b>Apply only to row middles do not apply over strawberries</b>  Apply prior to weed emergence  Crop spotting may occur if an adjuvant is added  <b>Application after fruit set may result in spotting of fruit and should be avoided</b>  Do not allow spray drift to come in contact with fruit or foliage

Table 10 Weeds Controlled by Preemergence Application of Valor Herbicide

BROADLEAF WEED SPECIES				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	VALOR HERBICIDE RATE
Bristly Starbur	<i>Acanthospermum hispidum</i>	Up to 10% <sup>1</sup>	All Soil Types <sup>2</sup>	Asparagus Garlic Hops 6 oz/A
Carpetweed	<i>Mollugo verticillata</i>			
Chickweeds				
Common	<i>Stellaria media</i>			Sugarcane 6 to 8 oz /A
Mouseear	<i>Cerastium vulgatum</i>			
Coffee Senna	<i>Cassia occidentalis</i>			
Dandelion	<i>Taraxacum officinale</i>			Bushberries Grapes Nut Trees
Eclipta	<i>Eclipta prostrata</i>			
Eveningprimrose Cutleaf	<i>Oenothera laciniata</i>			
False Chamomile	<i>Tripleurospermum maritima</i>			(Including Pistachio)
Filaree				
Redstem	<i>Erodium cicutarium</i>			
Whitestem	<i>Erodium moschatum</i>			Pome Fruit Stone Fruit and Non Bearing Fruit Trees
Fiddleneck Coast	<i>Amsinckia menziesii</i>			
Fleabane Hairy	<i>Conyza bonariensis</i>			
Field Pennycress	<i>Thlaspi arvense</i>			6 to 12 oz/A <sup>2</sup>
Florida Beggarweed	<i>Desmodium tortuosum</i>			To Maintain Bare Ground on Non Crop Areas of Farms
Florida Pusley	<i>Richardia scabra</i>			
Golden Crownbeard	<i>Verbesina encelioides</i>			
Groundsel Common	<i>Senecio vulgaris</i>			Orchards & Vineyards
Hairy Indigo	<i>Indigofera hirsuta</i>			
Hemp Sesbania	<i>Sesbania exaltata</i>			
Henbit	<i>Lamium amplexicaule</i>			6 to 12 oz/A
Jimsonweed	<i>Datura stramonium</i>			
Kochia	<i>Kochia scoparia</i>			
Lambsquarters Common	<i>Chenopodium album</i>			
Mallow				
Common (Cheeseweed)	<i>Malva neglecta</i>			
Little	<i>Malva parviflora</i>			
Horseweed/Marestail	<i>Conyza canadensis</i>			
Mayweed/False Chamomile	<i>Matricaria maritima</i>			
Morningglories				
Entireleaf	<i>Ipomoea hederacea</i> var <i>integruscula</i>			
Ivyleaf	<i>Ipomoea hederacea</i>			
Red/Scarlet	<i>Ipomoea coccinea</i>			
Smallflower	<i>Jacquemontia tamnifolia</i>			
Tall	<i>Ipomoea purpurea</i>			

continued

<sup>1</sup> Valor Herbicide can be used on soils with greater than 10% organic matter however length of residual control may be shorter than on soils with lower organic matter content

<sup>2</sup> A maximum Valor Herbicide rate of 6 oz /A per application should be used on any soil that has a sand plus gravel content over 80% if bushes trees or vines are under 3 years of age



Table 10 Weeds Controlled by Preemergence Application of *Valor* Herbicide (continued)

BROADLEAF WEED SPECIES				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	VALOR HERBICIDE RATE
Mustards		Up to 10% <sup>1</sup>	All Soil Types <sup>2</sup>	Asparagus Garlic Hops 6 oz/A  Sugarcane 6 to 8 oz /A  Bushberries Grapes Nut Trees (Including Pistachio) Pome Fruit Stone Fruit and Non Bearing Fruit Trees 6 to 12 oz/A <sup>2</sup>  To Maintain Bare Ground on Non Crop Areas of Farms Orchards & Vineyards 6 to 12 oz/A
London Rocket	<i>Sisymbrium ideo</i>			
Tansey	<i>Desurainia pinnata</i>			
Tumble	<i>Sisymbrium altissimum</i>			
Wild	<i>Brassica kaber</i>			
Nettle Burning	<i>Urtica urens</i>			
Nightshades				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Hairy	<i>Solanum sarrachoides</i>			
Pigweeds				
Palmer Amaranth	<i>Amaranthus palmeri</i>			
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Lettuce (China Lettuce)	<i>Lactuca serriola</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane				
Common	<i>Portulaca oleracea</i>			
Horse	<i>Trinantha portulacastrum</i>			
Radish Wild	<i>Raphanus raphanistrum</i>			
Ragweed Common	<i>Ambrosia artemisiifolia</i>			
Redmaids	<i>Calandrinia ciliata var menziessi</i>			
Redweed	<i>Melochia corchorifolia</i>			
Shepherd s purse	<i>Capsella bursa pastonis</i>			
Smellmelon	<i>Cucumis melo</i>			
Sowthistle Annual <sup>3</sup>	<i>Sonchus oleraceus</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Spurred Anoda	<i>Anoda cristata</i>			
Thistle Russian	<i>Salsola iberica</i>			
Tropic Croton	<i>Croton glandulosus</i>			
Venice Mallow	<i>Hibiscus trionum</i>			
Waterhemp				
Common	<i>Amaranthus rudis</i>			
Tall	<i>Amaranthus tuberculatus</i>			
Wild Poinsettia	<i>Euphorbia heterophylla</i>			
White Cockle	<i>Silene latifolia</i>			
Wormwood Biennial	<i>Artemisia biennis</i>			
Yellow Rocket	<i>Barbarea vulgaris</i>			

continued

<sup>1</sup> *Valor* Herbicide can be used on soils with greater than 10% organic matter however length of residual control may be shorter than on soils with lower organic matter content

<sup>2</sup> A maximum *Valor* Herbicide rate of 6 oz /A per application should be used on any soil that has a sand plus gravel content over 80 / if bushes trees or vines are under 3 years of age

<sup>3</sup> Except CA

**Table 10 Weeds Controlled by Preemergence Application of Valor Herbicide (continued)**

<b>BROADLEAF WEED SPECIES</b>				
<b>COMMON NAME</b>	<b>SCIENTIFIC NAME</b>	<b>ORGANIC MATTER</b>	<b>SOIL TYPE</b>	<b>VALOR HERBICIDE RATE</b>
<b>GRASS WEED SPECIES</b>		Up to 10% <sup>1</sup>	All Soil Types <sup>2</sup>	Asparagus Garlic Hops 6 oz/A
Barnyardgrass	<i>Echinochloa crus galli</i>			Sugarcane 6 to 8 oz /A
Bluegrass Annual	<i>Poa annua</i>			
Crabgrass				
Large	<i>Digitaria sanguinalis</i>			Bushberries Grapes Nut Trees (including Pistachio) Pome Fruit Stone Fruit and Non Bearing Fruit Trees 6 to 12 oz/A <sup>2</sup>
Smooth	<i>Digitaria ischaemum</i>			
Foxtails				
Bristly	<i>Setaria verticillata</i>			
Giant	<i>Setaria faberi</i>			
Green	<i>Setaria viridis</i>			
Yellow	<i>Setaria glauca</i>			
Goosegrass	<i>Eleusine indica</i>			
Guineagrass	<i>Panicum maximum</i>			
Johnsongrass Seedling	<i>Sorghum halepense</i>			
Lovegrass California	<i>Eragrostis diffusa</i>			
Panicum				
Fall	<i>Panicum dichotomiflorum</i>			
Texas	<i>Panicum texaum</i>			
Ryegrass Italian	<i>Lolium multiflorum</i>			
Signalgrass Broadleaf	<i>Brachiaria platyphylla</i>			
				To Maintain Bare Ground on Non Crop Areas of Farms Orchards & Vineyards 6 to 12 oz/A

<sup>1</sup>Valor Herbicide can be used on soils with greater than 10% however length of residual control may be shorter than on soils with lower organic matter content

<sup>2</sup>A maximum Valor Herbicide rate of 6 oz /A per application should be used on any soil that has a sand plus gravel content over 80% if bushes trees or vines are under 3 years of age

## DIRECTIONS FOR USE IN SUGARCANE

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 8 oz of VALOR per acre per application
- Do not make a sequential application within 14 days of the first application
- Do not apply more than 12 oz of VALOR per acre during a single growing season
- Do not apply within 90 days of harvest

### TIMING TO SUGARCANE

VALOR may be applied from 2 weeks prior to planting to before the sugarcane emerges post directed or at layby. Select the proper VALOR rate from Table 10 according to anticipated weed spectrum and soil organic matter content for preemergence applications. Select VALOR rate from Table 11 according to emerged weed spectrum and weed heights for post directed and layby applications.

### TIMING TO WEEDS

#### Burndown – Preemergence to Sugarcane Postemergence to Weeds

VALOR may be used for preemergence control and to assist in postemergence burndown of many annual broadleaf weeds in sugarcane. For control of emerged weeds, choose the most appropriate tank mix partner from Table 12. Apply VALOR **before the crop emerges**. To ensure thorough coverage, use a minimum of 15 gals of spray solution per acre. All VALOR tank mixes applied to assist in the control of emerged weeds must be applied with crop oil concentrate or methylated seed oil at 1 qt /A or a non ionic surfactant at 0.25% v/v. Some tank mix products such as ROUNDUP Original Max (glyphosate) may be formulated with a suitable adjuvant and do not require additional adjuvant.

#### Preemergence – Preemergence to Sugarcane Preemergence to Weeds

VALOR may be used for preemergence control of many annual broadleaf and grassy weeds in sugarcane. Select rate based on anticipated weed spectrum and soil organic matter content from Table 10. Apply VALOR **before the crop emerges**.

#### Post Directed – Postemergence to Sugarcane Postemergence to Weeds

Post directed applications should only be made to upright sugarcane varieties after the sugarcane has exceeded 24 inches in height and has begun to joint. Post directed applications should not be made to PINEAPPLE varieties. Post directed applications to PINEAPPLE varieties or to upright varieties that have not exceeded 24 inches in height and have not begun to joint may result in unacceptable crop injury. To ensure thorough coverage, use a minimum of 15 gals of spray solution per acre. Post directed applications of VALOR must include a crop oil concentrate or methylated seed oil at 1 qt /A or a non ionic surfactant at 0.25% v/v. Select the proper VALOR rate based on weed spectrum and weed height from Table 11.

#### Layby – Postemergence to Sugarcane Postemergence to Weeds

Layby applications can be made to upright and PINEAPPLE varieties after the sugarcane has exceeded 30 inches in height and the spray solution will not contact foliage above 6 inches from the base of the sugarcane. To ensure thorough coverage, use a minimum of 15 gals of spray solution per acre. Layby applications of VALOR must be applied with crop oil concentrate or methylated seed oil at 1 qt /A or a non ionic surfactant at 0.25% v/v. Select the proper VALOR rate based on weed spectrum and weed height from Table 11.

Table 11 Broadleaf Weeds Controlled by Post Directed or Layby Application of VALOR in Sugarcane

BROADLEAF WEED SPECIES		WEED HEIGHT (inches)	
COMMON NAME	SCIENTIFIC NAME	3 oz /A	4 oz /A
Bindweed Field <sup>1</sup>	<i>Convolvulus arvensis</i>	4	8
Carpetweed	<i>Mollugo verticillata</i>	4	4
Cocklebur Common	<i>Xanthium strumarium</i>	4	4
Florida Beggarweed	<i>Desmodium tortuosum</i>	2	2
Hemp Sesbania	<i>Sesbania exaltata</i>	6	8
Jimsonweed	<i>Datura stramonium</i>	4	4
Lambsquarters Common	<i>Chenopodium album</i>	4	4
Morningglories			
Entireleaf	<i>Ipomoea hederacea</i> var <i>integruscula</i>		4
Ivyleaf	<i>Ipomoea hederacea</i>	4	4
Pitted	<i>Ipomoea lacunosa</i>	4	6
Red	<i>Ipomoea coccinea</i>		4
Tall	<i>Ipomoea purpurea</i>	2	4
Mustard Wild	<i>Brassica kaber</i>	6	6
Pigweeds			
Palmer Amaranth	<i>Amaranthus palmeri</i>	4	6
Redroot	<i>Amaranthus retroflexus</i>	4	6
Smooth	<i>Amaranthus hybridus</i>	4	6
Plantain Broadleaf	<i>Plantago major</i>	6	6
Prickly Sida	<i>Sida spinosa</i>	4	6
Purslanes			
Common	<i>Portulaca oleracea</i>	2	4
Rock	<i>Calandrinia</i> spp		2
Ragweeds			
Common	<i>Ambrosia artemisiifolia</i>	2	2
Giant	<i>Ambrosia trifida</i>	4	4
Rice Flatsedge	<i>Cyperus iria</i>	2	4
Sicklepod	<i>Senna obtusifolia</i>	4	4
Smartweeds			
Ladysthumb	<i>Polygonum persicaria</i>	4	4
Pale	<i>Polygonum lapathifolium</i>	4	4
Pennsylvania	<i>Polygonum pennsylvanicum</i>	4	4
Spotted Spurge	<i>Euphorbia maculata</i>	4	4
Velvetleaf	<i>Abutilon theophrasti</i>	4	6
Venice Mallow	<i>Hibiscus trionum</i>	2	2
Waterhemp			
Common	<i>Amaranthus rudis</i>	2	2
Tall	<i>Amaranthus tuberculatus</i>	2	2

<sup>1</sup> VALOR tank mixes will only control the above ground portion of field bindweed. Repeated applications will be needed to control regrowth

## TANK MIXES

VALOR may be tank mixed with the herbicides listed in Table 12 for additional weed control in burndown preemergence post directed and layby applications Refer to tank mix partner s label for adjuvant recommendations

**Table 12 Tank Mixes with VALOR for Post Directed or Layby Use in Sugarcane**

TANK MIX PARTNER <sup>1</sup>	TARGET WEEDS	BURNDOWN	POST DIRECTED <sup>2</sup>	LAYBY
2 4 D amine	Annual and Perennial Broadleaf Weeds	X		
atrazine	Pigweeds Cocklebur	X	X	X
Asulox <sup>®3</sup>	Annual Grasses		X	X
Evik <sup>®4</sup>	Annual Grasses		X	X
glyphosate <sup>5</sup>	Annual and Perennial Weeds	X		X
metribuzin <sup>6</sup>	Broadleaf Panicum Goosegrass		X	X
Sempre <sup>®</sup>	Purple Nutsedge Yellow Nutsedge	X	X	X
Weedmaster <sup>®</sup>	Annual and Perennial Broadleaf Weeds	X		

<sup>1</sup> Refer to tank mix product labels for specific recommendations for control of emerged weeds present not listed in Table 11

<sup>2</sup> Post directed applications should only be made to upright sugarcane varieties after the sugarcane has exceeded 24 inches in height Post directed applications should not be made to PINEAPPLE varieties Post directed applications to PINEAPPLE varieties or to upright varieties that have not exceeded 24 inches in height may result in unacceptable crop injury

<sup>3</sup> Apply to sugarcane at least 24 inches tall

<sup>4</sup> Apply before weeds are greater than 6 inches tall

<sup>5</sup> Glyphosate applications must be made with a hooded sprayer Sugarcane must be at least 3 ft tall Contact with the sugarcane foliage by either the spray mixture or the treated weed foliage will result in sugarcane injury

<sup>6</sup> Refer to metribuzin label for restrictions based on soil type

## ADDITIONAL PREEMERGENCE BROADLEAF CONTROL

VALOR can be tank mixed with atrazine or diuron for additional preemergence broadleaf control

## ADDITIONAL PREEMERGENCE GRASS CONTROL

VALOR can be tank mixed with PROWL (or other pendimethalin products) for additional preemergence grass control provided sugarcane has not emerged

## DIRECTIONS FOR USE IN SUNFLOWER AND SAFFLOWER

### **HARVEST AID**

#### **RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season
- Do not harvest within 5 days of application

Desiccation from *Valor* Herbicide requires the addition of an agronomically approved adjuvant to the spray mixture. A methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A should be used. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* Herbicide with glyphosate or paraquat will increase control of emerged weeds and aid in harvest for sunflowers. Tank mixing *Valor* Herbicide with glyphosate will increase control of emerged weeds and aid in harvest for safflower.

#### **TIMING TO SUNFLOWER AND SAFFLOWER**

Apply *Valor* Herbicide at 1.5 to 2 oz/A when crop is mature (when seed is 35% moisture or less). For many varieties, this is when the backs of the heads are turning yellow and the bracts are turning brown. Sunflower and safflower can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure recommendations for postemergence application.

## DIRECTIONS FOR USE IN SWEET POTATO [For Use in the States of Arizona, California and Hawaii Only]

#### **RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 3 oz of *VALOR* per acre during a single growing season
- Do not apply postemergence to sweet potatoes
- Do not use greenhouse grown transplants
- Do not use transplants harvested more than 2 days prior to transplanting
- Do not use on any sweet potato variety other than *BEAUREGARD* unless user has tested *VALOR* on other variety and has found crop tolerance to be acceptable
- Do not apply as a part of any tank mix except with labeled rates of *COMMAND* if tank mix is applied prior to transplanting

#### **TIMING TO SWEET POTATOES**

*VALOR* must be applied prior to transplanting sweet potatoes.

#### **TIMING TO WEEDS**

##### **Preemergence To Weeds**

Apply *VALOR* to soil prior to transplanting sweet potato slips for the preemergence control of the weeds listed in Table 1.

## DIRECTIONS FOR USE IN WHEAT

[For use in the states of  
DE ID KY MD MN MT NC ND NJ OR SC SD TN VA and WA Only]

### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 2 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 2 oz of *Valor* Herbicide per acre during a single growing season

### PRE PLANT APPLICATIONS, PRE EMERGENCE WEED CONTROL

#### RESTRICTIONS AND LIMITATIONS

- For pre plant weed control use only on no till or minimum tillage fields where the previous year's crop residue has not been incorporated into the soil
- [Plant wheat no sooner than 7 days after *Valor* Herbicide application in the states of DE ID KY MD MN MT NC ND NJ OR SC SD TN VA or WA]
- [Plant wheat no sooner than 14 days after *Valor* Herbicide application in the states of DE ID KY MD MN MT NC ND NJ OR SC SD TN VA or WA]
- [Do not use on Durum wheat]
- Do not irrigate between emergence and spike
- Wheat must be planted a minimum of 1" deep
- Do not graze until wheat has reached 5 inches in height

#### Burndown Use Directions

*Valor* Herbicide applied as part of a burndown program at 2 oz/A may be used for residual weed control as well as to assist in postemergence burndown of many weeds where wheat will be planted directly into the residue of the previous crop. See Directions for Use in Fall Burndown Programs in Fields to be Planted to Barley Field Pea Flax Lentil Safflower Sunflower and Wheat for rates and timing of applications. For control of emerged weeds *Valor* Herbicide must be applied with an appropriate burndown tank mix partner. To ensure thorough coverage use a minimum of 15 gallons of spray solution per acre. Refer to tank mix partner's label for recommended application pressure and recommended adjuvant systems.

### POST PLANT, PRE EMERGENCE WEED CONTROL

#### RESTRICTIONS AND LIMITATIONS

- For post plant pre emergence weed control use only on no till or minimum tillage fields where the previous crop residue has not been incorporated into the soil
- Apply *Valor* Herbicide up to 2 days after planting
- [Do not use on Durum wheat]
- Do not irrigate between emergence and spike
- Wheat must be planted a minimum of 1" deep
- Do not graze until wheat has reached 5 inches in height

#### Use Directions

*Valor* Herbicide applied at 2 oz/A may be used for residual weed control where wheat has been planted directly into the residue of the previous year. Application must be made no later than 2 days after planting.]

### HARVEST AID

#### RESTRICTIONS AND LIMITATIONS

- Do not harvest within 10 days of application

#### Use Directions

*Valor* Herbicide applied at 2 oz/A for desiccation requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* Herbicide with glyphosate will increase control of emerged weeds and aid in harvest.

To ensure thorough coverage use a minimum of 10 gallons spray solution per acre by ground application and a minimum of 5 gallons per acre by aerial application. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for postemergence application.

#### TIMING TO WHEAT

Apply *Valor* at 1.5 to 2 oz/A after wheat reaches the hard dough stage and grain has no more than 30% moisture. Wheat can be harvested 10 days after application. Valent recommends tank mixing with glyphosate.

## DIRECTIONS FOR USE IN BUSHBERRIES, GRAPE, NUT TREES (INCLUDING PISTACHIO), POME FRUIT, STONE FRUIT AND NON BEARING FRUIT TREES

### RESTRICTIONS AND LIMITATIONS

Do not apply more than 12 oz of *Valor* Herbicide per acre during a single application  
 Do not apply more than 24 oz of *Valor* Herbicide per acre during a 12 month period except Bushberries  
 for Bushberries do not apply more than 12 oz of *Valor* Herbicide per acre during a 12 month period  
 Do not make a sequential application within 30 days of the first application except nut trees do not make  
 a sequential application within 60 days of the first application  
 A maximum *Valor* Herbicide rate of 6 oz/A per application should be used on any soil that has a sand plus  
 gravel content over 80% if bushes trees or vines are less than 3 years of age (Two applications of 6 oz/A in  
 a 12 month period can still be made as long as there have been 60 days between applications)  
 Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other  
 desirable vegetation  
 Do not apply within 300 yards of non dormant pears  
 Raise mower height during all mowing to reduce dust Dust created by mowing can drift onto desirable  
 vegetation resulting in injury  
 Do not apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be  
 applied immediately after application  
 [Do not mow treated areas between bud break and final harvest Dust created by mowing may drift onto  
 desirable vegetation resulting in injury ]  
 Follow the most restrictive label limitations and precautions of the tank mix product(s) being used  
 Avoid direct or indirect spray contact to foliage and green bark (non barked trunk and non barked vines  
 with the exception of undesirable suckers)

For bushberries grape nut trees (including pistachio) and non bearing fruit trees *Valor* Herbicide should be  
 applied as a uniform broadcast application to the orchard or vineyard floor or as a uniform band directed at the  
 base of the bush trunk or vine For pome fruit and stone fruit *Valor* Herbicide can only be applied as a uniform  
 band directed at the base of the trunk prior to pink bud in apple and bud break in stone fruit and pear The  
 preferred application timing for *Valor* Herbicide is in the fall to maximize the potential for rainfall to activate and set  
 the herbicide Do not apply over the top of crop or allow spray to come in contact with crop as a result of  
 application or drift

### Preemergence Application

Apply 6 to 12 oz (0.188 to 0.38 lb ai/A) of *Valor* Herbicide per broadcast acre as a preemergence application  
 Preemergence (to weed emergence) applications of *Valor* Herbicide should be made to a weed free soil surface  
 Preemergence applications of *Valor* Herbicide must be completed prior to weed emergence Moisture is  
 necessary to activate *Valor* Herbicide on soil for residual weed control Dry weather following application of *Valor*  
 Herbicide may reduce effectiveness However when adequate moisture is received after dry conditions *Valor*  
 Herbicide will control susceptible germinating weeds

### Postemergence Application

Apply 6 to 12 oz (0.188 to 0.38 lb ai/A) of *Valor* Herbicide per broadcast acre plus an adjuvant (0.25% v/v  
 non ionic surfactant or 1 qt/A crop oil concentrate) The addition of an adjuvant enhances *Valor* Herbicide  
 activity on emerged weeds Thorough spray coverage is necessary to maximize the postemergence activity of  
*Valor* Herbicide

Refer to Table 10 for weeds controlled by the residual activity of *Valor* Herbicide *Valor* Herbicide should be tank  
 mixed with a labeled burndown herbicide for control of the emerged weeds listed in Table 13 Refer to tank mix  
 partner's label for additional weed species and increased weed heights claimed Refer to tank mix partner's label  
 for additional restrictions including minimum carrier volume and crops in which tank mix partner may be used  
 Burndown tank mix partners include glyphosate paraquat 2,4-D and RELY® Tank mixes with glyphosate or 2,4-  
 D containing products are not recommended during the period after bloom through final harvest to ensure crop  
 safety from drift

Residual weed control will be reduced if vegetation prevents the *Valor* Herbicide from reaching the soil  
 surface If vegetation is heavy it is recommended to use a burndown herbicide with *Valor* Herbicide and  
 make a sequential *Valor* Herbicide application prior to the emergence of new weeds



### Carrier Volume and Spray Pressure

To ensure thorough coverage in burndown applications use a minimum of 15 gallons of spray solution per acre  
Use higher gallonage if dense vegetation or heavy crop residue is present

Nozzle selection should meet manufacturer's gallonage and pressure recommendations

### Banded Application

Rates listed in Table 13 refer to a broadcast application covering the entire acre. When making a banded application the rate must be reduced according to the following formula

Amount Needed per Acre for Banded Application	=	$\frac{\text{Band Width in inches}}{\text{Row Width in inches}}$	X	Rate per Broadcast Acre
--------------------------------------------------	---	------------------------------------------------------------------	---	----------------------------

### USE PRECAUTIONS FOR BUSHBERRIES

Bushberries Aronia Berry Black Currant Blueberry (Highbush Rabbit eye and Lowbush) Buffalo Currant  
Chilean Guava Cranberry (Highbush) Elderberry European Barberry Gooseberry Honeysuckle (edible)  
Huckleberry Jostaberry Juneberry Lingonberry Native Currant Red Currant Salal and Sea Buckthorn

Do not use in the states of Idaho Oregon or Washington except west of the Cascade Mountains in the following counties

**Oregon** Benton Clackamas Clatsop Columbia Coos Curry Douglas Jackson Josephine Lane  
Lincoln Linn Marion Multnomah Polk Tillamook Umatilla Yamhill and Washington

**Washington** Benton Clallam Clark Cowlitz Franklin Grant Grays Harbor King Jefferson Kitsap  
Lewis Pacific Pierce Skagit Snohomish Thurston Wahkiakum Walla Walla and Whatcom

Do not apply to Bushberries established less than 2 years unless they are protected from spray contact by non porous wrap grow tubes or waxed containers

Do not apply within 7 days of harvest

### USE PRECAUTIONS FOR GRAPES

Do not apply within 60 days of harvest

Do not apply to grapes established less than 2 years unless they are trellised at least 3 ft from the soil surface or are protected from spray contact by non porous wrap grow tubes or waxed containers

Do not apply to grapes that are not trellised or staked unless they are free standing

Avoid direct or indirect spray contact to foliage and green bark (non barked vines with the exception of undesirable suckers)

New plantings of own rooted varieties such as Concord should be planted so that all roots are a minimum 8 inches below the soil surface to be treated. In some situations this may require hilling soil around newly planted vines so that the settled depth of the hill will be 4 to 5 inches above the vineyard floor

### Juice Raisin and Wine Grapes

Do not apply during the period after bud break through final harvest unless using shielded application equipment and applicator can ensure spray drift will not come in contact with crop fruit or foliage. Shielded applications during this time period should not be made with glyphosate or products containing glyphosate

### Table Grapes

VALOR may be applied during the period following final harvest up to bud break

Do not apply after bud break

## USE PRECAUTIONS FOR NUT TREES (INCLUDING PISTACHIO) POME FRUIT AND STONE FRUIT

Nut Trees Almond Beechnut Betelnut Black Walnut Brazil Nut Butternut Cashew Chestnut Chinquapin Coconut English Walnut Filbert (Hazelnut) Ginkgo Heartnut Hickory Nut Macadamia Nut Oak Pecan Pili Nut Pine Nut Pistachio and Tropical Almond Pome Fruit Apple Crabapple Loquat Mayhaw Pear Pear (Oriental) and Quince

Stone Fruit Apricot Cherries (Sweet and Tart) Nectarine Peach Plum (Chickasaw Damson Japanese) Plumcot and Prune

- **California only** For almonds and stone fruit in the counties of Merced San Joaquin and Stanislaus follow supplemental labeling provided by Valent U S A Corporation
- For pome fruit and stone fruit *Valor* Herbicide can only be applied as a uniform band directed at the base of the trunk prior to silver tip in apples and bud break in stone fruit
- Do not apply to pears in the states of Oregon or Washington
- For pome fruit and stone fruit do not apply to row middles (area between berms)
- For nut trees (including Pistachio) apply after bud break through final harvest using shielded application equipment if the applicator can ensure the spray drift will not come into contact with non target vegetation crop fruit and/or foliage Shielded application equipment is not required if the following application parameters are followed
  - Application pressure (at boom) < 30 PSI
  - Application speed < 5 MPH
  - Applicator can ensure the spray drift will not come into contact with non target vegetation crop fruit and/or foliage

Do not apply within 60 days prior to harvest

Do not apply to trees established less than one year unless protected from spray contact by non porous wraps grow tubes paint or waxed containers

Do not use in the states of Oregon or Washington except in the following counties unless the additional restrictions listed below are followed

**Oregon** Benton Clackamas Clatsop Columbia Coos Curry Douglas Jackson Josephine Lane Lincoln Linn Marion Morrow Multnomah Polk Tillamook Umatilla Yamhill and Washington

**Washington** Clallam Cowlitz Grays Harbor King Jefferson Kitsap Lewis Pacific Pierce Skagit Snohomish Thurston Wahkiakum and Whatcom

- For apples east of the Cascade Mountains in Washington (counties not listed above) follow the restrictions above plus
  - o Apply between final harvest and January 1
  - o Apply only to apple blocks with an established (2 years or older) permanent cover crop that covers a minimum of 60% of the surface area in the block
  - o Application must be incorporated with a minimum of one half inch of water within 48 hours after application
  - o Do not apply to powdery soils or soils susceptible to wind displacement
  - o Apply only to orchard berms
  - o Do not mow the treated berm areas of the orchard

## USE PRECAUTIONS FOR NON BEARING FRUIT TREES

Non Bearing Avocado Fig Grapefruit Lemon Olive Orange Pomegranate and Tangerine

Do not apply more than 12 oz of *Valor* Herbicide per acre during a single application

Do not apply more than 24 oz of *Valor* Herbicide per acre during a 12 month period

Do not harvest fruit from treated trees within one year of application

Do not apply to trees established less than one year unless protected from spray contact by non porous wraps grow tubes or waxed containers

Do not apply during the period after flowering through leaf drop unless using shielded application equipment and the applicator can ensure spray drift will not come in contact with the crop foliage

Table 13 Weeds Controlled by Postemergence Activity of Valor Herbicide Tank mixes

BROADLEAF WEED SPECIES			
COMMON NAME	SCIENTIFIC NAME	WEED HEIGHT/LENGTH (inches)	VALOR HERBICIDE RATE
Bindweed Field <sup>1</sup>	<i>Convolvulus arvensis</i>	8	6 to 12 oz /A
Carpetweed	<i>Mollugo verticillata</i>	4	
Chickweeds			
Common	<i>Stellaria media</i>	4	
Mouseear	<i>Cerastium vulgatum</i>	4	
Cocklebur Common	<i>Xanthium strumarium</i>	4	
Eveningprimrose Cutleaf <sup>2</sup>	<i>Oenothera laciniata</i>	12	
Filaree			
Broadleaf	<i>Erodium botrys</i>	4	
Redstem	<i>Erodium cicutarium</i>	4	
Florida Beggarweed	<i>Desmodium tortuosum</i>	2	
Hemp Sesbania	<i>Sesbania exaltata</i>	8	
Jimsonweed	<i>Datura stramonium</i>	4	
Lambsquarters Common	<i>Chenopodium album</i>	4	
Morningglories			
Entireleaf	<i>Ipomoea hederacea</i> var <i>integruscula</i>	4	
Ivyleaf	<i>Ipomoea hederacea</i>	4	
Pitted	<i>Ipomoea lacunosa</i>	6	
Red/Scarlet	<i>Ipomoea coccinea</i>	4	
Tall	<i>Ipomoea purpurea</i>	4	
Mustard Wild	<i>Brassica kaber</i>	6	
Pigweeds			
Palmer Amaranth	<i>Amaranthus palmeri</i>	6	
Redroot	<i>Amaranthus retroflexus</i>	6	
Smooth	<i>Amaranthus hybridus</i>	6	
Plantain Broadleaf	<i>Plantago major</i>	6	
Prickly Sida (Teaweed)	<i>Sida spinosa</i>	6	
Purslanes			
Common	<i>Portulaca oleracea</i>	4	
Rock	<i>Calandrinia</i> spp	2	
Ragweeds			
Common	<i>Ambrosia artemisiifolia</i>	2	
Giant	<i>Ambrosia trifida</i>	4	
Rice Flatsedge	<i>Cyperus iria</i>	4	
Sicklepod	<i>Senna obtusifolia</i>	4	
Smartweeds			
Ladysthumb	<i>Polygonum persicaria</i>	4	
Pale	<i>Polygonum lapathifolium</i>	4	
Pennsylvania	<i>Polygonum pensylvanicum</i>	4	
Spotted Spurge	<i>Euphorbia maculata</i>	4	
Velvetleaf	<i>Abutilon theophrasti</i>	4	
Venice Mallow	<i>Hibiscus trionum</i>	4	
Waterhemp			
Common	<i>Amaranthus rudis</i>	2	
Tall	<i>Amaranthus tuberculatus</i>	2	

<sup>1</sup> Valor Herbicide will only provide control of the above ground portion of bindweed. Repeated applications will be needed to control regrowth.

<sup>2</sup> For acceptable control, cutleaf eveningprimrose should be 12 inches or less and in the rosette stage. Crop oil concentrate at 1 pt /A or non ionic surfactant at 0.25 / v/v should be added to glyphosate tank mixes for cutleaf eveningprimrose control, including glyphosate formulations that contain a built in adjuvant system.

**ADDITIONAL RESIDUAL WEED CONTROL**

*Valor* Herbicide may be tank mixed with oryzalin (SURFLAN®), simazine or diuron for additional residual weed control. Always read and follow label use directions for all products being used.

## **DIRECTIONS FOR USE TO MAINTAIN BARE GROUND ON NON CROP AREAS OF FARMS, ORCHARDS AND VINEYARDS**

**RESTRICTIONS AND LIMITATIONS**

- Do not apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation
- Do not apply to ditch banks

*VALOR* when used as directed can be used on farms, orchards and vineyards for non selective vegetation control to maintain bare ground on non crop areas that must be kept weed free. Follow all applicable directions as outlined above under **USE INFORMATION**.

*VALOR* offers residual and postemergence control of susceptible broadleaf and grass weeds as well as an additional mode of action to assist in the control of ALS (acetolactate synthase) resistant weeds. *VALOR* can be tank mixed with the herbicides listed in Table 14 for increased residual or postemergence control. The length of residual control is dependent on the rate applied as well as on rainfall and temperature conditions. Length of residual control will decrease as temperature and precipitation increase. *VALOR* rates of 6 to 12 oz /A are required to provide residual control of the weeds listed in Table 10.

**PREEMERGENCE APPLICATION**

Apply 6 to 12 oz (0.188 to 0.38 lb ai/A) of *VALOR* per broadcast acre as a preemergence application. Preemergence (to weed emergence) applications of *VALOR* should be made to a weed free soil surface. Preemergence applications of *VALOR* must be completed prior to weed emergence. Moisture is necessary to activate *VALOR* on soil for residual weed control. Dry weather following application of *VALOR* may reduce effectiveness. However, when adequate moisture is received after dry conditions, *VALOR* will control susceptible germinating weeds.

**POSTEMERGENCE APPLICATION**

Apply 6 to 12 oz (0.188 to 0.38 lb ai/A) of *VALOR* per broadcast acre plus an adjuvant (0.25% v/v non ionic surfactant or 1 qt /A crop oil concentrate). The addition of an adjuvant enhances *VALOR* activity on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of *VALOR*. Emerged weeds are controlled postemergence with *VALOR*; however, translocation of *VALOR* within a weed is limited and control is affected by spray coverage and by the addition of an adjuvant. The most effective postemergence weed control with *VALOR* occurs when applied in combination with a surfactant to weeds less than 2 inches in height. A tank mix partner should be used in combination with *VALOR* for the postemergence control of weeds larger than 2 inches. Recommended tank mix partners are listed in Table 14.

**IMPORTANT** Completely read and follow the label of any potential tank mix partner with *VALOR*. When using tank mixtures, use conditions must be in accordance with the most restrictive of the label limitations and precautions on either herbicide label.

**Table 14 Tank Mix Combinations to Maintain Bare Ground on Non Crop Areas**

glyphosate	2,4-D	Rely	paraquat
------------	-------	------	----------

## STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage disposal or cleaning of equipment

### PESTICIDE STORAGE

Keep pesticide in original container

Store in a cool dry secure place

Do not put formulation or dilute spray solution into food or drink containers

Do not contaminate food or foodstuffs

Do not store or transport near feed or food

Not for use or storage in or around the home

For help with any spill leak fire or exposure involving this material call day or night (800) 892 0099

### PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

### CONTAINER HANDLING

[Nonrefillable container Do not reuse or refill this container Offer for recycling if available Clean container promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill ]

or

***Note to EPA reviewer if this product is shipped in containers greater than 50 lbs the following container handling statement will be added to the label***

[Container statement for nonrefillable container with liner]

**[Nonrefillable bag]** Do not reuse or refill this bag Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles Empty residue into equipment Do not reuse bag Dispose of bag in a sanitary landfill or by incineration if allowed by State and local authorities Offer for recycling if available **Liner** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles Empty residue into equipment Do not reuse liner Dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities ]

or

[Container statement for nonrefillable drum with liner]

**[Nonrefillable container]** Do not reuse or refill this container Offer for recycling if available Clean container promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times **Liner** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles Empty residue into equipment Do not reuse liner Dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities ]

Copyright© 2012 by Valent U S A Corporation

*Cobra Phoenix Select Max* and *Valor* are trademark and registered trademarks of Valent U S A Corporation

*Valent Tank Cleaner* is a product of Valent U S A Corporation

*Basis Classic Express Lorox* and *Resolve* are trademarks and registered trademarks of E I du Pont de Nemours and Company

*Command* is a registered trademark of FMC Corporation

*Asulox* is a registered trademark of UPI USA Corp Protection

*Rely* is a registered trademark of Bayer

*Banvel Pursuit Scepter Squadron Steel* and *Weedmaster* are registered trademarks of BASF

*Boundary Dual Magnum Dual II Magnum* and *Evik* are registered trademarks of Syngenta

*FirstRate Hornet Python Sonalan* and *Surpass* are registered trademarks of Dow AgroSciences LLC

*Frontier Outlook Prowl* and *Raptor* are registered trademarks of BASF Ag

*Harness IntRRo Lasso Micro Tech Roundup Roundup Ready Roundup Original Roundup Power Max* and *Sempre* are registered trademarks of Monsanto Co

*Surflan* is a registered trademark of United Phosphorus Inc

Manufactured for

**Valent U S A Corporation**

P O Box 8025

Walnut Creek CA 94596 8025

www.valent.com

Made in U S A

EPA Reg No 59639 99

EPA Est

059639 00099 20120914 VLR FTAMEND

20120830/20120118

# Supplemental Label



Valent U.S.A. Corporation  
P.O. Box 8025  
Walton, CA 94596-8025

## VALOR<sup>®</sup> Herbicide EPA Reg No 59639 99

### PREEMERGENCE APPLICATION IN PEANUT (NORTH CAROLINA OKLAHOMA AND VIRGINIA ONLY)

This supplemental label expires on June 30 2014 and must not be used or distributed after this date

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

##### ENVIRONMENTAL HAZARDS

This product is toxic to non target plants and aquatic invertebrates. Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff may be hazardous to non target plants and aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

This pesticide is toxic to plants and should be used strictly in accordance with the drift and run off precautions on this label in order to minimize off site exposures.

Under some conditions this product may have a potential to run off to surface water or adjacent land. Where possible use methods which reduce soil erosion such as no till limited till and contour plowing. These methods also reduce pesticide run off. Use of vegetation filter strips along rivers creeks streams wetlands or on the downhill side of fields where run off could occur will minimize water run off and is recommended.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

#### PREEMERGENCE APPLICATION IN PEANUTS IN THE STATES OF NORTH CAROLINA OKLAHOMA AND VIRGINIA

VALOR at 3 oz per acre can be applied within 2 days of planting to control common ragweed tropic croton and entireleaf ivyleaf and tall/scarlet morningglories.

Cool temperatures near emergence (2 consecutive nighttime lows in the 50 s F) in combination with heavy rainfall may result in severe crop injury. VALOR at 3 oz /A should only be used in these states when other alternatives are not available for adequate control of the weeds listed above and the user acknowledges the risks associated with this use rate under the adverse environmental conditions listed above.

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICATION DIRECTIONS RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL.**

**PLEASE CONTACT VALENT U S A CORPORATION AT 1 800 6 VALENT (682 5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE**

**Copyright © 2004 by Valent U S A Corporation**

Manufactured for  
Valent U S A Corporation  
P O Box 8025  
Walnut Creek CA 94596 8025  
[www.valent.com](http://www.valent.com)

Made in U S A

Form 2004 VLR 0017



65/84

# Supplemental Label



## VALOR<sup>®</sup> Herbicide

EPA Reg No 59639 99

(For Use in Arizona, California and Hawaii Only)

### VALOR<sup>®</sup> HERBICIDE FALLOWBED USE ON TRANSPLANTED MELON, PEPPER AND TOMATO BEDS

This supplemental label expires on June 30 2014 and must not be used or distributed after this date

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION READ THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE BEFORE APPLYING USE OF VALOR ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE**

#### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 4 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 4 oz of *Valor* Herbicide per acre during a single growing season

Many weather related factors including high wind or heavy rains or cool conditions at or near crop transplanting may result in crop injury in fields treated with *Valor* Herbicide On occasion this has resulted in a delay in maturity User should assume these risks before using *Valor* Herbicide

#### TIMING TO CROP

#### VALOR HERBICIDE FALLOWBED USE PRIOR TO TRANSPLANTING

VALOR HERBICIDE RATES	ADJUVANT	GPA	TRANSPLANTING INTERVAL
4 oz/A	Required by burndown tank mix partner	Ground – 20 to 40	2 Months
<b>Application Method</b> Apply with a burndown herbicide labeled for the control of emerged weeds <i>Valor</i> Herbicide when used alone will not provide satisfactory control of emerged weeds			

## USE RESTRICTIONS FOR VALOR HERBICIDE FOR PREEMERGENCE FALLOWBED WEED CONTROL PRIOR TO TRANSPLANTING

- 1 Always read and follow all label directions when using any pesticide alone or in tank mix combinations
- 2 The top 4 inches of the bed from a horizontal and vertical perspective where the crop will be transplanted must be removed prior to transplanting
- 3 Use only healthy transplants Do not use on direct seeded crops



Beds are formed and *Valor* Herbicide is applied with a burndown herbicide



A minimum of 2 months after *Valor* Herbicide application the tops of the beds are removed and the soil from the tops of the beds is placed in the area between the beds



Crops are transplanted into beds

- 4 [On flat beds (tomato only) the soil must be incorporated to a depth of at least 4 inches twice prior to transplanting Failure to incorporate may result in stand reduction and/or crop injury ]
- 5 This use pattern makes no claim for in season weed control after the beds have been disturbed
- 6 Do not apply when weather conditions favor spray drift

**PLEASE CONTACT VALENT U S A CORPORATION AT 800 6 VALENT (682 5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE**

Copyright © 2010 by Valent U S A Corporation

*Valor* is a registered trademark of Valent U S A Corporation

Manufactured for  
Valent U S A Corporation  
P O Box 8025  
Walnut Creek CA 94596 8025  
www.valent.com

Made in U S A

# Supplemental Label



## VALOR® Herbicide

EPA Reg No 59639 99

This supplemental label expires on June 30 2014 and must not be used or distributed after this date

### VALOR® HERBICIDE USE IN ONION (DRY BULB) WITH CHEMIGATION

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION  
READ THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE BEFORE APPLYING  
USE OF VALOR HERBICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE  
PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR  
VALOR HERBICIDE**

#### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 2 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season  
Do not make sequential application within 14 days of the first application
- Do not apply more than 1 oz of *Valor* Herbicide per season on soils that contain greater than 90% sand plus gravel
- Do not apply as part of a tank mix other than with Prowl® H<sub>2</sub>O Herbicide or unacceptable injury may result Other formulations of pendimethalin should not be tank mixed with *Valor* Herbicide for use in onions
- Do not apply with any type of adjuvant
- Do not apply within 45 days of harvest

Use of *Valor* Herbicide may result in necrotic spotting of onion leaves that come in contact with the spray  
User should assume this potential crop response before using *Valor* Herbicide

#### Microrate Application

Sequential applications of *Valor* Herbicide may be applied to onions (dry bulb) between the 2 leaf and 6 leaf stage at rates of 0.5 to 1 oz/A on a 7 day interval

#### TIMING TO ONIONS (dry bulb)

Apply *Valor* Herbicide to transplanted onions (dry bulb) between the 2 leaf and 6 leaf stage and on direct seed onions (dry bulb) between the 3 leaf and 6 leaf stage

#### TIMING TO WEEDS

##### Preemergence – Emerged Onions (dry bulb) Preemergence To Weeds

Apply *Valor* Herbicide to weed free onions (dry bulb) for preemergence control of the weeds listed in Table A  
Broadleaf Weeds Controlled by Residual Activity of *Valor* Herbicide

**Read tank mix product label for rates and weeds controlled Always read and follow label directions for all tank mix products before using The most restrictive labeling of any tank mix product must be followed *Valor* Herbicide when applied according to label use directions will control the weeds listed in Table A This label makes no claims concerning control of other weed species**

## CHEMIGATION

*Valor* Herbicide may be applied through sprinkler irrigation systems in onions (dry bulb) Follow all label recommendations for these crops regarding rates timing of application special instructions and precautions

Apply this product only through center pivot systems End guns must be turned off due to uneven application Do not apply this product through any other type of irrigation system

Crop injury lack of efficacy or illegal pesticide residues in the crop can result from non uniform distribution of treated water

The system must be properly calibrated (with water only) to ensure that the amount of *Valor* Herbicide applied corresponds to the recommended rate

Apply *Valor* Herbicide in 1/2 to 3/4 inches of water during the first sprinkler set Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water To ensure the lines are flushed and free of remaining herbicide a dye indicator may be injected into the lines to mark the end of the application period Once chemigation has begun the run must be completed to ensure no product is left in the system

If you have any questions about calibration you should contact your State Extension Service Specialist equipment manufacturers or other experts

### Special Precautions for Chemigation

- 1 Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place
- 2 A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise
- 3 The system must be free of leaks and clogged nozzles
- 4 The pesticide must be supplied continuously for the duration of the aqueous application An uneven application may cause injury to the crop or poor weed control
- 5 Agitation must be maintained in the nurse tank
- 6 The sprinkler chemigation system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow
- 7 The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump
- 8 The pesticide injection pipeline must contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down
- 9 The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in the case where there is no water pump when the water pressure decreases to the point where pesticide distribution is adversely affected
- 10 The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected
- 11 Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with the pesticides and capable of being fitted with a system interlock
- 12 Do not apply when wind speed favors drift beyond the area intended for treatment

### Chemigation Systems Connected to Public Water Systems

- 1 Public water system means a system for the provision to the public of piped water for human consumption if such a system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year
- 2 Chemigation systems connected to the public water system must contain a functional reduced pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction As an option to the RPZ the water from the public water system should be

discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

- 3 All chemigation systems connected to the public water system must also follow restrictions listed in the preceding section titled **Special Precautions for Chemigation**

**Table A Broadleaf Weeds Controlled by Residual Activity of VALOR HERBICIDE**

<b>BROADLEAF WEED SPECIES</b>				
<b>Common Name</b>	<b>Scientific Name</b>	<b>Organic Matter</b>	<b>Soil Type</b>	<b>Valor Herbicide Rate</b>
Carpetweed	<i>Mollugo verticillata</i>	Up to 5%	All Soil Types	2 oz/A
Chickweeds				
Common	<i>Stellaria media</i>			
Mouseear	<i>Cerastium vulgatum</i>			
Dandelion	<i>Taraxacum officinale</i>			
Eclipta	<i>Eclipta prostrata</i>			
Eveningprimrose Cutleaf	<i>Oenothera laciniata</i>			
Florida Pusley	<i>Richardia scabra</i>			
Henbit	<i>Lamium amplexicaule</i>			
Lambsquarters Common	<i>Chenopodium album</i>			
Little Mallow	<i>Malva parviflora</i>			
Marestail/Horseweed	<i>Conyza canadensis</i>			
Nightshades				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Hairy	<i>Solanum sarrachoides</i>			
Pigweeds				
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane Common	<i>Portulaca oleracea</i>			
Radish Wild	<i>Raphanus raphanistrum</i>			
Redmaids	<i>Calandrinia ciliata</i> var <i>menziesii</i>			
Shepherd's Purse	<i>Capsella bursa pastoris</i>			
Smallflower Morningglory	<i>Jacquemontia tamnifolia</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Venice Mallow	<i>Hibiscus trionum</i>			

**PLEASE CONTACT VALENT U S A CORPORATION AT 800 6 VALENT (682 5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE**

Copyright © 2010 by Valent U S A Corporation

Valor is a registered trademark of Valent U S A Corporation  
Prowl H<sub>2</sub>O is a registered trademark of BASF

Manufactured for  
Valent U S A Corporation  
P O Box 8025  
Walnut Creek CA 94596 8025  
www.valent.com

Made in U S A

# Supplemental Label



## VALOR<sup>®</sup> Herbicide

EPA Reg No 59639 99

(For Use Only in Arizona, California, Colorado, Delaware, Florida, Hawaii, Idaho Maryland Minnesota Montana, Nebraska, Nevada, New Jersey New Mexico, North Carolina, North Dakota, Oregon, South Dakota, Texas, Utah, Virginia, Washington, Washington DC and Wyoming)

This supplemental label expires on June 30 2014 and must not be used or distributed after this date

### VALOR<sup>®</sup> HERBICIDE USE IN POTATO WITH CHEMIGATION

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION  
READ THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE BEFORE APPLYING  
USE OF VALOR HERBICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE  
PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR  
VALOR HERBICIDE**

#### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 1.5 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 1.5 oz of *Valor* Herbicide per acre during a single growing season
- Do not apply to Rill (furrow) irrigated potatoes

Many weather related factors including high wind splashing or heavy rains or cool conditions at or near potato emergence may result in potato injury in fields treated with *Valor* Herbicide. On occasion this has resulted in a delay in maturity. User should assume these risks before using *Valor* Herbicide.

#### TIMING TO POTATOES

*Valor* Herbicide may be applied to potatoes after hilling for the preemergence suppression of the weeds listed in Table A. Weeds Suppressed by Residual Activity of *Valor* Herbicide at 1.5 oz/A. Tank mix *Valor* Herbicide with other labeled herbicides for broad spectrum weed control. A minimum of 2 inches of settled soil must cover the vegetative portion of the potato plant at the time of *Valor* Herbicide application. Application to potatoes with less than 2 inches of soil covering the vegetative portion of the potato may result in crop injury. In areas with historically higher amounts of rainfall during the time of preemergence herbicide applications such as the Red River Valley Minnesota and North Dakota the requirement for 2 inches of settled soil is critical to avoid crop injury. Mechanical incorporation of *Valor* Herbicide will result in decreased weed control and should be avoided. In areas with sprinkler irrigation *Valor* Herbicide should be incorporated with 0.25 to 0.75 inches of irrigation after application and before any sprouts are within 2 inches of the settled soil surface if a rainfall event has not yet occurred.

#### TIMING TO WEEDS

##### Preemergence – Soil Covered Potatoes Preemergence to Weeds

Apply *Valor* Herbicide to soil covered potatoes for the preemergence suppression of the weeds listed in Table A. Harrowing cultivation or corrugating after *Valor* Herbicide application will reduce weed control.

**Read tank mix product label for rates and weeds controlled. Always read and follow label directions for all tank mix products before using. The most restrictive labeling of any tank mix product must be followed.**

## CHEMIGATION

*Valor* Herbicide may be applied through sprinkler irrigation systems in potatoes. Follow all label recommendations for this crop regarding rates, timing of application, special instructions and precautions.

Apply this product only through center pivot systems. End guns must be turned off due to uneven application. Do not apply this product through any other type of irrigation system.

Crop injury, lack of efficacy or illegal pesticide residues in the crop can result from non uniform distribution of treated water.

The system must be properly calibrated (with water only) to ensure that the amount of *Valor* Herbicide applied corresponds to the recommended rate.

Apply *Valor* Herbicide in 1/2 to 3/4 inches of water during the first sprinkler set. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide, a dye indicator may be injected into the lines to mark the end of the application period. Once chemigation has begun, the run must be completed to ensure no product is left in the system.

If you have any questions about calibration, you should contact your State Extension Service Specialist, equipment manufacturers or other experts.

### Special Precautions for Chemigation

- 1 Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.
- 2 A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.
- 3 The system must be free of leaks and clogged nozzles.
- 4 The pesticide must be supplied continuously for the duration of the aqueous application. An uneven application may cause injury to the crop or poor weed control.
- 5 Agitation must be maintained in the nurse tank.
- 6 The sprinkler chemigation system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- 7 The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
- 8 The pesticide injection pipeline must contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 9 The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in the case where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 10 The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 11 Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with the pesticides and capable of being fitted with a system interlock.
- 12 Do not apply when wind speed favors drift beyond the area intended for treatment.

### Chemigation Systems Connected to Public Water Systems

- 1 Public water system means a system for the provision to the public of piped water for human consumption, if such a system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- 2 Chemigation systems connected to the public water system must contain a functional reduced pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

- 3 All chemigation systems connected to the public water system must also follow restrictions listed in the preceding section titled **Special Precautions for Chemigation**

**Table A Weeds Suppressed by Residual Activity of Valor Herbicide at 1.5 oz/A**

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	VALOR HERBICIDE RATE
Lambsquarters Common	<i>Chenopodium album</i>	Up to 5%	1.5 oz/A
Mustard Wild	<i>Brassica kaber</i>		
Nightshades			
Black	<i>Solanum nigrum</i>		
Eastern Black	<i>Solanum ptycanthum</i>		
Hairy	<i>Solanum sarrachoides</i>		
Palmer Amaranth	<i>Amaranthus palmeri</i>		
Pigweeds			
Redroot	<i>Amaranthus retroflexus</i>		
Smooth	<i>Amaranthus hybridus</i>		
Spiny Amaranth	<i>Amaranthus spinosus</i>		
Tumble	<i>Amaranthus albus</i>		
Radish Wild	<i>Raphanus raphanistrum</i>		

**PLEASE CONTACT VALENT U S A CORPORATION AT 800 6 VALENT (682 5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE**

Copyright © 2010 by Valent U S A Corporation

Valor is a registered trademark of Valent U S A Corporation

Manufactured for  
Valent U S A Corporation  
P O Box 8025  
Walnut Creek CA 94596 8025  
www.valent.com

Made in U S A



# Supplemental Label



## VALOR<sup>®</sup> Herbicide

EPA Reg No 59639 99

(For Use Only in Arizona, California, Colorado,  
Hawaii, Idaho, Nebraska, Oregon and Washington)

This supplemental label expires on June 30 2014 and must not be used or distributed after this date

### VALOR<sup>®</sup> HERBICIDE USE IN DRY BEAN FOR WEED SUPPRESSION

Dried cultivars of bean (*Lupinus*) bean (*Phaseolus*) (includes field bean kidney bean lima bean (dry) navy bean pinto bean tepary bean) bean (*Vigna*) (includes adzuki bean blackeyed pea catjang cowpea crowder pea moth bean mung bean rice bean southern pea urd bean) broad bean (dry) chickpea guar lablab bean and lentil

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION READ THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE BEFORE APPLYING USE OF VALOR HERBICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE**

#### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 1.5 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 1.5 oz of *Valor* Herbicide per acre during a single growing season

**Many weather related factors including high wind splashing or heavy rains or cool conditions at or near crop emergence may result in dry bean injury in fields treated with *Valor* Herbicide. On occasion this has resulted in a delay in maturity. User should assume these risks before using *Valor* Herbicide.**

#### TIMING TO DRY BEAN

*Valor* Herbicide may be applied to dry beans within 2 days after planting for the preemergence suppression of the weeds listed in Table A Weeds Suppressed by Residual Activity of *Valor* Herbicide at 1.5 oz/A. Tank mix *Valor* Herbicide with other labeled herbicides for broad spectrum weed control.

#### TIMING TO WEEDS

*Valor* Herbicide may be applied to dry beans prior to planting or preemergence (after planting). Preemergence application of *Valor* Herbicide must be made within 2 days after planting and prior to dry bean emergence. To avoid severe crop injury, do not apply to dry beans after beans begin to crack or have emerged.

Preplant incorporation (PPI) applications may result in reduced weed control.

**ADDITIONAL RESIDUAL GRASS CONTROL**

Valor Herbicide can be tank mixed with pendimethalin for additional grass control

**Table A Weeds Suppressed by Residual Activity of Valor Herbicide at 1.5 oz/A**

COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	VALOR HERBICIDE RATE
Lambsquarters Common	<i>Chenopodium album</i>	Up to 5%	1.5 oz/A
Mustard Wild	<i>Brassica kaber</i>		
Nightshades			
Black	<i>Solanum nigrum</i>		
Eastern Black	<i>Solanum ptycanthum</i>		
Hairy	<i>Solanum sarrachoides</i>		
Palmer Amaranth	<i>Amaranthus palmeri</i>		
Pigweeds			
Redroot	<i>Amaranthus retroflexus</i>		
Smooth	<i>Amaranthus hybridus</i>		
Spiny Amaranth	<i>Amaranthus spinosus</i>		
Tumble	<i>Amaranthus albus</i>		
Prickly Lettuce (China Lettuce)	<i>Lactuca serriola</i>		
Radish Wild	<i>Raphanus raphanistrum</i>		

**PLEASE CONTACT VALENT U S A CORPORATION AT 800 6 VALENT (682 5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE**

Copyright © 2010 by Valent U S A Corporation

Valor is a registered trademark of Valent U S A Corporation

Manufactured for  
Valent U S A Corporation  
P O Box 8025  
Walnut Creek CA 94596 8025  
www.valent.com

Made in U S A

# Supplemental Label



## VALOR<sup>®</sup> Herbicide

EPA Reg No 59639 99

(For Use Only in Arizona, California, Hawaii, Idaho, Oregon and Washington)

### VALOR<sup>®</sup> HERBICIDE USE ON CHICKPEA (GARBANZO BEAN)

This supplemental label expires on June 30 2014 and must not be used or distributed after this date

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION READ THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE BEFORE APPLYING USE OF VALOR HERBICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE**

#### RESTRICTIONS AND LIMITATIONS

- Do not apply more than 2 0 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 2 0 oz of *Valor* Herbicide per acre during a single growing season

**Many weather related factors, including high wind splashing or heavy rains or cool conditions at or near crop emergence, may result in garbanzo bean injury in fields treated with *Valor* Herbicide On occasion this has resulted in a delay in maturity User should assume these risks before using *Valor* Herbicide**

#### TIMING TO CHICKPEA (GARBANZO BEAN)

*Valor* Herbicide may be applied to garbanzo beans within 2 days after planting for the preemergence suppression of the weeds listed in Table A Broadleaf Weeds Controlled by Residual Activity of *Valor* Herbicide Tank mix *Valor* Herbicide with other labeled herbicides for broad spectrum weed control

#### TIMING TO WEEDS

*Valor* Herbicide may be applied to garbanzo beans prior to planting or preemergence (after planting) Preemergence application of *Valor* Herbicide must be made within 2 days after planting and prior to garbanzo bean emergence Application after the garbanzo beans have begun to crack or are emerged will result in severe crop injury Application should not be made when garbanzo beans have begun to crack

Preplant incorporation (PPI) applications may result in reduced weed control

## ADDITIONAL RESIDUAL GRASS CONTROL

Valor Herbicide can be tank mixed with pendimethalin for additional grass control

**Table A Broadleaf Weeds Controlled by Residual Activity of Valor Herbicide**

BROADLEAF WEED SPECIES				
SECTION A				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	VALOR Herbicide RATE
Carpetweed	<i>Mollugo verticillata</i>	Up to 5%	All Soil Types	2 oz/A
Chickweeds				
Common	<i>Stellaria media</i>			
Mouseear	<i>Cerastium vulgatum</i>			
Dandelion	<i>Taraxacum officinale</i>			
Eclipta	<i>Eclipta prostrata</i>			
Eveningprimrose Cutleaf	<i>Oenothera laciniata</i>			
Field Pennycress	<i>Thlaspi arvense</i>			
Florida Pusley	<i>Richardia scabra</i>			
Henbit	<i>Lamium amplexicaule</i>			
Lambsquarters Common	<i>Chenopodium album</i>			
Little Mallow	<i>Malva parviflora</i>			
Marestail/Horseweed	<i>Conyza canadensis</i>			
Mayweed/False Chamomile	<i>Matricaria inodora</i>			
Nightshades				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum elaeagnifolium</i>			
Hairy	<i>Solanum elaeagnifolium</i>			
Pigweeds				
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Lettuce	<i>Lactuca scariola</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane Common	<i>Portulaca oleracea</i>			
Radish Wild	<i>Raphanus raphanistrum</i>			
Redmaids	<i>Calandrinia ciliata</i> var <i>menziesii</i>			
Shepherd's purse	<i>Capsella bursa-pastoris</i>			
Smallflower Morningglory	<i>Jacquemontia tamnifolia</i>			
Sowthistle Prickly	<i>Sonchus asper</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Venice Mallow	<i>Hibiscus trionum</i>			

continued

**Table A Broadleaf Weeds Controlled by Residual Activity of Valor Herbicide(continued)**

<b>SECTION B</b>				
All weeds listed in Section A plus				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	VALOR Herbicide RATE
Coffee Senna	<i>Cassia occidentalis</i>	Up to 3%	All Soil Types	2 oz/A
Common Ragweed	<i>Ambrosia artemisiifolia</i>			
False Chamomile	<i>Tripleurospermum</i>			
Florida Beggarweed	<i>Desmodium tortuosum</i>			
Golden Crownbeard	<i>Verbesina encelioides</i>			
Hairy Indigo	<i>Indigofera hirsuta</i>	3 to 5%	Coarse and Medium Soils (sandy loam loamy sand loamy silt loam silt sandy clay sandy clay loam)	2 oz/A
Hemp Sesbania	<i>Sesbania exaltata</i>			
Jimsonweed	<i>Datura stramonium</i>			
Kochia	<i>Kochia scoparia</i>			
London Rocket	<i>Sisymbrium irio</i>			
Morningglories				
Entireleaf	<i>Ipomoea hederacea</i> var <i>integruscula</i>			
Ivyleaf	<i>Ipomoea hederacea</i>			
Red/Scarlet	<i>Ipomoea coccinea</i>			
Tall	<i>Ipomoea purpurea</i>			
Mustard, Wild	<i>Brassica kaber</i>		Fine Soils (silty clay silty clay loam clay clay loam)	2 oz/A
Palmer Amaranth	<i>Amaranthus palmeri</i>			
Spurred Anoda	<i>Anoda cristata</i>			
Tropic Croton	<i>Croton glandulosus</i>			
Waterhemp				
Common	<i>Amaranthus rudis</i>			
Tall	<i>Amaranthus tuberculatus</i>			
Wild Poinsettia	<i>Euphorbia heterophylla</i>			
Yellow Rocket	<i>Barbarea vulgaris</i>			

**PLEASE CONTACT VALENT U S A CORPORATION AT 800 6 VALENT (682 5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE**

Copyright © 2010 by Valent U S A Corporation

VALOR is a registered trademark of Valent U S A Corporation

Manufactured for  
Valent U S A Corporation  
P O Box 8025  
Walnut Creek CA 94596 8025  
www.valent.com

Made in U S A

# Supplemental Label



## VALOR<sup>®</sup> Herbicide

EPA Reg No 59639 99

This supplemental label expires on June 30 2014 and must not be used or distributed after this date

### VALOR<sup>®</sup> HERBICIDE FOR USE PRECAUTIONS ON ALMOND AND STONE FRUIT IN A DEFINED AREAS OF MERCED, SAN JOAQUIN AND STANISLAUS COUNTIES OF CALIFORNIA


#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION READ THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE BEFORE APPLYING USE OF VALOR HERBICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE**

The use of *Valor* Herbicide in soils common in parts of Merced San Joaquin and Stanislaus counties in California is known to have resulted in injury to almonds under drought stress conditions These soils are characterized by having been cut or filled high sand content low clay content and shallow profiles Growers in the Defined Area must be aware and assume the risk of using *Valor* Herbicide on almond or stone fruit crops The Defined Area can be seen on the Map or by the description that follows

- Intersection of Highway 4 and Escalon Bellota Road at Farmington in San Joaquin County
- Directly South on Escalon Bellota to the Santa Fe Avenue and railroad tracks at Escalon
- Southeast on Santa Fe Avenue down to the Merced River
- East following the Merced River to the Merced/Mariposa County line
- Northwest following the Merced County line through the intersection of Merced and Stanislaus County line following the Stanislaus/Tuolumne County and Calaveras County line to Highway 4
- West on Highway 4 back to the Farmington intersection of Escalon – Bellota Road



**PLEASE CONTACT VALENT U S A CORPORATION AT 800 6 VALENT (682 5368) TO  
DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE**

Copyright © 2010 by Valent U S A Corporation

*Valor* is a registered trademark of Valent U S A Corporation

Manufactured for  
Valent U S A Corporation  
P O Box 8025  
Walnut Creek CA 94596 8025  
[www.valent.com](http://www.valent.com)

Made in U S A

# Supplemental Label

## VALOR<sup>®</sup> HERBICIDE

EPA Reg No 59639 99

### VALOR<sup>®</sup> HERBICIDE FOR USE IN FIELD PEAS, FLAX, LENTILS, SUNFLOWER AND SAFFLOWER AND WHEAT

**VALENT<sup>®</sup>**  
**ACCEPTED**  
 with COMMENTS  
 In EPA Letter Dated  
 SEP 28 2012  
 Under the Federal Insecticide  
 Fungicide and Rodenticide Act  
 as amended for the pesticide  
 registered under EPA Reg No  
 59639-99

This supplemental label expires September 30 2015 and must not be used or distributed after this date

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION  
 READ THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE BEFORE APPLYING USE  
 OF VALOR HERBICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS  
 AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR VALOR HERBICIDE**

#### DIRECTIONS FOR USE IN FIELD PEAS

##### WEED CONTROL

##### **RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 2 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 2 oz of *Valor* Herbicide per acre during a single growing season
- [For use in Idaho Montana Oregon and Washington only ]

**Many weather related factors including high wind splashing or heavy rains or cool conditions at or near crop emergence may result in field pea injury in fields treated with *Valor* Herbicide On occasion this has resulted in a delay in maturity User should assume these risks before using *Valor* Herbicide**

##### **TIMING TO FIELD PEAS**

*Valor* Herbicide may be applied to field peas within 2 days after planting for the preemergence control of the weeds listed in Table 1 Broadleaf Weeds Controlled by Residual Activity of *Valor* Herbicide or Table 8 Weeds Suppressed by Residual Activity of *Valor* Herbicide Tank mix *Valor* Herbicide with other labeled herbicides for broad spectrum weed control

##### **TIMING TO WEEDS**

Apply *Valor* Herbicide to field peas prior to planting or preemergence (after planting) Preemergence application of *Valor* Herbicide must be made within 2 days after planting and prior to field pea emergence To avoid severe crop injury do not apply to field peas after beans begin to crack or have emerged

Preplant incorporation (PPI) applications may result in reduced weed control



**ADDITIONAL RESIDUAL GRASS CONTROL**

*Valor* Herbicide can be tank mixed with pendimethalin for additional grass control

**HARVEST AID****RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season
- Do not harvest within 5 days of application

Desiccation from *Valor* Herbicide requires the addition of an agronomically approved adjuvant to the spray mixture. A methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A should be used. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* Herbicide with glyphosate will increase control of emerged weeds and aid in harvest.

**TIMING TO FIELD PEAS**

Apply *Valor* Herbicide at 1.5 to 2 oz/A when crop is physiologically mature and a minimum of 80% of the pods are yellow to tan in color and 20% are yellow in color. If field peas are treated too early, a reduction in seed quality may occur. Do not spray *Valor* Herbicide on any area of the field with a significant amount of plants with green color. Peas can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure recommendations for postemergence application.

**DIRECTIONS FOR USE IN FLAX****HARVEST AID****RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season
- Do not harvest within 5 days of application

Desiccation from *Valor* Herbicide requires the addition of an agronomically approved adjuvant to the spray mixture. A methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A should be used. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil.

**TIMING TO FLAX**

Apply *Valor* Herbicide at 1.5 to 2 oz/A when crop is physiologically mature and at least 75% of the bolls are brown in color. Flax can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure recommendations for postemergence application.

## DIRECTIONS FOR USE IN LENTILS

### **HARVEST AID**

#### **RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season
- Do not harvest within 5 days of application

Desiccation from *Valor* Herbicide requires the addition of an agronomically approved adjuvant to the spray mixture. A methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A should be used. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* Herbicide with glyphosate or paraquat will increase control of emerged weeds and aid in harvest.

### **TIMING TO LENTILS**

Apply *Valor* Herbicide at 1.5 to 2 oz/A when crop is physiologically mature and a minimum of 80% of the pods are yellow to tan in color and 20% are yellow in color. If lentils are treated too early, a reduction in seed quality may occur. Do not spray *Valor* Herbicide on any area of the field with a significant amount of plants with green color. Lentils can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure recommendations for postemergence application.

## DIRECTIONS FOR USE IN SUNFLOWER AND SAFFLOWER

### **HARVEST AID**

#### **RESTRICTIONS AND LIMITATIONS**

- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single application
- Do not apply more than 3 oz of *Valor* Herbicide per acre during a single growing season
- Do not harvest within 5 days of application

Desiccation from *Valor* Herbicide requires the addition of an agronomically approved adjuvant to the spray mixture. A methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A should be used. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* Herbicide with glyphosate or paraquat will increase control of emerged weeds and aid in harvest for sunflowers. Tank mixing *Valor* Herbicide with glyphosate will increase control of emerged weeds and aid in harvest for safflower.

### **TIMING TO SUNFLOWER AND SAFFLOWER**

Apply *Valor* Herbicide at 1.5 to 2 oz/A when crop is mature (when seed is 35% moisture or less). For many varieties, this is when the backs of the heads are turning yellow and the bracts are turning brown. If a sufficient amount of large weeds (harvest limiting) are present, Sunflower and safflower can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure recommendations for postemergence application.

## DIRECTIONS FOR USE IN WHEAT

[For use in the states of

DE ID KY MD MN MT NC ND NJ OR SC SD TN VA and WA Only]

### RESTRICTIONS AND LIMITATIONS

Do not apply more than 2 oz of *Valor* Herbicide per acre during a single application

- Do not apply more than 2 oz of *Valor* Herbicide per acre during a single growing season

### PRE PLANT APPLICATIONS, PRE EMERGENCE WEED CONTROL

#### RESTRICTIONS AND LIMITATIONS

- For pre plant weed control use only on no till or minimum tillage fields where the previous year's crop residue has not been incorporated into the soil
- [Plant wheat no sooner than 7 days after *Valor* Herbicide application in the states of DE ID KY MD MN MT NC ND NJ OR SC SD TN VA or WA]
- [Plant wheat no sooner than 14 days after *Valor* Herbicide application in the states of DE ID KY MD MN MT NC ND NJ OR SC SD TN VA or WA]
- [Do not use on Durum wheat]
- Do not irrigate between emergence and spike
- Wheat must be planted a minimum of 1" deep
- Do not graze until wheat has reached 5 inches in height

#### Burndown Use Directions

*Valor* Herbicide applied as part of a burndown program at 2 oz/A may be used for residual weed control as well as to assist in postemergence burndown of many weeds where wheat will be planted directly into the residue of the previous crop. See Directions for Use in Fall Burndown Programs in Fields to be Planted to Barley Field Pea Flax Lentil Safflower Sunflower and Wheat for rates and timing of applications. For control of emerged weeds *Valor* Herbicide must be applied with an appropriate burndown tank mix partner. To ensure thorough coverage use a minimum of 15 gallons of spray solution per acre. Refer to tank mix partner's label for recommended application pressure and recommended adjuvant systems.

### IPOST PLANT, PRE EMERGENCE WEED CONTROL

#### RESTRICTIONS AND LIMITATIONS

- For post plant pre emergence weed control use only on no till or minimum tillage fields where the previous crop residue has not been incorporated into the soil
- Apply *Valor* Herbicide up to 2 days after planting
- [Do not use on Durum wheat]
- Do not irrigate between emergence and spike
- Wheat must be planted a minimum of 1" deep
- Do not graze until wheat has reached 5 inches in height

#### Use Directions

*Valor* Herbicide applied at 2 oz/A may be used for residual weed control where wheat has been planted directly into the residue of the previous year. Application must be made no later than 2 days after planting.]

### HARVEST AID

#### RESTRICTIONS AND LIMITATIONS

- Do not harvest within 10 days of application

#### Use Directions

*Valor* Herbicide applied at 2 oz/A for desiccation requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lbs/A or a 28 to 32% nitrogen solution at 1 to 2 qts/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* Herbicide with glyphosate will increase control of emerged weeds and aid in harvest.

To ensure thorough coverage use a minimum of 10 gallons spray solution per acre by ground application and a minimum of 5 gallons per acre by aerial application. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for postemergence application.

### TIMING TO WHEAT

Apply *Valor* at 1.5 to 2 oz/A after wheat reaches the hard dough stage and grain has no more than 30% moisture. Wheat can be harvested 10 days after application. Valent recommends tank mixing with glyphosate.

PLEASE CONTACT VALENT U S A CORPORATION AT 800 6 VALENT (682 5368) TO  
DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE

Copyright © 2012 by Valent U S A Corporation

*Valor* is a registered trademark of Valent U S A Corporation

Manufactured for  
**Valent U S A Corporation**  
P O Box 8025  
Walnut Creek CA 94596 8025  
[www.valent.com](http://www.valent.com)

Made in U S A

059639 00119 20110914 Valor wheatupdate