



ESTEEM[®] Insect Growth Regulator

Active Ingredient	By Wt.
*Pyriproxyfen (2-[1-methyl-2-(4-phenoxyphenoxy) ethoxy]pyridine)	33%
Other Ingredients	67%
Contains 2.9 pound a.i. per gallon.	
Contains aromatic petroleum distillates.	

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS 1 GALLON

ACCEPTED
 NOV 23 1999
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 59639-104

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye irritation. Causes skin irritation. Do not get in eyes or on clothing. Avoid contact with skin. Harmful if absorbed through skin, if swallowed or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT:

- If in eyes:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.
- If on skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.
- If swallowed:** Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.
- If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
- Note to Physician:** Probable mucosal damage may contraindicate the use of gastric lavage. This product contains a light petroleum solvent and an aspiration hazard may exist. Emergency information - call 1-800-892-0099.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: coveralls over short-sleeved shirt, short pants, chemical-resistant gloves, such as Barrier Laminate or Viton \geq 14 mils, socks, chemical-resistant footwear, protective eyewear, for overhead exposure chemical-resistant headgear and for mixers/loaders a chemical-resistant apron.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment washwaters or rinsate. Avoid direct application and/or spray drift to bee hives.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over short sleeve shirt, short pants, chemical-resistant gloves, such as Barrier Laminate or Viton \geq 14 mils, socks, chemical-resistant footwear, and protective eyewear.

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. EXCEPT AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

In no event shall Valent or Seller be liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is later, so that an immediate inspection of the affected property and growing crops can be made.

If Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

GENERAL INFORMATION

Do not apply this product through any type of irrigation system.

ESTEEM Insect Growth Regulator is intended for use in Integrated Pest Management (IPM) or Integrated Resistance Management (IRM) programs. ESTEEM Insect Growth Regulator will not control insect adults, and it is recommended to be used in combination and/or rotation with other IPM or IRM materials. Contact your local state extension service for details.

ESTEEM Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM Insect Growth Regulator has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM Insect Growth Regulator is an Insect Growth Regulator (IGR) and activity depends on the insects' development, evidence of activity will be slower than typical contact insecticides.

COTTON

GENERAL

ESTEEM Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM is an Insect Growth Regulator (IGR) and activity depends on the insects development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	DOSAGE PER ACRE	SPECIAL INSTRUCTIONS
Cotton	Silverleaf Whitefly Sweetpotato Whitefly	2.5 to 3.0 fl. oz. (25 to 30 gms. ai) Do not apply less than 2.5 fl. oz./A	Apply with air or ground equipment in adequate water for uniform coverage: By Air: 3 to 10 gals./A By Ground: 10 to 50 gals./A HIGHER WATER VOLUMES PROVIDE IMPROVED INSECT CONTROL. Make only one application per growing season. Begin applications when whiteflies reach threshold levels (3 to 5 adults per leaf and/or equivalent levels of immatures). Under severe pressure, use the higher recommended rates. To support resistance management only whole fields should be treated.
<p>ESTEEM Insect Growth Regulator exhibits translaminar movement in leaves, which means that ESTEEM Insect Growth Regulator applied to the upper surface of the leaves will move to the lower surface of the leaf where whitefly are present.</p> <ul style="list-style-type: none"> Do not apply within 28 days of harvest. Do not apply more than 3.0 fl. oz. (30 gms. ai) ESTEEM Insect Growth Regulator per acre per season. Do not make more than 1 application of ESTEEM Insect Growth Regulator per growing season. Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after last application. 			

MANAGING WHITEFLY RESISTANCE: Repeated use of the same class of insecticides or insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting ESTEEM Insect Growth Regulator applications to a single application per growing season. ESTEEM Insect Growth Regulator however, may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 2.5 fl. oz. per acre for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local, state, and federal IPM and Integrated Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

APPLE

GENERAL

ESTEEM Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM is an Insect Growth Regulator (IGR) and activity depends on the insects development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	DOSAGE PER ACRE	SPECIAL INSTRUCTIONS
Apple	Codling Moth	4.0 to 5.0 fl. oz. (40 to 50 gms. ai)	1st Generation Codling Moth: APPLY JUST PRIOR TO EGG HATCH. Make a second application 14 to 21 days later. Apply with ground equipment in 100 - 400 gals./A. For additional generations apply other insecticides registered for Codling Moth control.
	San Jose Scale	3.0 to 5.0 fl. oz. (30 to 50 gms. ai)	Apply at delayed dormant with oil or in cover sprays without oil. Use the manufacturers recommended rate of oil. Under heavy scale pressure, use the higher rate. Make application in cover sprays when scale crawlers first emerge.
	Spotted Tentiform Leafminer APHID SUPPRESSION: Green Apple Aphid Rosy Apple Aphid Spirea Aphid	3.0 to 4.0 fl. oz. (30 to 40 gms. ai)	Apply when insects first appear. Usually this occurs in Prepink apple stage for the first generation. Make a separate application for the second generation if necessary. Application made for Spotted Tentiform Leafminer will aid in the suppression of aphids. Apply with ground equipment in 100 - 400 gals./A.
	<p>ESTEEM Insect Growth Regulator may be mixed and/or alternated with commonly used insecticides to comply with local IPM programs and resistance management programs.</p> <ul style="list-style-type: none"> • Do not apply within 45 days of harvest. • Do not exceed 10.0 fl. oz. (100 gms. ai) between petal fall and harvest or 15.0 fl. oz. (150 gms. ai) of ESTEEM Insect Growth Regulator per acre per season. • Do not apply earlier than 14 days after last ESTEEM treatment. 		

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM Insect Growth Regulator to two applications per growing season. ESTEEM may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

PEAR

GENERAL

ESTEEM Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM is an Insect Growth Regulator (IGR) and activity depends on the insects development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	DOSAGE PER ACRE	SPECIAL INSTRUCTIONS
Pear	Pear Psylla San Jose Scale	4.0 to 5.0 fl. oz. (40 to 50 gms. ai)	Apply at delayed dormant with a Supreme oil at manufactures recommended rates with ground equipment in 100 to 250 gals./A. For Pear Psylla a second application may be made at petal fall in addition to the delayed dormant application. <i>For control of scale in season make application in cover sprays when scale crawlers first emerge.</i>
	Codling Moth San Jose Scale (crawler stage)	4.0 to 5.0 fl. oz. (40 to 50 gms. ai)	First Generation Codling Moth: APPLY JUST PRIOR TO EGG HATCH. Make a second application 14 to 21 days later. Apply with ground equipment in 100 to 400 gals./A. <i>For additional generations apply other insecticides registered for Codling Moth control.</i>
<p>ESTEEM Insect Growth Regulator may be mixed and/or alternated with commonly used insecticides to comply with local IPM and resistance management programs.</p> <ul style="list-style-type: none"> • Do not apply within 45 days of harvest. • Do not exceed 10.0 fl. oz. (100 gms. ai) between petal fall and harvest or 15.0 fl. oz. (150 gms. ai) of ESTEEM Insect Growth Regulator per acre per season. • Do not apply earlier than 14 days after last ESTEEM treatment. 			

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM Insect Growth Regulator to two applications per growing season. ESTEEM may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

WALNUT

GENERAL

ESTEEM Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM is an Insect Growth Regulator (IGR) and activity depends on the insects development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	DOSAGE PER ACRE	SPECIAL INSTRUCTIONS
Walnut	Codling Moth	4.0 to 5.0 fl. oz. (40 to 50 gms. ai)	1st Generation Codling Moth: APPLY JUST PRIOR TO EGG HATCH. Make a second application 14 to 21 days later. Apply with ground equipment in 100 - 400 gals./A.
	Codling Moth Suppression of Navel Orangeworm	4.0 to 5.0 fl. oz. (40 to 50 gms. ai)	Late Season Codling Moth: Apply at egg deposition. Make a second application if needed 14 to 21 days later.
	<p>ESTEEM Insect Growth Regulator may be mixed and/or alternated with commonly used insecticides to comply with local IPM programs and resistance management programs.</p> <ul style="list-style-type: none"> • Do not apply within 21 days of harvest. • Do not exceed 10.0 fl. oz. (100 gms. ai) between petal fall and harvest or 15.0 fl. oz. (150 gms. ai) of ESTEEM Insect Growth Regulator per acre per season. • Do not apply earlier than 14 days after last ESTEEM treatment. 		

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM Insect Growth Regulator to two applications per growing season. ESTEEM may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

ALMOND

GENERAL: ESTEEM Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	DOSAGE PER ACRE	SPECIAL INSTRUCTIONS
Almond	San Jose Scale Suppression of: Peach Twig Borer	4.0 to 5.0 fl. oz. (40 to 50 gms. ai)	Apply with ground equipment in 100 to 400 gals./A. Apply at delayed dormant or in cover sprays. Delayed Dormant: Use oil at 1-1/2 to 2 gals./100 gals. as dilute spray or 4 to 6 gals./acre as concentrate spray. Follow the manufacturers' directions and recommendations on timing and rates. Under heavy scale pressure, use the higher rate. Make application in cover sprays when scale crawlers first emerge. Follow local recommendations regarding in season use and rates of oil in your area. ESTEEM has provided good suppression of Peach Twig Borer when applied delayed dormant through bloom sprays.
<p>ESTEEM Insect Growth Regulator may be mixed and/or alternated with commonly used insecticides to comply with local IPM programs and resistance management programs.</p> <ul style="list-style-type: none"> • Do not apply within 21 days of harvest. • Do not exceed 49 fl. oz. (150 gms. ai) of ESTEEM Insect Growth Regulator per acre per season. • Do not apply earlier than 14 days after last ESTEEM treatment. 			

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM Insect Growth Regulator to three applications per growing season. ESTEEM may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

CITRUS

GENERAL: ESTEEM Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM is an Insect Growth Regulator (IGR) and activity depends on the insects development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	DOSAGE PER ACRE	SPECIAL INSTRUCTIONS
Citrus	Scale California Red Black Brown Soft	5.0 fl. oz. (50 gms. ai)	<p>Apply with ground equipment in 200 to 1500 gals./A.</p> <p>Apply anytime prior to fruit becoming infested when scale is in the crawler stage. Thorough coverage is essential for good control.</p> <p>ESTEEM IGR may be applied with superior spray oils at 1/2 to 1%. Follow the manufacturers' directions and recommendations on timing and rates.</p> <p>NOTE: For scale apply only one application during emergence of scale crawlers per calendar year.</p>
	Citrus Whitefly Citrus Leafminer Citrus Black Fly	2.5 to 3.0 fl. oz. (25 to 30 gms. ai)	<p>Apply with ground equipment in 200 to 1500 gals./A.</p> <p>Apply when pests are just beginning to build and before infestation has occurred. Thorough coverage is essential for good control.</p> <p>ESTEEM IGR may be mixed and/or alternated with commonly used insecticides to comply with local IPM programs and resistance management programs.</p>
<ul style="list-style-type: none"> • Do not apply within 1 day of harvest. • Do not exceed 49 fl. oz. (150 gms. ai) of ESTEEM Insect Growth Regulator per acre per season. • Do not make more than three (3) ESTEEM applications per season. • Do not apply earlier than 21 days after last ESTEEM treatment. 			

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM Insect Growth Regulator to three applications per growing season. ESTEEM may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

PEPPER

GENERAL: ESTEEM Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM is an Insect Growth Regulator (IGR) and activity depends on the insects development, evidence of activity will be slower than typical contact insecticides.

ESTEEM does exhibit translaminar movement in pepper leaves which means that ESTEEM applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROP	PESTS	DOSAGE PER ACRE	SPECIAL INSTRUCTIONS
Pepper	Sweetpotato/Silverleaf Whitefly (immatures only)	2.5 to 3.0 fl. oz. (25 to 30 gms. ai)	EARLY SEASON: Apply with ground equipment in adequate water for uniform coverage (10 to 50 gals./A). Higher water volumes may provide improved insect control. Make application when pest activity is first noticed.
	Sweetpotato/Silverleaf Whitefly (immatures & adults) Cabbage Looper Green Peach Aphid Tobacco Hornworm	2.0 to 2.5 fl. oz. (20 to 25 gms. ai) + ORTHENE® 75 S Soluble Powder (0.5 - 1.0 lb. ai)	Apply as a tank mix with ground equipment for uniform coverage (25 to 150 gals./A by ground. Make application when eggs or insects first appear. Do not make a second ESTEEM application sooner than 30 days following the first ESTEEM application. If control is needed during this period use an alternate insecticide registered for the target pest. ESTEEM may also be mixed or alternated with commonly used insecticides to comply with local IPM programs. Comply with all applicable directions, restrictions and precautions on the registered labels for ORTHENE 75S Soluble Powder.
<ul style="list-style-type: none"> • Do not apply within 14 days of harvest. • Do not exceed 20 fl. oz. (80 gms. ai) of ESTEEM Insect Growth Regulator per acre per season. • Do not make more than three (3) ESTEEM applications per growing season. • Do not apply earlier than 14 days after last ESTEEM treatment. • Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after last application. 			

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM Insect Growth Regulator to three applications per growing season. ESTEEM may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before

applying any insecticide.

TOMATO

GENERAL: ESTEEM Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM is an Insect Growth Regulator (IGR) and activity depends on the insects development, evidence of activity will be slower than typical contact insecticides.

ESTEEM does exhibit translaminar movement in tomato leaves which means that ESTEEM applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROP	PESTS	DOSAGE PER ACRE	SPECIAL INSTRUCTIONS
Tomato	Sweetpotato/Silverleaf Whitefly (immatures only)	2.5 to 3.0 fl. oz. (25 to 30 gms. ai)	EARLY SEASON: Apply with ground equipment in adequate water for uniform coverage (10 to 50 gals./A). Higher water volumes may provide improved insect control. Make application when pest activity is first noticed.
	Sweetpotato/Silverleaf Whitefly (immatures & adults) Tomato Pinworm Fruitworms Potato Aphid Thrips Yellowstriped Armyworm	2.0 to 2.5 fl. oz. (20 to 25 gms. ai) + MONITOR [®] 4 Spray 24 oz. (0.75 lb. ai) + DANITOL [®] 2.4 EC 10 2/3 oz. (0.2 lb. ai)	EARLY SEASON WHITEFLY CONTROL PROGRAM FOR REDUCTION OF GEMINIVIRUS: Apply as a tank mix with ground equipment for uniform coverage (10 to 120 gals./A). Make application when scouting first indicates adult whiteflies are on young plants. NOTE: This may result in treatment immediately following transplanting. MID TO LATE SEASON WHITEFLY CONTROL PROGRAM FOR PREVENTION OF IRREGULAR FRUIT RIPENING: Apply as a tank mix with ground equipment for uniform coverage (20 to 120 gals./A by ground). Make application when scouting indicates whitefly are present. Do not make a second ESTEEM application sooner than 30 days following the first ESTEEM application. If control is needed during this period use an alternate insecticide registered for the target pest. ESTEEM may also be mixed or alternated with commonly used insecticides to comply with local IPM programs. Comply with all applicable directions, restrictions and precautions on the registered labels for MONITOR 4 Spray and DANITOL 2.4 EC.
<ul style="list-style-type: none"> • Do not apply within 14 days of harvest. • Do not exceed 20 fl. oz. (80 gms. ai) of ESTEEM Insect Growth Regulator per acre per season. • Do not make more than three (3) ESTEEM applications per growing season. • Do not apply earlier than 14 days after last ESTEEM treatment. • Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after last application. 			

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM Insect Growth Regulator to three applications per growing season. ESTEEM may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Store in a cool dry place.

Keep pesticide in original container.

Keep container closed when not in use.

Do not put concentrate or dilute into food or drink containers.

Not for use or storage in or around the home.

For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-892-0099.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Do not reuse container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Copyright© 1999 by Valent U.S.A. Corporation

DANITOL® - Reg. TM of Sumitomo Chemical Company, Ltd.

KNACK® - Reg. TM of Valent U.S.A. Corporation

MONITOR® - Reg. TM of Valent U.S.A. Corporation

ORTHENE® - Reg. TM of Monsanto Company

Manufactured for:

Valent U.S.A. Corporation

P.O. Box 8025

Walnut Creek, CA 94596-8025

Made in U.S.A.

Form 991110 - ESTEEM-8.REG

EPA Reg. No. 59639-104

EPA Est.