

59639-105

11/22/2002

1/15

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

 Registration
 Amendment
 Other

OPP Identifier Number
283410

Application for Pesticide - Section I

1. Company/Product Number 59639-105	2. EPA Product Manager Marion Johnson	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Reimment Herbicide	PM# 10	
5. Name and Address of Applicant (Include ZIP Code) Valent U.S.A. Corporation 1333 N. California Ave. Suite 600 Walnut Creek, CA 94596-8025 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION
NOV 22 2002

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION: ALTERNATE BRAND NAME OF: VELOCITY Herbicide

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt. No. per container		<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Linda Obrestad	Title Labeling Specialist	Telephone No. (Include Area Code) (925) 256-2791
2. Signature <i>Linda Obrestad</i>		3. Title Labeling Specialist
4. Typed Name Linda Obrestad		5. Date November 18, 2002
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)

2/15

1333 North California Blvd.
Suite 600
P.O. Box 8025
Walnut Creek, CA 94596-8025
(925) 256-2700



November 18, 2002
Regiment™ Herbicide
EPA Reg No. 59639-105
Notification: Alternate Brand Name of
Velocity® Herbicide
OPP. ID.No. 283410

NOTIFICATION
NOV 22 2002

CERTIFICATION STATEMENT

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

A handwritten signature in black ink that reads "Linda Obrestad".

Linda Obrestad
Labeling Specialist
Valent U.S.A. Corporation



Regiment™ Velocity® (HERBICIDE)

NOTIFICATION

NOV 22 2002

Active Ingredient	By Wt.
*Bispyribac-sodium	80.0%
Other Ingredients	<u>20.0%</u>
	100.0%


*Sodium 2,6-bis[(4,6-dimethoxypyrimidin-2-yl)oxy]benzoate

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS



4/15

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Avoid breathing dust or spray mist.

FIRST AID

- If on swallowed :**
 - Call poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- If in eyes:**
 - Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- If on skin or clothing:**
 - Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- If inhaled:**
 - Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-892-0099** for emergency medical treatment information

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow instructions for category A on an EPA chemical resistant category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical resistant gloves such as Barrier Laminate or Butyl Rubber ≥ 14 mils or Nitril Rubber ≥ 14 mils or Viton ≥ 14 mils, shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

5/15

ENVIRONMENTAL HAZARDS

This product is toxic to non-target plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves, such as Barrier Laminate or Butyl Rubber \geq 14 mils or Nitril Rubber \geq 14 mils or Viton \geq 14 mils and shoes plus socks.

6/15

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. EXCEPT AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

In no event shall Valent or Seller be liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

If Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

7/15

GENERAL INFORMATION

Do not apply this product through any type of irrigation system.

~~REGIMENTVELOCITY~~ is a selective, postemergence contact herbicide which provides control of many weeds infesting rice. It has an exceptionally wide window of application and should be used as an integral part of a weed control program in conjunction with a resistance management strategy.* The mode of action is the inhibition of the acetolactate synthase (ALS) enzyme, and thus, activity is relatively slow, 14 to 21 days for complete control. Susceptible weeds turn yellow and stop growing 3 to 7 days after treatment. Browning of sensitive weeds is evident in 7 to 14 days after treatment with death of the stem and roots occurring within 14 to 21 days after treatment. ~~REGIMENTVELOCITY~~ is not a residual/soil active herbicide and will not prevent reinfestation of weeds which germinate after application. ~~REGIMENTVELOCITY~~ is rainfast 8 hours after application. Temporary injury to rice may occur after application, but injury is transient and yields are not adversely affected. Fertilizer top-dressing will speed temporary injury recovery.

*See resistance management statement on this label.

8/15

Table 1


PRODUCT USE RATE/WEED FOR USE IN RICE GROWING REGIONS (EXCEPT CALIFORNIA)			
Weeds Controlled	Scientific Name	Weed Size	Rate Acres/Packet (grams product/Acre)
Barnyardgrass (including propanil and/or Facet® resistant barnyardgrass)	<i>Echinochloa crus-galli</i>	1 leaf up to 3 tillers	5 to 3.75 (11.25g. - 15g.)
Junglerice	<i>Echinochloa colonum</i>	1 leaf up to 3 tillers	
Johnsongrass	<i>Sorghum halepense</i>	3 to 24 inches	
Hemp Sesbania	<i>Sesbania exaltata</i>	3 to 18 inches	
Indian Jointvetch	<i>Aeschynomene indica</i>	3 to 18 inches	
Northern Jointvetch	<i>Aeschynomene virginica</i>	3 to 18 inches	
Annual Rice Flatsedge	<i>Cyperus iria</i>	1 leaf up to 4 leaf	
Gooseweed	<i>Sphenoclea zeylanica</i>	1 leaf up to 4 leaf	
Dayflower	<i>Commelina communis</i>	1 leaf up to 4 leaf	
Waterhyssop	<i>Bacopa rotundifolia</i>	1 leaf up to 4 leaf	
Ducksalad	<i>Heteranthera spp.</i>	1 leaf up to "spoon leaf"	
Pennsylvania Smartweed	<i>Polygonum pennsylvanicum</i>	1 to 4 inches	3.125 (18g.)
Late Application Barnyardgrass Suppression	<i>Echinochloa crus-galli</i> <i>Echinochloa colonum</i>	4 tiller up to booting	

Weeds Suppressed	Scientific Name	Weed Size	Rate Acres/Packet (grams product/Acre)
Alligatorweed	<i>Alternanthera philoxeroides</i>	Up to 10 inch runners	3.75 (15g.)
Knotgrass-POST FLOOD ONLY	<i>Paspalum ditichum</i>	Up to heading	
Perennial Barnyardgrass	<i>Echinochloa polystachya</i>	Up to 2 tillers	
Redstem	<i>Ammannia spp.</i>	1 to 4 inches	5 to 3.75 (11.25g. - 15g.)
Pennsylvania smartweed	<i>Polygonum pennsylvanicum</i>	4 to 24 inches	
Pigweeds	<i>Amaranthus spp.</i>	1 to 12 inches	
Pitted Morningglory	<i>Ipomoea lacunosa</i>	1 to 4 inches	
Entireleaf Morningglory	<i>Ipomoea hederacea</i>	1 to 4 inches	
Texas/Mexicanweed	<i>Caperonia spp.</i>	1 leaf up to 4 leaf	
Eclipta	<i>Eclipta spp.</i>	1 leaf up to 4 leaf	

Adjuvant: Application of "ECHO 9000" must include a surfactant unless otherwise specified in another section of this label or in supplemental labeling. Refer to the Valent bulletin "Approved Surfactants for use with ECHO 9000" for a list of approved surfactants and

9/15

rates. Use of surfactants other than the ones specified is done at the sole risk of the user.



10/15

DRY-SEEDED OR WATER-SEEDED RICE - U.S. RICE GROWING REGIONS (Except California)

- **REGIMENTVELOCITY** may be applied to rice after the 3-leaf (3rd leaf fully expanded) stage of development until the panicle initiation (green ring/just prior to joint movement) stage of development. Do not apply to rice before the 3rd leaf is fully expanded or after panicle initiation. Regardless of seeding method, rice must have the 3rd leaf fully expanded and the root system must be completely below the soil surface prior to **REGIMENTVELOCITY** application. Medium grain varieties may be more sensitive to **REGIMENTVELOCITY** than long grain varieties. Pubescent (hairy) leaf varieties may be more sensitive to **REGIMENTVELOCITY** than glabrous (smooth) leaf varieties.
- **Pre-Flood Application:** Following application, wait at least one day for herbicide uptake, then establish the permanent flood as soon as the rice will tolerate flooding. Under conditions in which the permanent flood is delayed, flush as necessary to maintain rice growth. Reinfestation of weeds may occur if a permanent flood is not established in a timely manner.
- **Post-Flood Application:** Prior to application, the floodwater must be lowered so that at least 70% of the weed plant surface is above the floodwater. Bring the field to normal flood level 2 to 3 days after application.
- Refer to Table 1: "Product Use Rate/Weed".
- Use the upper end of the recommended use rate range when weed populations are approaching the maximum controllable size and/or weed infestation is severe.
- Multiple applications of **REGIMENTVELOCITY** may be made as long as the total seasonal amount does not exceed 30 grams of product (24 grams ai) per acre and at least 3 weeks elapse between applications.
- **Late Application Barnyardgrass Suppression:** When barnyardgrass develops to stages between 4-tiller and booting, a negative influence on yield has already occurred. Controlling or suppressing barnyardgrass at these stages will maximize the remaining yield potential and reduce weed seed production.
- **Suppression of Knotgrass:** Make application after the rice is in permanent flood and 70% of the knotgrass is above the flood level. Make application prior to knotgrass heading.

Table 2

PRODUCT USE RATE/WEED FOR USE IN CALIFORNIA			
Weeds Controlled	Scientific Name	Weed Size	Rate Acres/Packet (grams Product/Acre)
Watergrass	<i>Echinochloa crus-galli</i> var. <i>oryzicola</i>	1 leaf up to 2 tillers	4.5 to 3.75 (12g. - 15g.)
	<i>Echinochloa oryzoides</i>	1 leaf up to 2 tillers	
Barnyardgrass	<i>Echinochloa crus-galli</i>	1 leaf up to 3 tillers	
Watergrass resistant biotypes	Including <i>Echinochloa phyllopogon</i> , <i>Echinochloa crus-galli</i> var. <i>oryzicola</i> <i>Echinochloa oryzoides</i>	1 to 2 tillers	3 to 2.5 (18.75g. - 22.5g.)
Ricefield Bulrush	<i>Scirpus mucronatus</i>	2 leaf up to flower initiation	3.75 (15g.)
California Arrowhead	<i>Sagittaria montevidensis</i> spp. <i>Calycina</i>	1 leaf up to flower initiation	4.5 to 3.75 (12g. - 15g.)
Waterhyssop	<i>Bacopa rotundifolia</i>	1 leaf up to 4 leaf	
Ducksalad	<i>Heteranthera</i> spp.	1 leaf up to "spoon leaf"	
Waterplantain	<i>Alisma triviale</i>	1 leaf up to flower initiation	
Adjuvant: Application of REGIMENTVELOCITY must include a surfactant unless otherwise specified in another section of this label or in supplemental labeling. Refer to the Valent bulletin "Approved Surfactants for use with REGIMENTVELOCITY" for a list of approved surfactants and rates. Use of surfactants other than the one specified is done at the sole risk of the user.			

11/15

DRY-SEEDED OR WATER-SEEDED (California)

- ~~REGIMENTVELOCITY~~ may be applied to rice after the 3-leaf stage of development until the panicle initiation (green ring/just prior to joint movement) stage of development. Do not apply to rice before the 3rd leaf is fully expanded or after panicle initiation. Regardless of seeding method, rice must have the 3rd leaf fully expanded and the root system must be completely below the soil surface prior to ~~REGIMENTVELOCITY~~ application.
- **Pinpoint or Leathers Flood Culture:** After seeding rice into the initial (seedling) flood, drain the field when the root is approximately ¼ inch long, but before shoot growth has been initiated. This will allow the root to peg (tack) to the soil. Apply ~~REGIMENTVELOCITY~~ after draining when the rice seedling is in at least the 3-leaf (fully expanded) stage and the root system is completely covered with soil. Bring the field to pinpoint flood level 2 to 3 days after application. Prolonged drainage can stress the rice plant and/or allow for subsequent weed germination which could potentially result in yield reduction.
- **Dry-Seeded Pre-Flood Culture:** Following application, wait at least one day for herbicide uptake then establish the permanent flood as soon as the rice will tolerate flooding. Under conditions in which flooding is delayed, flush as necessary to maintain rice growth. Reinfestation of weeds may occur if a permanent flood is not established in a timely manner.
- **Post-Flood Application:** Prior to application, the floodwater must be lowered so that 70% of the weed plant surface is above the floodwater. Bring the field to normal flood level 2 to 3 days after application. Refer to Table 2: "Product Use Rate/Weed".
- Use the upper end of the recommended use rate range when weed populations are approaching the maximum controllable size and/or weed infestation is severe.
- Multiple applications of ~~REGIMENTVELOCITY~~ may be made as long as the total seasonal amount does not exceed 30 grams of product (24 grams ai) per acre and at least 3 weeks elapse between applications.
- **Herbicide Resistant Watergrass Biotypes (Early and Late):** In localized specific areas, these species have exhibited resistance to various herbicides. If resistant species have been documented in your field(s), consult your crop advisor or local extension service for further information. Resistance management will prolong the usefulness of all rice herbicides.
- Do not use ~~REGIMENTVELOCITY~~ on the rice variety CM 101. Pubescent (hairy) leaf varieties are more sensitive to ~~REGIMENTVELOCITY~~ than glabrous (smooth) leaf varieties.

TANK MIX APPLICATIONS

~~REGIMENTVELOCITY~~ may be tank mixed with BOLERO® 8 EC, ABOLISH™, PROWL® 3.3 EC, COMMAND®, 3ME, WHIP® 360, Dimilin®, FACET®, Permit®, GRANDSTAND® R, or GRANDSTAND® CA, 2,4-D, MCPA, BLAZER®, and KARATE®, or KARATE® Z. Tank mixing with AIM may cause antagonism to the activity of ~~REGIMENTVELOCITY~~. If this tank mixture is used, increase the rate of ~~REGIMENTVELOCITY~~ 1 packet for 4 acres to 1 packet for 3.75 acres and do not exceed 1.5 oz. of AIM per acre. Read and carefully observe the label claims, cautionary statements, rates and all other information on the labels of products to be used in tank mixture. Use according to the most restrictive label directions of each product in the mixture. DO NOT tank mix ~~REGIMENTVELOCITY~~ with malathion, methyl parathion, propanil or herbicidal mixtures which contain propanil because antagonism and/or injury will occur. Tank mixing or use of ~~REGIMENTVELOCITY~~ with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of the user, applicator and/or application advisor. For further information regarding tank mixtures see the "Resistance Management" section of the label.

METHOD OF APPLICATION

Uniformly apply ~~REGIMENTVELOCITY~~ or ~~REGIMENTVELOCITY~~ tank mixes by aircraft in no less than 10 gallons of water per acre total spray volume or by ground equipment in 10 to 20 gallons of water per acre total spray volume. Any factor, such as reduced spray volume, which adversely affects coverage and canopy penetration will have a negative effect on the performance of ~~REGIMENTVELOCITY~~. Use nozzle types and nozzle arrangements which will provide maximum coverage and minimize the potential for off target movement of spray particles. Refer to label section "Spray Drift".

12/15

MIXING AND SPRAYING EQUIPMENT PREPARATION AND CLEANUP

PRECAUTION: DO NOT USE CHLORINE BLEACH WITH AMMONIA. REMOVE ALL TRACES OF LIQUID FERTILIZER CONTAINING ANY FORM OF AMMONIA OR AMMONIUM BEFORE ADDING ANY CHLORINE SOURCE SUCH AS CHLORINE BLEACH.

Prior to using REGIMENTVELOCITY, thoroughly drain, clean, and rinse all mixing and spraying equipment that will come in contact with REGIMENTVELOCITY. Follow the cleanup procedures recommended by the manufacturer of the previously sprayed product. Failure to remove all deposits of previously sprayed products may result in collection of REGIMENTVELOCITY residues and inhibit cleanup of mixing and spraying equipment after REGIMENTVELOCITY use.

Residual amounts of herbicide in or on mixing or spraying equipment may have an adverse effect on subsequently sprayed crops. Thoroughly drain, clean and rinse all mixing and spraying equipment including tanks, booms, hoses, strainers, screens, and nozzles immediately after use. Use the following procedure:

1. Remove all physical residues
2. Thoroughly drain and rinse tanks, booms, and hoses with clean water.
3. Fill the tank one half full of clean water and use a spraying/mixing tank cleaner that DOES NOT contain chlorine. Fill the remainder of the tank with clean water. Let agitate/recirculate according to the directions of the cleaner manufacturer. Thoroughly flush the boom and hoses before draining.
4. Rinse all hoses, tanks, nozzles, strainers, and booms with clean water to remove the tank cleaner. Follow the directions provided by the tank cleaner manufacturer.
5. Fill the tank half full of clean water and add one (1) gallon of 3% active household ammonia for every 100 gallons of water the tank will hold. Fill the remainder of the tank with clean water and allow the solution to agitate/recirculate for 15 minutes. Thoroughly flush the ammonia cleaning solution through the boom, hoses, nozzles, screens, and strainers before draining the tank.
6. Remove the strainers, nozzles, and screens and clean separately in a solution of household ammonia and water.
7. Replace the strainer(s), nozzles, and screens.
8. Repeat Step 5.
9. Thoroughly rinse the tank with clean water and flush the water through the boom, nozzles and hoses in order to remove all traces of ammonia.
10. Dispose of the rinsate on site or at an approved waste disposal facility.

SPRAY DRIFT

Do not allow spray from ground or aerial equipment to drift onto adjacent land or crops. When drift may be a problem, do everything possible to reduce spray drift, including:

- Do not spray if wind speeds are or become excessive. Do not spray if wind speed is 8 MPH or greater. If sensitive crops or plants are downwind, extreme caution must be used under all conditions. Do not spray if winds are gusty.
- Use extreme caution when conditions are favorable for drift (high temperatures, drought, low relative humidity), especially when sensitive plants are located nearby.
- Do not apply when a temperature inversion exists. If inversion conditions are suspected, consult with local weather services before making an application.
- Further reductions in drift can be obtained by:
 - Using large droplet size sprays. Do not use nozzles that produce small droplets. Orient nozzles downward and slightly backward as needed to reduce drift for ground applications.
 - Orienting nozzles straight back with the windstream, using straight stream orifices for aerial applications.
 - Use the lowest number of nozzles practical with the largest possible orifice size to obtain the minimum 10 GPA volume. Application height and boom length should be set according to manufacturer's instructions to minimize drift.
 - Increasing the volume of spray mixture (for example a minimum of 10-20 GPA for ground applications) by using higher flow rate nozzles. Using lower pressure with the appropriate nozzle to obtain higher volumes will also reduce drift.
 - Apply as close to target plants as practical while maintaining a good spray pattern for adequate coverage.

13/15

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

MIXING INSTRUCTIONS

Water Soluble Packaging

This bag contains water soluble packets of REGIMENTVELOCITY. Do not handle the packets with wet gloves or allow the packets to become wet prior to mixing. If all packets are not used, close and reseal outer container to protect remaining packet(s). Do not add any liquid fertilizers, micronutrients or adjuvants to the spray solution until after the water soluble packets and their contents have completely dissolved. Water soluble packet(s) should completely dissolve in approximately five minutes. Dissolution rate may be slowed by cold water, lack of agitation, or water containing high concentrations of boron or sulfur. High concentration of boron or sulfur may result in spray screen or nozzle clogging due to the incomplete dissolution of the water soluble packet material.

- Partially fill tank with water to approximately the half way mark.
- Begin agitation
- Add REGIMENTVELOCITY water-soluble packets and make sure that they have dissolved completely before proceeding.
- Add approved surfactant*
- Add tank mix partner (if any) in the following order
 - water soluble packets (preferably added before the surfactant)
 - water dispersible granules/wettable powder
 - soluble powders
 - suspension concentrate
 - emulsifiable concentrate
- Fill remainder of tank

*If foaming is anticipated, add defoamer prior to the addition of the surfactant.

RESISTANCE MANAGEMENT

Any weed population may contain or develop plants naturally resistant to herbicides in various mode of action classes. Resistant biotypes may eventually dominate the weed population if the same class of chemistry/mode of action herbicides are used repeatedly in the same field or in successive years. These resistant biotypes may not be adequately controlled by herbicides in a mode of action class for which resistance has developed. A gradual or total loss of weed control may occur over time. Other resistance mechanisms that are not linked to site of action, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

TO DELAY HERBICIDE RESISTANCE

- Avoid the use of herbicides that have a similar target site mode of action in consecutive years. Herbicide use should be based on an IPM program that includes scouting, record keeping, and consideration of cultivation practices, water management, weed free crop seed, crop rotation, and other chemical or cultural control practices.
- Monitor treated weed population for resistance development and report suspected resistance.
- Contact your local extension or crop expert (advisor) for any additional pesticide resistance management and/or IPM recommendations for specific crops and weed biotypes.
- For further information contact Valent USA Corporation at the following toll free number 1-800-682-5368.

USE PRECAUTIONS

- Water drained directly from treated fields must not be used to irrigate other crops.
- DO NOT overlap or double spray ends of field.
- DO NOT apply more than 30 grams of REGIMENTVELOCITY (24 grams ai) per acre per year. DO NOT apply to second crop (stubble/ratoon crop) rice.
- REGIMENTVELOCITY is a contact herbicide which is not soil active and does not provide residual activity. Reinfestation of weeds may occur if a permanent flood is not established in a timely manner.
- Any environmental (ie, temperature, drought, etc.) or other stress (ie, herbicide injury, fertilizer injury or nutrient deficiencies, etc.) factors which decrease plant metabolism and growth may reduce

14/15

REGIMENTVELOCITY efficacy and increase rice injury. DO NOT APPLY TO STRESSED RICE.

- Temporary injury, chlorosis and/or stunting may occur after application but injury is transient. Fertilizer top-dressing will speed temporary injury recovery. Medium grain varieties may be more sensitive than long grain varieties. Pubescent (hairy) leaf varieties may be more sensitive to RegimentVelocity than glabrous (smooth) leaf varieties.
- Water-seeded rice that has not fully pegged (rice root system not completely below the soil surface) is susceptible to significant injury from REGIMENTVELOCITY, regardless of number of leaves.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Store in a cool dry place.

Keep pesticide in original container.

Keep container closed when not in use.

Do not put concentrate or dilute into food or drink containers.

Not for use or storage in or around the home.

For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-892-0099.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Do not reuse the outer bag. Dispose of outer bag in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke.

Copyright © 2001 by Valent U.S.A. Corporation

AIM™ - Trademark of FMC Corporation
 ABOLISH™ - Trademark of Valent USA Corporation
 BLAZER® - Reg. TM of BASF Corporation
 BOLERO® - Reg. TM of Kumiai Chemical Industry Co. Ltd.
 COMMAND® 3 ME - Reg. TM of FMC Corporation
 Dimilin® - Reg. TM of Uniroyal Chemical
 FACET® - Reg. TM of BASF Corporation
 GRANDSTAND® - Trademark of Dow AgroSciences
 KARATE® - Reg. TM of Syngenta
 Permit® - Reg. TM of Monsanto
 PROWL® - Reg. TM of BASF Corporation
 REGIMENTVELOCITY® - Trademark of Valent U.S.A. Corporation
 WHIP® - Reg. TM of Aventis Crop Science

Manufactured for:
Valent U.S.A. Corporation
 P.O. Box 8025
 Walnut Creek, CA 94596-8025
www.valent.com

Made in U.S.A.

EPA Reg. No. 59639-105
EPA Est.

059639.000105.20021118.Reg03Velocitynotif.doc

15/15

1333 North California Blvd.
Suite 600
PO. Box 8025
Walnut Creek, CA 94596-8025
(925) 256-2700



November 18, 2002

REGIMENT™ Herbicide
EPA Reg. No. 59639-105
Notification for Alternate Brand Name
OPP Identifier No. 283410

Mr. Marion J. Johnson, Jr. – PM-10
Office of Pesticide Programs (7504C)
Document Processing Desk (NOTIF)
U.S. Environmental Protection Agency
401 M Street S.W.
Washington D.C. 20460-0001

Dear Mr. Johnson,

Valent U.S.A. Corporation would like to notify EPA of an Alternate Brand Name for our product Regiment Herbicide (59639-105).

Alternate Brand Name: Velocity® Herbicide

Enclosed is the EPA Form 8570-1 pesticide application, a certification statement and a label clearly marked in red.

If you have any questions or need further information, please contact me at (925)256-2791 or Eric Maurer in our Washington DC office at (202)872-4682.

Sincerely

Linda Obrestad
Labeling Specialist
Registration and Regulatory Affairs

Enclosures

