

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number
283396

Application for Pesticide - Section I

1. Company/Product Number 59639-115	2. EPA Product Manager Joseph M. Tavano	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) ESTEEM 35 WP Insect Growth Regulator	PM# 10	
5. Name and Address of Applicant (Include ZIP Code) Valent U.S.A. Corporation 1333 N. California Blvd., Suite 600 Walnut Creek, CA 94596-8025 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION OCT 18 2002
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
NOTIFICATION: Changed Net Contents from 5 oz. to 4 oz. Changed the number of water soluble packets to treat 5 acres to 4 acres. Corrected type in the Almond Crop section of "Tree Nuts".

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Labels: Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Linda Obrestad	Title Labeling Specialist	Telephone No. (Include Area Code) (925) 256-2791
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Linda Obrestad</i>	3. Title Labeling Specialist	
4. Typed Name Linda Obrestad	5. Date <i>October 9, 2002</i>	

1333 North California Blvd.
 Suite 600
 P.O. Box 8025
 Walnut Creek, CA 94596-8025
 (925) 256-2700



October 9, 2002

Notification
ESTEEM 35 WP Insect Growth Regulator
 EPA Reg. No. 59639-115
 OPP ID No. 283396

Mr. Joseph M. Tavano – PM 10
 Insecticide Branch Registration Division (7505C)
 Document Processing Desk (NOTIF)
 U.S. Environmental Protection Agency
 401 M Street S.W.
 Washington D.C. 20460-0001

Dear Mr. Tavano:

Valent USA is submitting this notification to update our product label Esteem 35 WP Insect Growth Regulator (EPA Reg. No. 59639-115) as follows:

- Change the Net Contents from 5 ounces to 4 ounces
- Changed the number of water soluble packets to treat from 5 acres to 4 acres in the Table of Equivalents.
- Corrected a typo in the Almond crop section of "Tree Nut s".

I have enclosed the above mentioned label (clearly marked in red), EPA Form 8570-1, and a certification statement.

If you have any questions or need further information, please contact me at (925)256-2791.

Sincerely,

A handwritten signature in black ink that reads "Linda Obrestad".

Linda Obrestad
 Labeling Specialist
 Registration and Regulatory Affairs

Enclosures

3/16



REDLINE
COPY

ESTEEM[®] 35 WP (INSECT GROWTH REGULATOR)

NOTIFICATION
OCT 18 2002


	By Wt.
Active Ingredient	
Pyriproxyfen (2-[1-methyl-2-(4-phenoxyphenoxy) ethoxy]pyridine)	35.0%
Other Ingredients	65.0%
Total	100.0%

ESTEEM 35 WP Insect Growth Regulator is a wettable powder formula containing 35% active ingredient.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS



NET CONTENTS 54 OUNCES

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Causes moderate eye irritation. Harmful if inhaled, swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact **1-800-892-0099** for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long sleeved shirt, long pants, and socks plus shoes.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by disposing of equipment washwaters or rinsate. Avoid direct application and/or spray drift to bee hives.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.


Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over long-sleeved shirt and long pants, waterproof gloves, and socks plus shoes.



**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. EXCEPT AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

In no event shall Valent or Seller be liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

If Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

WATER SOLUBLE PACKAGING

MULTIPLE PACKAGING

This bag contains multiple water soluble packets of ESTEEM 35 WP Insect Growth Regulator. Do not handle the packets with wet gloves or allow the packets to become wet prior to addition to spray tank. Do not break open packets. Refer to the Tables of Equivalents to calculate the number of packets to use. If all packets are not used, close and reseal outer container to protect remaining packet(s).

SINGLE PACKAGING

This bag contains one water soluble packet of ESTEEM 35 WP Insect Growth Regulator. Do not handle the packet with wet gloves or allow the packet to become wet prior to addition to spray tank. Do not break open packets. Refer to the Table of Equivalents to calculate the number of packets to use. Open outer bag by pulling perforated tearstrip and drop the unopened inner packet of ESTEEM 35 WP Insect Growth Regulator into spray tank.

TABLE OF EQUIVALENTS


RATES OF ESTEEM 35 WP PER ACRE	# OF WATER SOLUBLE PACKETS TO TREAT 5.04 ACRES
3 oz.	3
4 oz.	4
5 oz.	5

GENERAL INFORMATION

Do not apply this product through any type of irrigation system.

ESTEEM 35 WP is intended for use in Insect Pest Management (IPM) or Insect Resistance Management (IRM) programs. ESTEEM 35 WP will not control insect adults, and it is recommended to be used in combination and/or rotation with other IPM or IRM materials. Contact your local state extension service for details.

ESTEEM 35 WP acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM 35 WP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM 35 WP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.




APPLE

GENERAL: ESTEEM 35 WP Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM 35 WP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM 35 WP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Apple	Codling Moth Suppression of: Leafroller Pandemis Oblique Banded	4 to 5 oz.	<p>Apply by ground with airblast equipment in a minimum of 50 gals./A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>1st Generation Codling Moth: Apply at Biofix plus 100 degree days (this is usually close to petal fall). Make a second application 14 to 21 days later or timed to coincide with peak moth flights. For subsequent applications, if needed, apply other insecticides registered for Codling Moth control. The addition of oil has been shown to increase codling moth activity of ESTEEM 35 WP. Follow local recommendations regarding in season use and rates of oil in your area.</p> <p>Oblique Banded and Pandemis Leafroller: Apply to the overwintering larvae at pink to petal fall tree growth stage. ESTEEM 35 WP does not directly control the larvae but inhibits normal pupation and adult emergence.</p>
	Florida Red Scale Lecanium Scale San Jose Scale	4 to 5 oz.	<p>Apply by ground with airblast equipment in a minimum of 50 gals./A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply at delayed dormant through pink. For delayed dormant use a superior spray oil at the manufacturer's recommended rate of oil. Under heavy scale pressure, use the higher rate of ESTEEM 35 WP.</p> <p>Scale: For control in season make application in cover sprays when scale crawlers first emerge. The use of oil at the crawler emergence stage has been shown to improve control. Follow local recommendations regarding in season use and rates of oil in your area.</p>

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Apple (cont.)	Aphid Green Apple Rosy Apple Spirea Leafminer Spotted Tentiform Western Tentiform	3 to 5 oz.	Apply by ground with airblast equipment in a minimum of 50 gals./A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage. Leafminers: Apply at peak moth flight for control of 1 st or 2 nd generation. This usually occurs at pre-pink to pink apple stage for 1 st generation and 2 nd cover for 2 nd generation. Rosy Apple Aphid: Apply at 1/2" green apple growth stage. Green Apple & Spirea Aphid: Apply when aphids first appear, but before aphids are colonized. TIMING IS CRITICAL FOR ACCEPTABLE APHID CONTROL.
ESTEEM 35 WP Insect Growth Regulator may be mixed and/or alternated with commonly used insecticides to comply with local IPM programs and resistance management programs. •Do not apply within 45 days of harvest. •Do not exceed 2 ESTEEM 35 WP applications per growing season. •Do not exceed 10 oz. of ESTEEM 35 WP per acre per season. •Do not apply earlier than 14 days after last ESTEEM 35 WP treatment.			

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM 35 WP Insect Growth Regulator to two applications per growing season. ESTEEM 35 WP may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.




PEAR

GENERAL: ESTEEM 35 WP Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM 35 WP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM 35 WP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Pear	Pear Psylla	4 to 5 oz.	<p>Apply by ground with airblast equipment in 75 to 400 gals./A with a superior spray oil at manufacturer's recommended rates for the stage of growth. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply a single application during the delayed dormant to pink stage of growth at 5 oz./A or make two applications from delayed dormant through petal fall stage of growth at 4 to 5 oz./A. Make application just before sustained egg lay is anticipated.</p> <p>DO NOT USE SKIP ROWS DURING APPLICATION.</p> <p>For in-season control of pear psylla, use an insecticide recommended for this use.</p>
	Lecanium Scale San Jose Scale	4 to 5 oz.	<p>Apply by ground with airblast equipment in 100 to 400 gals./A at delayed dormant with a superior spray oil at the manufacturer's recommended rates. Under heavy scale pressure, use the higher rate.</p> <p>Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>For control of scale in season make application in cover sprays when scale crawlers first emerge. The use of oil at the crawler emergence stage has been shown to improve performance. Follow local recommendations regarding in season use and rates of oil in your area.</p>

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Pear (cont.)	Codling Moth Suppression of: Leafroller Pandemis Oblique Banded	4 to 5 oz.	Apply by ground with airblast equipment in 100 to 400 gals./A. 1st Generation Codling Moth: Apply at Biofix plus 100 degree days. Make a second application 14 to 21 days later or timed to coincide with peak moth flights. For subsequent applications, if needed, apply other insecticides registered for Codling Moth control. The addition of oil has been shown to increase codling moth activity of ESTEEM 35 WP. Follow local recommendations regarding in season use and rates of oil in your area. For additional generations apply other insecticides registered for Codling Moth Control.
<p>ESTEEM 35 WP Insect Growth Regulator may be mixed and/or alternated with commonly used insecticides to comply with local IPM programs and resistance management programs.</p> <ul style="list-style-type: none"> •Do not apply within 45 days of harvest. •Do not exceed 2 ESTEEM 35 WP applications per growing season. •Do not exceed 10 oz. of ESTEEM 35 WP Insect Growth Regulator per acre per season. •Do not apply earlier than 14 days after last ESTEEM 35 WP treatment. 			

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM 35 WP Insect Growth Regulator to two applications per growing season. ESTEEM 35 WP may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.



CITRUS

GENERAL: ESTEEM 35 WP Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM 35 WP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM 35 WP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Citrus	Scale Black Brown Soft California Red Florida Red Snow	5 oz.	<p>Apply with ground equipment in 200 to 1500 gals./A. Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply anytime prior to fruit becoming infested when scale is in the crawler stage. Thorough coverage is essential for good control.</p> <p>ESTEEM 35 WP may be applied with superior spray oils at 1/2 to 1%. Follow the manufacturer's directions and recommendations on timing and rates.</p> <p>NOTE: For scale apply only one application during emergence of scale crawlers per calendar year.</p>
	Citrus Black Fly Citrus Leafminer Citrus Whitefly	3 oz.	<p>Apply with ground application in 200 to 1500 gals./A. Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply when pests are just beginning to build and before infestation has occurred. Thorough coverage is essential for good control.</p> <p>ESTEEM 35 WP may be mixed and/or alternated with commonly used insecticides to comply with local IPM programs and resistance management programs.</p>
<ul style="list-style-type: none"> •Do not apply within 1 day of harvest. •Do not make more than 2 ESTEEM 35 WP applications per season. •Do not exceed 8 oz. of ESTEEM 35 WP per acre per season. •Do not apply earlier than 21 days after last ESTEEM 35 WP treatment. 			


MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM 35 WP Insect Growth Regulator to two applications per growing season. ESTEEM 35 WP may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

TREE NUTS

GENERAL: ESTEEM 35 WP acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM 35 WP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM 35 WP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Almond	Black Scale San Jose Scale	4 to 5 oz.	Apply with ground equipment in 100 to 400 gals./A. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will assure better coverage.
	Suppression of: Peach Twig Borer		Apply at either dormant, delayed dormant or an in season cover spray. At dormant, delayed dormant use a spray oil at the manufacturer's recommended oil rate. In a cover spray make applications when scale crawlers first emerge. Follow local recommendations regarding in season use and rates of oil in your area. ESTEEM 35 WP has provided good suppression of Peach Twig Borer when applied delayed dormant through bloom sprays.
Almonds: •Do not apply within 21 days of harvest. •Do not exceed 1 ESTEEM 35 WP application per growing season. •Do not exceed 5 oz. of ESTEEM 35 WP per acre per season.			

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM 35 WP Insect Growth Regulator to one application per growing season. ESTEEM 35 WP may be used in alternation with other IGR materials possessing dissimilar modes of action and/or other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.



CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
<p>Tree Nuts (cont.)</p> <p>Beechnut Brazil Butternut Cashew Chestnut Chinquapin Filbert (Hazelnut) Hickory Macadamia Pecan Walnut</p>	<p>Codling Moth Dusky Veined Aphid Walnut Aphid</p>	<p>4 to 5 oz.</p>	<p>Apply with ground equipment in 100 to 400 gals./A. Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>1st Generation Codling Moth: Apply at Biofix plus 100 degree days. Make a second application 14 to 21 days later or timed to coincide with peak moth flights. For subsequent applications, if needed, apply other insecticides registered for Codling Moth control. The addition of oil has been shown to increase codling moth activity of ESTEEM 35 WP. Follow local recommendations regarding in season use and rates of oil in your area.</p>
	<p>Black Scale San Jose Scale Walnut Scale</p> <p>Suppression of: Peach Twig Borer</p>	<p>4 to 5 oz.</p>	<p>Apply with ground equipment in 100 to 400 gals./A at at either dormant, delayed dormant or an in season cover spray. At dormant, delayed dormant use a spray oil at the manufacturer's recommended oil rate. In a cover spray make applications when scale crawlers first emerge. Follow local recommendations regarding in season use and rates of oil in your area.</p> <p>Under heavy scale pressure, use the higher rate. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will assure better coverage.</p>
<p>ESTEEM 35 WP Insect Growth Regulator may be mixed and/or alternated with commonly used insecticides to comply with local IPM programs and resistance management programs.</p> <ul style="list-style-type: none"> •Do not apply within 21 days of harvest. •Do not exceed 2 ESTEEM 35 WP applications per growing season. •Do not exceed 10 oz. of ESTEEM 35 WP per acre per season. •Do not apply ESTEEM 35 WP earlier than 14 days after the last ESTEEM 35 WP application. 			

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM 35 WP Insect Growth Regulator to two applications per growing season. ESTEEM 35 WP may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

STONE FRUIT

GENERAL: ESTEEM 35 WP Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. ESTEEM 35 WP has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since ESTEEM 35 WP is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Stone Fruit (Including, but not limited to): Apricot Cherry Nectarine Peach Plum Prune	Lecanium Scale San Jose Scale Suppression of: Peach Twig Borer Oriental Fruit Moth	4 to 5 oz.	Apply in 100 to 400 gals./A. with ground equipment. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will assure better coverage. Scale: Apply at dormant/delayed dormant. At dormant/delayed dormant, use a spray oil at the manufacturer's recommended oil rate. Under heavy scale pressure, use the higher rate of ESTEEM 35 WP. Peach Twig Borer: ESTEEM 35 WP has provided good suppression of Peach Twig Borer when applied delayed/dormant through bloom sprays. Use a spray oil at the manufacturer's recommended rate for the stage of crop growth.
In-season sprays of ESTEEM 35 WP may not protect fruit from blemishes by scale. ESTEEM 35 WP Insect Growth Regulator may be mixed and/or alternated with commonly used insecticides to comply with local IPM programs and Resistance Management Programs. •Do not apply within 14 days of harvest. •Do not exceed 3 ESTEEM 35 WP application per growing season. •Do not exceed 15 oz. of ESTEEM 35 WP per acre per season. •Do not apply earlier than 14 days after last ESTEEM 35 WP treatment.			

MANAGING INSECT RESISTANCE: Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent USA Corporation is restricting ESTEEM 35 WP Insect Growth Regulator to three applications per growing season. ESTEEM 35 WP may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

STORAGE AND DISPOSAL PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Store in a cool dry place.

Keep pesticide in original container.

Keep container closed when not in use.

Do not put concentrate or dilute into food or drink containers.

Not for use or storage in or around the home.

For help with any spill, leak, fire or exposure involving this material, call day or night 1-800-892-0099.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Do not reuse the outer bag. Dispose the outer bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Copyright© 2002 by Valent U.S.A. Corporation

ESTEEM® - Reg. TM of Valent U.S.A. Corporation

Manufactured for:

Valent U.S.A. Corporation

P.O. Box 8025

Walnut Creek, CA 94596-8025

www.valent.com

Made in U.S.A.

Form 000913 – ESTEM35R.PND

EPA Reg. No. 59639-115

EPA Est. No. _____