



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

May 12, 2026

Zenna Scarlett
Senior Manager, Regulatory
Valent U.S.A. LLC
4600 Norris Canyon Road
San Ramon, CA 94583

Subject: Label Amendment - Registration Review Mitigation for Etoxazole
Product Name: Zeal WDG Miticide
EPA Registration Number: 59639-123
Case Number: 474817
Application Dates: December 22, 2020

Dear Zenna Scarlett:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Etoxazole Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must

submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Rachel Fletcher by phone at 202-566-2354, or via email at fletcher.rachel@epa.gov.

Sincerely,

A handwritten signature in blue ink that reads "Cathryn Britton". The signature is written in a cursive style with a large initial 'C'.

Cathryn Britton
Chief, Risk Management and Implementation
Branch V
Pesticide Re-evaluation Division (7508M)
Office of Pesticide Programs

ENCLOSURE: Stamped label



ACCEPTED
 05/12/2026
 Under the Federal Insecticide, Fungicide
 and Rodenticide Act as amended, for the
 pesticide registered under
 EPA Reg. No. 59639-123

[Bracketed Information is Optional Text]

GROUP 10B ACARICIDE

Zeal[®] WDG Miticide

Active Ingredient	By Wt.
Etoxazole*	72%
Other Ingredients	28%
Total	100%

*2-(2,6-difluorophenyl)-4-[4-(1,1-dimethylethyl)-2-ethoxyphenyl]-4,5-dihydrooxazole

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 59639-123
EPA Est. No.

NET WEIGHT _____

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
<p>HOT LINE NUMBER</p> <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 800-892-0099 for emergency medical treatment information.</p>	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Causes moderate eye irritation. Avoid contact with eyes, skin and clothing. Avoid breathing dust.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Applicators and other handlers must wear: Long-sleeved shirt and long pants, shoes plus socks and chemical-resistant gloves made of: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride ≥14 mils, or viton ≥14 mils.

Mixer/loaders must also wear: a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N*, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to freshwater and marine/estuarine aquatic invertebrates, including oysters and shrimp. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: long sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material and shoes plus socks.

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product, which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label **and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

Valent must be provided with prompt notice of any claim so that an immediate inspection of the affected property or plants can be made. To the extent applicable law allows such requirement, notice of a claim shall be given promptly to Valent, whether the claim is based on contract, negligence, strict liability, or other tort or otherwise be barred from any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law.

Read and follow the entire label of each product to be used in the tank mix with this product.

CHEMIGATION

RESTRICTION: Do not apply this product through any type of irrigation or chemigation system.

MANDATORY SPRAY DRIFT MANAGEMENT

SPRAY DRIFT

Aerial Applications:

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select a nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572.1).
- If the windspeed is 10 miles per hour or less, applicators must use $\frac{1}{2}$ swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 miles per hour, applicators must use $\frac{3}{4}$ swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters
- Do not apply during temperature inversions.

Airblast Applications

- Sprays must be directed into the canopy.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at low ends and when spraying outer row.
- Do not apply during temperature inversions.

Ground Boom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to select a nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications

Applicators are required to select a nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572.1).

- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications

- Take precautions to minimize spray drift.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, TetraSan 5 WDG contains a Group 10B acaricide. Any mite population may contain individuals naturally resistant to TetraSan 5 WDG and other Group 10 B. The resistant individuals may dominate the mite population if this group of acaricide are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay acaricide resistance, take the following steps:

- Rotate the use of TetraSan 5 WDG or other Group 10B acaricide within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with acaricide from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of the mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual acaricide selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefits only for the period where both insecticides are active.
- Adopt an integrated pest management program for acaricide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance contact Valent U.S.A. LLC at 800-6-VALENT (682-5368).

CANEBERRY (Subgroup 13-07A)

Blackberry; Loganberry; Raspberry, Black; Raspberry, Red; Raspberry, Wild; Cultivars varieties and/or hybrids of these

PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
European Red Spider Mite McDaniel Spider Mite Two-spotted Spider Mite Yellow Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	0

*Pre-Harvest Interval

SPECIAL INSTRUCTIONS

Apply by ground as a full coverage spray in a minimum of 50 gals/A of water. Applications of *Zea/* WDG Miticide must be done using enough carrier (water) to ensure thorough coverage of the crop's vegetative and reproductive parts, which mites are using for dispersal, feeding and reproduction. Use higher water volumes on older trees and varieties that have more compact and dense foliage. Higher volumes of water coupled with well calibrated equipment will help to provide better coverage.

Best results are achieved when mite populations are low. *Zea/* WDG Miticide is predominately an ovicide/larvicide. Apply *Zea/* WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.

Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.

See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to *Zea/* WDG Miticide or other acaricides.

FIELD CORN, POPCORN, CORN (Grown For Seed Production)

PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
Banks Grass Mite ¹ Carmine Spider Mite Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite	1.0 to 3.0 (0.045 to 0.135 lb ai/A)	6.0 (0.27 lb ai/A)	2	21

*Pre-Harvest Interval

SPECIAL INSTRUCTIONS

Apply with [air or] ground equipment in adequate water for uniform coverage ([a minimum of 3 gals/A by air or] a minimum of 10 gals/A [by ground]). Coverage is essential for good control. Use of higher water volume will assure better coverage. Applications of *Zea/* WDG Miticide are recommended before tasseling stage to allow good coverage and provide better control.

For field corn utilized for seed production apply *Zea/* WDG Miticide before or at tasseling growth stage. If a second application is needed wait for 14 days to re-treat with *Zea/* WDG Miticide.

Best results are achieved when mite populations are below threshold or beginning to build up on the plants. *Zea/* WDG Miticide is predominately an ovicide/larvicide and should be used early in the life cycle of mites. Always follow the recommended threshold for your area.

Do not use rates below 1.0 oz/A (0.045 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.

Restriction: Do not apply treatments less than 14 days apart.

See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to *Zea/* WDG Miticide or other acaricides.

¹ Applications targeting exclusively Banks grass mites may require higher rates particularly if populations have exceeded the established threshold in the fields to be treated with *Zea/* WDG Miticide. Also, under extreme drought conditions and higher populations Banks grass mites might require more than one application of a miticide.

COTTON				
PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
Carmine Spider Mite Pacific Spider Mite Two-spotted Spider Mite	0.66 to 1.0 (0.03 to 0.045 lb ai/A)	1.0 (0.045 lb ai/A)	1	28
*Pre-Harvest Interval				
<p>SPECIAL INSTRUCTIONS</p> <p>Apply with air or ground equipment in adequate water for uniform coverage (3 to 10 gals/A by air or 10 to 50 gals/A by ground). Coverage is essential for good control. Use of higher water volume will assure better coverage.</p> <p>Best results are achieved when mite populations are low. <i>Zea/</i> WDG Miticide is predominately an ovicide/larvicide and should be used early in the life cycle of mites. Apply <i>Zea/</i> WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Do not use below use rate 0.66 oz/A (0.03 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p> <p>See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to <i>Zea/</i> WDG Miticide or other acaricides.</p>				

CUCURBIT VEGETABLES (Crop Group 9)

Acorn Squash; Balsam Apple; Balsam Pear; Bittermelon; Butternut Squash; Calabaza Squash; Cantaloupe; Chayote (fruit); Chinese Cucumber; Chinese Okra; Chinese Waxgourd (Chinese Preserving Melon); Citron Melon; Cucumber; Cucuzza; Gherkin; Gourd, Edible; Hechima; Hubbard Squash; Hyotan; *Momordica* spp.; Muskmelon; Pumpkin; Spaghetti Squash; Summer Squash; Watermelon; Winter Squash

PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
Two-spotted Spider Mite Carmine Spider Mite Pacific Spider Mite Strawberry Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	7

*Pre-Harvest Interval

SPECIAL INSTRUCTIONS

Apply with [air or] ground equipment in adequate water for uniform coverage ([a minimum of 3 gals/A by air or] a minimum of 10 gals/A [by ground]). Coverage is essential for good control. Use of higher water volume will assure better coverage.

Applications of *Zea/* WDG Miticide must be made using enough carrier (water) to ensure thorough coverage of the crop's vegetative and reproductive parts, which mites use for dispersal, feeding and reproduction. Use higher water volumes on more mature plants and varieties that have more compact and dense foliage. Higher volumes of water coupled with well calibrated equipment will help to provide better coverage.

Best results are achieved when mite populations are low. *Zea/* WDG Miticide is predominately an ovicide/larvicide. Apply *Zea/* WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.

Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.

See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to *Zea/* WDG Miticide or other acaricides.

HOPS				
PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
Two-spotted Spider Mite	3.0 to 4.0 (0.135 to 0.18 lb ai/A)	4.0 (0.18 lb ai/A)	1	7
*Pre-Harvest Interval				
<p>SPECIAL INSTRUCTIONS</p> <p>Apply with ground equipment in a minimum of 50 gals/A of water. Coverage is essential for good control. Use of higher water volume will assure better coverage.</p> <p>Best results are achieved when mite populations are low. <i>Zea/</i> WDG Miticide is predominately an ovicide/larvicide. Apply <i>Zea/</i> WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Do not use below use rate 3.0 oz/A (0.135 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p> <p>See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to <i>Zea/</i> WDG Miticide or other acaricides.</p>				

LOW GROWING BERRY (Subgroup 13-07G)

Bearberry; Bilberry; Blueberry, Lowbush; Cloudberry; Cranberry; Lingonberry; Muntries; Partridgeberry; Strawberry; Cultivars, varieties, and/or hybrids of these

PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
European Red Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	1

*Pre-Harvest Interval

SPECIAL INSTRUCTIONS

Apply with ground equipment in a minimum of 100 gals/A of water. Coverage is essential for good control. Use of higher water volume will assure better coverage.

Best results are achieved when mite populations are low. *Zea/* WDG Miticide is predominately an ovicide/larvicide and should be used early in the life cycle of mites. Apply *Zea/* WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.

Zea/ WDG Miticide will not control Cyclamen Mite. Another miticide registered for this pest should be used if these mites are a problem.

Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.

See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to *Zea/* WDG Miticide or other acaricides.

PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A) + 10 2/3 fl oz Danitol® 2.4 EC Insecticide (0.2 lb ai/A)	3.0 (0.135 lb ai/A)	1	1

*Pre-Harvest Interval

Continued

SPECIAL INSTRUCTIONS

Apply with ground equipment in adequate water for uniform coverage (using a minimum of 100 gals/A).

Alternate with other non-pyrethroid insecticides if retreatment is needed in less than 30 days to comply with local IPM programs.

Comply with all applicable directions, restrictions and precautions on the registered label for *Danitol* 2.4 EC Insecticide.

See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to *Zeal* WDG Miticide or other acaricides.

MINT (Peppermint and Spearmint)				
PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
Pacific Spider Mite Strawberry Spider Mite Two-spotted Spider Mite	2.0 to 4.0 (0.09 to 0.18 lb ai/A)	4.0 (0.18 lb ai/A)	1	7
*Pre-Harvest Interval				
<p>SPECIAL INSTRUCTIONS Apply with [air or] ground equipment in adequate water for uniform coverage (minimum of [10 gals/A by air or] 50 gals/A [by ground]). Coverage is essential for good control. Use of higher water volume will assure better coverage.</p> <p>Best results are achieved when mite populations are low. <i>Zea/ WDG</i> Miticide is predominately an ovicide/larvicide. Apply <i>Zea/ WDG</i> Miticide at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p> <p>See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to <i>Zea/ WDG</i> Miticide or other acaricides.</p>				

PEPPER AND EGGPLANT (Subgroup 8-10B)

African Eggplant; Bell Pepper; Eggplant; Martynia; Non-Bell Pepper; Okra; Pea Eggplant; Pepino; Roselle; Scarlet Eggplant; Cultivars, varieties and/or hybrids of these

PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	7

*Pre-Harvest Interval

SPECIAL INSTRUCTIONS

Apply with ground equipment in a minimum of 20 gals/A of water. Applications of *Zea/* WDG Miticide must be made using enough carrier (water) to ensure thorough coverage of the crop's vegetative and reproductive parts, which mites use for dispersal, feeding and reproduction. Use higher water volumes on more mature plants and varieties that have more compact and dense foliage. Higher volumes of water coupled with well calibrated equipment will help to provide better coverage.

Best results are achieved when mite populations are low. *Zea/* WDG Miticide is predominately an ovicide/larvicide. Apply *Zea/* WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.

Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.

See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to *Zea/* WDG Miticide or other acaricides.

POME FRUIT (Crop Group 11)				
Apple; Crabapple; Loquat; Mayhaw; Pear; Pear, Asian; Quince				
PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
European Red Mite McDaniel Spider Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	14
*Pre-Harvest Interval				
SPECIAL INSTRUCTIONS				
Apply by ground with airblast equipment in a minimum of 100 gals/A. Coverage is essential for good control. Use of higher water volume will assure better coverage.				
Best results are achieved when mite populations are low. <i>Zea/</i> WDG Miticide is predominately an ovicide/larvicide and should be used early in the life cycle of mites. Apply <i>Zea/</i> WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.				
<i>Zea/</i> WDG Miticide will not control Rust Mites or Blister Mites. If these pests are a problem, use an alternative miticide registered for that use.				
Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.				
See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to <i>Zea/</i> WDG Miticide or other acaricides.				

SMALL FRUIT VINE CLIMBING, EXCEPT FUZZY KIWIFRUIT (Subgroup 13-07 F)

Amur River Grape; Gooseberry; Grape; Kiwifruit, Hardy; Maypop; Schisandra Berry; Cultivars, varieties, and/or hybrids of these

PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
European Red Mite Pacific Spider Mite Two-spotted Spider Mite Willamette Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	14

*Pre-Harvest Interval

SPECIAL INSTRUCTIONS

Apply by ground as a full coverage spray in a minimum of 25 gals/A of water. Coverage is essential for good control. Use of higher water volume will assure better coverage.

Best results are achieved when mite populations are low. *Zea/* WDG Miticide is predominately an ovicide/larvicide and should be used early in the life cycle of mites. Apply *Zea/* WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.

Zea/ WDG Miticide will not control Rust Mites or Blister Mites. If these pests are a problem, use an alternative miticide registered for that use.

Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.

See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to *Zea/* WDG Miticide or other acaricides.

STONE FRUIT (Crop Group 12)

Apricot; Cherry (sweet and tart); Nectarine; Peach; Plum; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plumcot; Prune (fresh)

PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
European Red Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	7

*Pre-Harvest Interval

SPECIAL INSTRUCTIONS

Apply with ground equipment in a minimum of 50 gals/A of water. Coverage is essential for good control. Use of higher water volume will assure better coverage.

Best results are achieved when mite populations are low. *Zeal* WDG Miticide is predominately an ovicide/larvicide. Apply *Zeal* WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.

Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.

See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to *Zeal* WDG Miticide or other acaricides.

TREE NUTS

Almond; Beech Nut; Brazil Nut; Butternut; Cashew; Chestnut; Chinquapin; Filbert (Hazelnut); Hickory Nut; Macadamia Nut; Pecan; Pistachio; Walnut (Black and English)

PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
European Red Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	28

*Pre-Harvest Interval

SPECIAL INSTRUCTIONS

Apply by ground with airblast equipment as a full coverage spray.

Best results are achieved when mite populations are low. *Zea/ WDG* Miticide is predominately an ovicide/larvicide. Apply *Zea/ WDG* Miticide at or prior to threshold for your area but not greater than the maximum rate listed.

Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.

See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to *Zea/ WDG* Miticide or other acaricides.

NON-BEARING FRUIT TREES				
PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI*
European Red Mite McDaniel Spider Mite Pacific Spider Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	One Year
*Pre-Harvest Interval				
SPECIAL INSTRUCTIONS				
Apply by ground with airblast equipment as a full coverage spray.				
Best results are achieved when mite populations are low. <i>Zea/</i> WDG Miticide is predominately an ovicide/larvicide. Apply <i>Zea/</i> WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.				
Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.				
See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to <i>Zea/</i> WDG Miticide or other acaricides.				

TROPICAL AND SUB-TROPICAL FRUIT (INEDIBLE PEEL)

Avocado; Canistel; Mango; Papaya; Sapodilla; Sapote, Black; Sapote, Mamey; Star Apple

PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
Avocado Brown Mite Persea Mite Two-spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	1

*Pre-Harvest Interval

SPECIAL INSTRUCTIONS

Apply with [air or] ground equipment in adequate water for uniform coverage ([minimum 20 gals/A by air or] minimum 50 gals/A [by ground]). Applications of *Zeal* WDG Miticide must be made using enough carrier (water) to ensure thorough coverage of the crop's vegetative and reproductive parts, which mites use for dispersal, feeding and reproduction. Use higher water volumes on more mature plants and varieties that have more compact and dense foliage. Higher volumes of water coupled with well calibrated equipment will help to provide better coverage.

Best results are achieved when mite populations are low. *Zeal* WDG Miticide is predominately an ovicide/larvicide. Apply *Zeal* WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.

Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.

See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to *Zeal* WDG Miticide or other acaricides.

CHRISTMAS TREES				
PESTS	Product Rate oz/Acre	Maximum Rate per Season oz/Acre	Maximum Applications per Season	PHI* Days
McDaniel Spider Mite Pacific Spider Mite Spruce Spider Mite Two- spotted Spider Mite	2.0 to 3.0 (0.09 to 0.135 lb ai/A)	3.0 (0.135 lb ai/A)	1	N/A
<p>*Pre-Harvest Interval</p> <p>Apply by ground as full coverage spray or by air in a minimum of 10 gals/A. Coverage is critical for effective control.</p> <p>Best results are achieved when mite populations are low. <i>Zea/</i> WDG Miticide is predominately an ovicide/larvicide. Apply <i>Zea/</i> WDG Miticide at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Do not use below use rate 2.0 oz/A (0.09 lb ai/A) as this may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p> <p>See label section titled RESISTANCE MANAGEMENT RECOMMENDATIONS for recommendations to delay mite resistance to <i>Zea/</i> WDG Miticide or other acaricides.</p>				

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.
Open dumping is prohibited.

PESTICIDE STORAGE

Keep pesticide in original container.

Do not put concentrate or dilute into food or drink containers.

Store in cool, dry place.

Do not store diluted spray.

For help with any spill, leak, fire or exposure involving this material, call day or night
(800) 892-0099.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.

Copyright © 2026 by Valent U.S.A. LLC

Zeal is a registered trademark of Valent U.S.A. LLC

Danitol is a registered trademark of Sumitomo Chemical Company, Ltd.

Danitol is a restricted use product.

Manufactured for:

Valent U.S.A. LLC

P.O. Box 5075

San Ramon CA 94583

(800)-6-VALENT (682-5368)

Made in U.S.A.

EPA Reg. No. 59639-123

EPA Est. No.

059639-00123.20260429.ZWDG.Etoxazole.ID.Clean