



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

59639-136

Date of Issuance:

6-23-05

Term of Issuance:

Conditional

Name of Pesticide Product:

Velocity SG Herbicide

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Valent USA Corp.
1600 Riviera Avenue
P.O. Box 8025
Walnut Creek, CA
94596-8025

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided you agree in writing to:

1. Submit the following outstanding product chemistry data requirement: one year storage stability and corrosion characteristics study, within one year from the date of this letter.

Signature of Approving Official:


Date: 6-23-05

James Tompkins, Product Manager (25)
Herbicide Branch, Registration Division (7505C)

2/18

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). A stamped copy of labeling is enclosed for your records. If you have any questions please contact Erik Kraft at 703-308-9358.

Sincerely,



James Tompkins
Product Manager (25)
Herbicide Branch
Registration Division (7505C)

3/18



ACCEPTED
with COMMENTS
in EPA Letter Dated

6-23-05

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

GROUP 2 HERBICIDE

59639-136

Velocity[®] SG Herbicide

SELECTIVE POSTEMERGENCE HERBICIDE FOR CONTROL OF ANNUAL BLUEGRASS, ROUGH STALK BLUEGRASS AND CERTAIN BROADLEAF WEEDS ON GOLF COURSE TURFGRASS AND SOD FARMS

SUPPRESSION OF DOLLAR SPOT

Active Ingredient	By Wt.
*Bispyribac-sodium.....	17.6%
Other Ingredients.....	<u>82.4%</u>
Total	100.0%

*Sodium 2,6-bis[(4,6-dimethoxypyrimidin-2-yl)oxy]benzoate

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS _____

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Avoid breathing dust or spray mist.

FIRST AID

- If on skin or clothing:**
 - Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- If in eyes:**
 - Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- If swallowed:**
 - Call a poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- If inhaled:**
 - Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-892-0099** for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow instructions for category A on an EPA chemical resistant category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material (such as butyl or nitrile), shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

5/18

ENVIRONMENTAL HAZARDS

This product is toxic to non-target plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters or rinsate.

This product may contaminate water through drift of spray in wind. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 24 hours.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, greenhouses or sodfarms.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift.
Do not enter treated areas without protective clothing until sprays have dried.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material (such as butyl or nitrile), shoes plus socks.

6/18

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. EXCEPT AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

In no event shall Valent or Seller be liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

If Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

7/18

TABLE OF CONTENTS

GENERAL INFORMATION

- Use Precautions.....
- Application.....
- Mixing and Spraying Equipment Preparation and Cleanup.....
- Mixing Instructions.....
- Resistance Management
- To Delay Herbicide Resistance.....
- Weeds Controlled by VELOCITY SG..... Table 1

DIRECTIONS FOR USE IN CREEPING BENTGRASS AND PERMANENTLY ESTABLISHED PERENNIAL RYEGRASS

- Crops and Use Site.....
- Reseeding, Overseeding or Sprigging.....
- VELOCITY SG Use in Creeping Bentgrass and Permanently Established Perennial Ryegrass.....
- Recommended Control Programs
- Slow Conversion to Creeping Bentgrass or Perennial Ryegrass
- Rapid Conversion to Creeping Bentgrass or Perennial Ryegrass.....
- Restrictions and Limitations
- Spray Drift
- Dollar Spot Suppression by VELOCITY SG in Creeping Bentgrass..... Table 2

DIRECTIONS FOR USE IN BERMUDAGRASS OVERSEEDED WITH PERENNIAL RYEGRASS

- Crop and Use Site.....
- VELOCITY SG Use in Bermudagrass Overseeded with Perennial Ryegrass.....
- Recommended Control Programs
- Control of Annual Bluegrass and Broadleaf Weeds.....
- Seed Head Suppression of Annual Bluegrass.....
- Restrictions and Limitations
- Spray Drift

STORAGE AND DISPOSAL.....

GENERAL INFORMATION

VELOCITY SG is a postemergence herbicide that will selectively control annual bluegrass (*Poa annua*), roughstalk bluegrass (*Poa trivialis*) and numerous broadleaf weeds that are growing within certain species of established turfgrass. VELOCITY SG will also suppress seedhead production by annual bluegrass. VELOCITY SG displays activity against emerged weeds, but has almost no preemergent activity. Therefore, VELOCITY SG will not control weeds that germinate after application.

VELOCITY SG inhibits the enzyme acetolactate synthase (ALS), which plants require to produce three key amino acids. Annual bluegrass and other susceptible weeds usually stop growing within 3 to 7 days after treatment, and turn yellow or brown within 3 to 14 days after treatment. Plant death typically occurs by 21 to 28 days after treatment. More than one application of VELOCITY SG is usually required for maximum weed control.

VELOCITY SG is absorbed by plant foliage and roots. Plant uptake and performance of VELOCITY SG is influenced by environmental conditions, cultural practices and spray coverage. For best results, only apply VELOCITY SG when turf and weeds are actively growing. Application of VELOCITY SG to control weeds will also suppress infection of creeping bentgrass by dollar spot, *Sclerotinia homeocarpa*. Suppression of dollar spot will be greatest when a weed control program is initiated in the late spring or early summer before the appearance of significant dollar spot infection. When VELOCITY SG is applied at this time, dollar spot suppression is usually evident for several weeks after the last application of Velocity SG. Therefore, early season application of VELOCITY SG may delay the initiation of a dollar spot control program with fungicides, and reduce overall fungicide application on creeping bentgrass.

VELOCITY SG will also provide some curative control of dollar spot, but should not be used in place of labeled fungicides to control established infections of this disease.

USE PRECAUTIONS

VELOCITY SG is a very active herbicide, and users should exercise good judgment and caution until familiarity is gained with this product. Due to variability of turfgrass varieties, growth stages, environmental conditions, cultural practices and application techniques, users should test this product under user growing conditions in a small area, and evaluate treated turf for 28 days to determine if the herbicide can be used safely in a widespread application.

APPLICATION

Apply VELOCITY SG using standard, low pressure (20 to 50 psi) spray equipment in a sufficient volume of water to provide thorough spray coverage and a uniform spray pattern. To ensure thorough coverage, apply a minimum of 20 gals. of spray solution per acre. Apply VELOCITY SG with ground equipment using standard commercial sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume. Avoid streaking, skips or excessive overlaps during application. Do not apply with flood jet nozzles, air induction nozzles or hand held sprayers, as equipment may not provide adequate or uniform coverage. Calibrate spray equipment before each use and check periodically during application. The addition of a spray indicator, such as dyes or foams, is recommended.

9/16

MIXING AND SPRAYING EQUIPMENT PREPARATION AND CLEANUP

Use well maintained and clean equipment to apply VELOCITY SG. Clean the spray tank, and all hoses and booms according to the manufacturer's directions for the last product used before applying VELOCITY SG. This will ensure that no residue from the previous application remains in the sprayer.

Trace amounts of VELOCITY SG in or on mixing or spraying equipment may have an adverse effect on subsequently sprayed plants. Therefore, it is important that the sprayer be properly cleaned after spraying VELOCITY SG. Thoroughly drain, clean and rinse all mixing and spraying equipment including tanks, booms, hoses, strainers, screens and nozzles immediately after use. Use the following procedure:

1. Remove all physical residues.
2. Thoroughly drain and rinse tanks, booms and hoses with clean water.
3. Fill the tank one half full with clean water and use a spraying or mixing tank cleaner that **does not** contain chlorine. Fill the remainder of the tank with clean water. Let agitate or recirculate according to the directions of the cleaner manufacturer. Thoroughly flush the boom and hoses before draining.
4. Rinse all hoses, tanks, nozzles, strainers and booms with clean water to remove the tank cleaner. Follow the directions provided by the tank cleaner manufacturer.
5. Fill the tank half full with clean water and add 1 gal. of 3% active household ammonia for every 100 gals. of water the tank will hold. Fill the remainder of the tank with clean water and allow the solution to agitate or recirculate for 15 minutes. Thoroughly flush the ammonia cleaning solution through the boom, hoses, nozzles, screens and strainers before draining the tank.
6. Remove the strainers, nozzles and screens and clean separately in a solution of household ammonia and water.
7. Replace the strainers, nozzles and screens.
8. Repeat Step 5.
9. Thoroughly rinse the tank with clean water and flush the water through the boom, nozzles and hoses in order to remove all traces of ammonia.
10. Dispose of the rinsate on site or at an approved waste disposal facility.

IMPORTANT: Do not use chlorine bleach with ammonia, or toxic chlorine gas may be released. Remove all traces of liquid fertilizer containing any form of ammonia or ammonium before adding any chlorine source such as chlorine bleach.

MIXING INSTRUCTIONS

VELOCITY SG should completely dissolve in the spray tank within approximately five minutes. Dissolution rate may be slowed by cold water, lack of agitation or water containing high concentrations of boron or sulfur.

1. Fill clean spray tank 1/3 to 1/2 of desired level with clean water.
2. While agitating, add VELOCITY SG and ensure it has dissolved completely before proceeding. Agitation should create a rippling or rolling action on the water surface.
3. Fill spray tank to desired level with water. Agitation should continue until spray solution has been applied.
4. Mix only the amount of spray solution that can be applied the day of mixing.
5. Apply VELOCITY SG within 6 hours after mixing with water in spray tank.

19/18

RESISTANCE MANAGEMENT

VELOCITY SG Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to VELOCITY SG and other Group 2 herbicides. Weed species with acquired resistance to Group 2 may eventually dominate the weed population if Group 2 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by VELOCITY SG or other Group 2 herbicides.

TO DELAY HERBICIDE RESISTANCE

- Avoiding the consecutive use of VELOCITY SG or other target site of action Group 2 herbicides that have similar target site of action, on the same weed species.
- Using tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on a comprehensive Integrated Pest Management (IPM) program.
- Monitoring treated weed populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.
- For further information contact Valent U.S.A. Corporation at the following toll free number: 1-800-682-5368.

11/18

Table 1. WEEDS CONTROLLED BY VELOCITY SG

COMMON NAME	SCIENTIFIC NAME	WEED SIZE	APPLICATION RATE oz./A
Bluegrass			6 to 9
Annual	<i>Poa annua</i>	Up to seedhead	
Roughstalk	<i>Poa trivialis</i>	Up to seedhead	
Chickweed			
Common	<i>Stellaria media</i>	Up to seedhead	
Sticky	<i>Cerastium glomeratum</i>	Up to seedhead	
Clover			
Large Hop	<i>Trifolium campestre</i>	Up to seedhead	
White	<i>Trifolium repens</i>	Up to seedhead	
Dandelion	<i>Taraxacum officinale</i>	Seedling stage	
Henbit	<i>Lamium amplexicaule</i>	Up to seedhead	
Lawn Burweed	<i>Soliva pterosperma</i>	Up to seedhead	
Parsley-Piert	<i>Alchemilla arvensis</i>	Up to seedhead	
Plantain			
Broadleaf	<i>Plantago major</i>	Up to seedhead	
Buckhorn	<i>Plantago lanceolata</i>	Up to seedhead	
Swinecress	<i>Coronopus didymus</i>	Up to seedhead	
Yellow Nutsedge	<i>Cyperus esculentus</i>	Seedling stage	
Yellow Woodsorrel	<i>Oxalis stricta</i>	Seedling stage	

DIRECTIONS FOR USE IN CREEPING BENTGRASS AND PERMANENTLY ESTABLISHED PERENNIAL RYEGRASS

CROPS AND USE SITE

- Creeping Bentgrass, *Agrostis palustris*
- Perennial Ryegrass, *Lolium perenne*
- Golf course fairways and tees mowed at 3/8" to 3/4" in height; sod farms mowed at 1/2" to 3/4" in height.

VELOCITY SG may injure creeping bentgrass or perennial ryegrass that is not well established or that has been weakened by moisture stress, pests, diseases, chemicals, low fertility, thatch, mechanical injury or other stresses. VELOCITY SG may also cause unacceptable injury to creeping bentgrass and perennial ryegrass mowed at greens height.

VELOCITY SG may initiate mild chlorosis when applied to creeping bentgrass or perennial ryegrass. The onset, intensity and persistence of symptoms are at least partially influenced by environmental conditions (i.e., temperature, cloud cover and rainfall) and cultural practices. Under cool and cloudy conditions, symptoms tend to appear more slowly than under warm sunny conditions. Symptoms may also be more persistent under cool cloudy conditions because turfgrass is growing less vigorously. Therefore, avoid application of VELOCITY SG in the spring before grass resumes active growth, or in the fall after growth slows.

VELOCITY SG can injure creeping bentgrass and perennial ryegrass that are under high heat stress. Therefore, do not apply VELOCITY SG to turfgrass that is exhibiting symptoms of heat stress, or if significant heat stress (temperatures above 90°F) is expected during the week following application. It is recommended that VELOCITY SG not be applied after June 30 in areas where creeping bentgrass or perennial ryegrass are typically exposed to prolonged periods of significant heat stress during the summer months.

In general, VELOCITY SG should only be applied during the time of year when creeping bentgrass, perennial ryegrass and target weeds are actively growing. The use season for VELOCITY SG will therefore vary according to location. VELOCITY SG will perform optimally (i.e. best weed control and least chlorosis to creeping bentgrass and perennial ryegrass) under sunny conditions when daytime high temperatures are between 65°F and 80°F during and after application.

VELOCITY SG may cause significant injury to other turf species, especially certain cultivars of Kentucky bluegrass, *Poa pratensis*. VELOCITY SG should not be applied in heavy traffic and/or heavily shaded turf areas, which are more prone to herbicide injury.

NOTE: Annual or roughstalk bluegrass chlorosis can be mistaken for creeping bentgrass or perennials ryegrass chlorosis, especially in sites with a moderate to heavy infestation. Turf chlorosis is usually more apparent when small patches of turf within a fairway or tees are treated with VELOCITY SG, and less apparent when entire fairways or tees are treated. Therefore, if temporary turf chlorosis is a concern, avoid treating small patches of turf that are surrounded by nontreated areas of turf.

RESEEDING, OVERSEEDING OR SPRIGGING

In turfgrass with >10% annual bluegrass, overseeding is recommended in conjunction with VELOCITY SG applications to promote conversion to creeping bentgrass or perennial ryegrass, and to avoid stand thinning due to loss of annual and roughstalk bluegrass.

VELOCITY SG may be applied to sodded or sprigged creeping bentgrass and to perennial ryegrass that is well established. Creeping bentgrass must have a developed root system and uniform stand and have received at least two mowings before the first application of Velocity SG.

13/18

Following an application of VELOCITY SG, wait 10 days before reseeding or sprigging. When reseeding or sprigging, always use proper cultural practices such as soil cultivation, irrigation and fertilization to ensure rapid turf establishment. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed-to-soil contact. Do not apply VELOCITY SG between 10 days before and 60 days after overseeding with creeping bentgrass or perennial ryegrass.

VELOCITY SG USE IN CREEPING BENTGRASS AND PERMANENTLY ESTABLISHED PERENNIAL RYEGRASS

VELOCITY SG should only be applied during the time of year when creeping bentgrass, perennial ryegrass and target weeds are actively growing. The use season for VELOCITY SG will therefore vary according to location.

Recommended Use Season:

- Northern States - May 15 to September 1
- Southern Temperate States - April 15 to June 30

Contact your Valent representative or your local extension specialist for a recommendation specific to your area.

Recommended Control Programs

The control program for VELOCITY SG should be determined by considering the desired level and speed of control, and the severity of infestation by annual and/or roughstalk bluegrass. Maximum control will usually require more than one application.

- **Slow Conversion to Creeping Bentgrass or Perennial Ryegrass**
Apply VELOCITY SG on a 7 day interval at the rate of 2.0 oz./A (10 g ai/A). Begin application early in the recommended use season, and continue until the desired level of control is achieved. This program should be considered for turf with a heavy infestation of annual and/or roughstalk bluegrass, where complete removal of these weeds during a single season could result in an unacceptable stand of creeping bentgrass or perennial ryegrass.
- **Rapid Conversion to Creeping Bentgrass or Perennial Ryegrass**
Apply VELOCITY SG up to four times at the rate of 6.0 oz./A (30 g ai/A) on a 14 to 21 day interval. Use 21 day interval if creeping bentgrass or perennial ryegrass is exhibiting undesirable chlorosis at 14 days after application. Efficacy may be decreased if application interval exceeds 21 days. This program should be considered for turf with light infestations of annual and/or roughstalk bluegrass, where removal of these weeds would not result in an unacceptable stand of turfgrass.

RESTRICTIONS AND LIMITATIONS

- Do not apply through any type of irrigation system.
- Do not apply to golf greens or roughs.
- Do not apply if rain is expected within 12 hrs. after application.
- Do not mow or irrigate turfgrass within 12 hrs. after application.
- Do not apply to moist or wet turfgrass (including dew).
- Do not mix with other pesticides, fertilizers, wetting agents, spreader stickers, surfactants or other adjuvants.
- Do not apply more than 9 oz. of VELOCITY SG per acre per application.
- Do not apply more than a total of 12 oz. of VELOCITY SG per acre in a 28 day period.
- Do not apply with flood jet nozzles, air induction nozzles or hand sprayers.
- Do not apply when temperatures are below 55°F or above 85°F.
- Do not apply to turfgrass under stress due to drought, temperature, disease, low fertility, heavy thatch, mechanical injury or other stresses.
- Do not apply VELOCITY SG between 10 days before and 60 days after overseeding with creeping bentgrass or perennial ryegrass.
- VELOCITY SG has not been evaluated for safety on all creeping bentgrass and perennial ryegrass cultivars.

14/18

- VELOCITY SG has not been evaluated under all microclimates or against all biotypes of annual and roughstalk bluegrass. Therefore, performance may be less effective in some locations, and against some biotypes of these weed species.
- VELOCITY SG should not be applied in heavy traffic and/or heavily shaded turf areas.
- VELOCITY SG may dramatically reduce overall turfgrass cover due to its high activity against annual and roughstalk bluegrass.
- Turf growth regulators may affect the efficacy and safety of VELOCITY SG.

SPRAY DRIFT

- Do not spray if winds are gusty or if wind speeds are greater than 5 mph.
- Do not apply within 15 ft. of native plant communities when sustained winds will carry VELOCITY SG towards these native plant communities.

Table 2. DOLLAR SPOT SUPPRESSION BY VELOCITY SG IN CREEPING BENTGRASS

DISEASES		APPLICATION RATE oz./A	SPECIAL INSTRUCTIONS
COMMON NAME	SCIENTIFIC NAME		
Dollar Spot	<i>Sclerotinia homeocarpa</i>	6 to 9	When used for weed control, VELOCITY SG can substantially suppress the development and severity of dollar spot in bentgrass fairways and tees. Suppression may be evident for several weeks after the final application. To maximize suppression of dollar spot, initiate weed control program in the late spring or early summer before or soon after the appearance of dollar spot symptoms in bentgrass.

USE PRECAUTIONS:

- See "RESTRICTIONS AND LIMITATIONS".
- VELOCITY SG can suppress dollar spot when applied before or soon after appearance of symptoms, but may not provide adequate curative control of established infections of dollar spot.
- Do not use VELOCITY SG in place of labeled fungicides for curative control of dollar spot.
- Do not use VELOCITY SG to suppress dollar spot on golf greens.

15/18

DIRECTIONS FOR USE IN BERMUDAGRASS OVERSEEDED WITH PERENNIAL RYEGRASS

CROP AND USE SITE

- Hybrid or Common Bermudagrass turf, *Cynodon dactylon*, that is fall overseeded with perennial ryegrass, *Lolium perenne*.
- Golf course fairways and tees mowed at 3/8" to 3/4" in height; sod farms mowed at 1/2" to 3/4".

VELOCITY SG may injure perennial ryegrass that is not well established or that has been weakened by moisture stress, pests, diseases, chemicals, low fertility, thatch, mechanical injury or other stresses. Do not apply VELOCITY SG until at least 60 days after perennial ryegrass seedling emergence. VELOCITY SG may also cause unacceptable injury to perennial ryegrass mowed at greens or rough height.

VELOCITY SG may initiate mild chlorosis when applied to perennial ryegrass. The onset, intensity and persistence of symptoms are at least partially influenced by environmental conditions (i.e., temperature, cloud cover and rainfall) and cultural practices. Under cool and cloudy conditions, symptoms tend to appear more slowly than under warm sunny conditions, but symptoms may also be more persistent under cool cloudy conditions because turfgrass is growing less vigorously.

NOTE: Annual bluegrass chlorosis can be mistaken for ryegrass chlorosis, especially under higher infestation levels of annual bluegrass, and when VELOCITY SG application is not initiated until mid-late flower. Turf chlorosis is usually more apparent when small patches of turf within a fairway or tee are treated with VELOCITY SG, and less apparent when entire fairways or tees are treated. Therefore, if temporary turf chlorosis is a concern, avoid treating small patches of ryegrass that are surrounded by larger areas of nontreated turf.

Application of a complete foliar fertilizer 3 to 4 days after application of VELOCITY SG may decrease the amount of ryegrass chlorosis. Some forms of iron can antagonize the performance of VELOCITY SG. Therefore, do not tank mix VELOCITY SG with foliar fertilizers that contain iron. Tank mixing VELOCITY SG with surfactants or other adjuvants may increase ryegrass chlorosis to unacceptable levels, and should therefore be avoided.

VELOCITY SG should not thin ryegrass when applied in accordance with the label, but can cause thinning if applied at excessive rates, especially when ryegrass is under heat or moisture stress, and mowed at less than 1/2". Therefore, do not exceed labeled rates, do not apply to ryegrass that is exhibiting symptoms of heat or moisture stress, and do not apply when air temperatures are greater than 80°F or are predicted to exceed 80°F in the three days after application.

To maximize performance, initiate treatment with VELOCITY SG when annual bluegrass first begins flowering in the mid-late winter and temperatures are high enough to promote active growth. Do not apply when air temperatures are below 55°F or are NOT predicted to exceed 55°F in any of the three days following application. In general, VELOCITY SG will perform optimally (i.e. best weed control and least chlorosis to ryegrass) under sunny conditions when daytime high temperatures are consistently between 60°F and 75°F during and after application.

VELOCITY SG may be less efficacious against annual bluegrass growing in thin stands of ryegrass. In thin ryegrass stands, annual bluegrass is exposed to less competition from ryegrass and therefore, more annual bluegrass will germinate, and the resulting plants will grow more vigorously and be more difficult to control than in denser stands of ryegrass. To maximize the efficacy of VELOCITY SG, broadcast ryegrass seed at a minimum of 300 lbs. per acre, and employ cultural practices that encourage the rapid formation of a dense stand of ryegrass.

16/18

VELOCITY SG can be applied in locations where bermudagrass does not go completely dormant and retains some green color during the winter. VELOCITY SG will not delay spring green-up, if applied before bermudagrass begins active growth (i.e. obvious tillering) in the late winter and spring. VELOCITY SG may temporarily discolor and regulate the growth of bermudagrass if applied after bermudagrass begins active growth, especially when applied at above the maximum-labeled rate.

Annual bluegrass density and vigor are higher in non-overseeded bermudagrass than in bermudagrass overseeded with perennial ryegrass. As a result, VELOCITY SG may not be effective against annual bluegrass growing in non-overseeded bermudagrass, especially when applied during the late winter and spring. In addition, if VELOCITY SG is applied after non-overseeded bermudagrass has resumed actively growing, any resulting discoloration or growth regulation will be more evident than in an overseeded site, where it would be masked by ryegrass. Therefore, application of VELOCITY SG is not recommended in non-overseeded bermudagrass.

VELOCITY SG may cause significant injury to other desirable turf species, especially certain cultivars of Kentucky bluegrass, *Poa pratensis*. VELOCITY SG should not be applied in heavy traffic and/or heavily shaded turf areas, which are more prone to herbicide injury.

VELOCITY SG USE IN BERMUDAGRASS OVERSEEDED WITH PERENNIAL RYEGRASS

VELOCITY SG should be applied in the winter or early spring when annual bluegrass first begins to actively flower in the late winter or early spring and temperatures are within the recommended range, but not until at least 60 days after ryegrass emergence. The use season will therefore vary according to location and environmental conditions.

Recommended Use Season: January 1 to April 15

Contact your Valent representative or your local extension specialist for a recommendation specific to your area.

RECOMMENDED CONTROL PROGRAMS:

The control program for VELOCITY SG should be determined by considering the desired level of control, and the tolerance for ryegrass chlorosis. Effective control or seed head suppression may require more than one application.

- **Control of Annual Bluegrass and Broadleaf weeds**

Apply VELOCITY SG two or three times on a 14 to 21 day interval at 6.0 oz./A (30 g ai/A). Use 21 day interval if perennial ryegrass is exhibiting undesirable chlorosis at 14 days after application. Do not make more than two applications per year where mowing height is less than 1/2". Efficacy may be decreased if application interval exceeds 21 days. This program should be considered for heavier infestations of annual bluegrass, and where there is a higher tolerance for temporary ryegrass chlorosis.

- **Seed head Suppression of Annual Bluegrass**

Apply VELOCITY SG three times on a 14 day interval at 3.0 oz./A (15 g ai/A). Efficacy may be decreased at longer applications intervals. While this program will strongly suppress production of seed heads, it may kill a small percentage of annual bluegrass and other labeled weeds. This program will cause less chlorosis than the maximum control program, and should be considered where there is a low tolerance for ryegrass chlorosis.

17/18

RESTRICTIONS AND LIMITATIONS

- Do not apply through any type of irrigation system.
- Do not apply to golf greens or roughs.
- Do not apply to ryegrass mowed at less than 3/8".
- Do not apply if rain is expected within 12 hrs. after application.
- Do not mow or irrigate turfgrass within 12 hrs. after application.
- Do not apply to moist or wet turfgrass (including dew).
- Do not mix with other pesticides, fertilizers, wetting agents, spreader stickers, surfactants or other adjuvants.
- Do not apply more than 9 oz. of VELOCITY SG per acre per application.
- Do not apply more than a total of 12 oz. of VELOCITY SG per acre in a 28 day period.
- Do not exceed three applications per acre per year.
- Do not apply with flood jet nozzles, air induction nozzles or hand sprayers.
- Do not apply when temperatures are below 55°F or above 85°F.
- Do not apply to turfgrass under stress due to drought, temperature, disease, low fertility, heavy thatch, mechanical injury or other stresses.
- Do not apply in spring after bermudagrass has begun actively tillering.
- Do not apply to perennial ryegrass until at least 60 days after seedling emergence.
- VELOCITY SG has not been evaluated for safety on all perennial ryegrass cultivars.
- VELOCITY SG has not been evaluated under all microclimates or against all biotypes of annual and roughstalk bluegrass. Therefore, performance may be less effective in some locations, and against some biotypes of these weed species.
- VELOCITY SG should not be applied in heavy traffic and/or heavily shaded turf areas.
- Turf growth regulators may affect the efficacy and safety of Velocity SG.

SPRAY DRIFT

- Do not spray if winds are gusty or if wind speeds are greater than 5 mph.
- Do not apply within 15 ft. of native plant communities when sustained winds will carry VELOCITY SG towards these native plant communities.

18/18

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Store in a cool dry place.
Keep pesticide in original container.
Keep container closed when not in use.
Do not put concentrate or dilute into food or drink containers.
Not for use or storage in or around the home.
For help with any spill, leak, fire or exposure involving this material, call day or night **1-800-892-0099**.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Copyright© 2004 by Valent U.S.A. Corporation

Velocity® - Reg.TM of Valent U.S.A. Corporation

Manufactured for:
Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025
www.valent.com

Made in U.S.A.

EPA Reg. No. 59639-XXX
EPA Est.