

 <p align="center">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7504P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p>	EPA Reg. Number: 59639-140	Date of Issuance: JAN 28 2008
	Term of Issuance: Conditional	
	Name of Pesticide Product: V-10161 4 SC	
NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)		
Name and Address of Registrant (include ZIP Code): Valent USA Corporation 1600 Riviera Avenue, Suite 200 P.O. Box 8025 Walnut Creek, CA 94596-8025 Attention: David Wustner		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(C) provided that you:</p> <p>A. Submit and/or cite all data required for registration of your product under FIFRA Section 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for re-registration of your product under FIFRA Section 4.</p> <p>B. Submit the following conditional data/information listed below, before January 31, 2009:</p> <p>1. For the phenyl-¹⁴C-labeled cow metabolism study (MRID 46708514), the petitioner provide complete sample history information for samples from the study, including not only dates of collection, but also dates of storage, radioassay, extraction, and analysis. If sample analyses were not completed within 6 months of sample collection, the data demonstrating that the metabolic profile was stable in the affected matrices during storage must be submitted.</p> <p align="right">(CONTINUED)</p>		
Signature of Approving Official: Tony Kish Product Manager, Team 22 Fungicide Branch Registration Division (7505P)	Date: JAN 28 2008	

2. For the pyridinyl-¹⁴C-labeled cow metabolism study (MRID 46708518):

A. Storage stability data are required to support the study. If samples were stored for greater than 6 months, provide data showing stability of the metabolic profile of the affected matrices for the duration of the storage period and under the conditions that the samples were stored.

B. Clarify that the identification of two peaks in liver methanol/water extract retention times are correct, and explain how the retention times for these metabolites were correlated with the identified metabolites in urine and kidney.

C. Correct the flowchart for fat to include the correct total radioactive residue (TRR) value for this matrix (0.039 ppm).

D. Recalculate the radioactivity levels and/or clarify the results for the HPLC analysis of any extract in which the calculated limit of quantitation (LOQ) was too high to allow meaningful interpretation of the chromatogram.

3. For the hen metabolism study reported in MRID 46708515, provide complete sample history information for samples from the study, including not only dates of collection but also dates of storage, radioassay, extraction, and analysis. If sample analyses were not completed within 6 months of sample collection, provide data demonstrating that the metabolic profile was stable in the affected matrices during the storage period and under the conditions that the samples were stored. Also submit copies of the LC/MS chromatograms of metabolite AE C653711 (BAM) in liver as well as the corresponding chromatogram of the reference standard. These chromatograms were referenced in the submission (MRID 46708515, page 113) but were not included.

4. The hen metabolism study conducted with [2,6-¹⁴C-pyridinyl]fluopicolide (MRID 46708519) is incomplete but upgradeable. The samples from this study were not extracted until >6 months after sample collection. Provide data showing stability of the metabolic profile for the duration of the storage period and under the conditions that the samples were stored.

5. For storage stability data for crops, submit additional storage stability data for celery and spinach reflecting a storage interval of 38 months. One study should be conducted on any representative leafy vegetable.

6. Submit one copy of a final printed label within 45 days from the date of this Notice which makes the following changes:

A. On page 2, change "Harmful if swallowed or absorbed through skin", to "Harmful if swallowed, inhaled, or absorbed through skin", and at the end of these statements add "Avoid breathing vapor or spray mist", which matches the acute toxicity review.

B. On Page 2, change the dollar signs (\$) and large letter X's (X) used for bullets to another symbol(s).

(CONTINUED)

C. On Page 6 under "Mode of Action", correct or clarify the disharmony between "The biochemical mode of action of fluopicolide is specific", vs just below this section "The target site of action....mode of action is not known". Deletion of the first sentence with "specific" or changing "mode of action to "site" in this sentence may harmonize these contradictions.

D. On page 6 under "General Restrictions" there are no listed tank mix partners for ornamentals as there is on the label for V-10161 VPP Fungicide. We recommend this be harmonized.

E. On page 9, there are no directions regarding using certain chemigation application methods for turf (eg. center pivot) versus using other methods for ornamentals (eg drip) as there is on the label for V-10161 VPP Fungicide. We recommend this be harmonized.

F. On page 10, in the rotational restriction box, change "Tuber vegetables" to "Vegetable, tuberous and corm, except potato, subgroup 1D".

G. On Page 11, either add something to the effect, "Chemigation is not recommended when only downy mildew control is desired", as it is on V-10162 Premix, or explain why it is not needed.

H. On page 14, delete "interiorscapes" as that indoor use was not assessed.

I. In the ornamental section, add the following restriction or equivalent, "The reentry interval for hand harvesting, thinning, pruning, and pinching of ornamental flowers is 6 days after the last product application".

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA Section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted with comments" is enclosed for your records.



GROUP 43 FUNGICIDE

V-10161 4 SC

(Fungicide)

FOR CONTROL AND/OR SUPPRESSION OF CERTAIN DISEASES IN CUCURBIT VEGETABLES, FRUITING VEGETABLES, GRAPES, LEAFY VEGETABLES AND SWEET POTATOES, AND FOR CONTROL OF PYTHIUM AND PHYTOPHTHORA ON TURFGRASS & ORNAMENTALS.

Active Ingredient	By Wt.
*Fluopicolide	39.5%
Other Ingredients	60.5%
Total	100.0%

*2,6-dichloro-N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl]benzamide

V-10161 4 SC is an aqueous flowable fungicide containing 4 lbs active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NET CONTENT _____

SHAKE WELL BEFORE USING
AGITASE BIEN ANTE USO

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 28 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

59639-

140

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION *inhalable*

Harmful if swallowed, or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Wear long-sleeved shirt and long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse. *Avoid BREATHING VAPOR OR SPRAY MIST.*

FIRST AID

- | | | |
|--------------------------------|----|--|
| If on skin or Clothing: | \$ | Take off contaminated clothing. |
| | \$ | Rinse skin immediately with plenty of water for 15-20 minutes. |
| | \$ | Call a poison control center or doctor for further treatment advice |
| If inhaled: | \$ | Move person to fresh air. |
| | \$ | If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. |
| | \$ | Call a poison control center or doctor for further treatment advice. |
| If swallowed: | \$ | Call a poison control center or doctor immediately for treatment advice. |
| | \$ | Have person sip a glass of water if able to swallow. |
| | \$ | Do not induce vomiting unless told to do so by the poison control center or doctor. |
| | \$ | Do not give anything by mouth to an unconscious person. |
| If in eyes: | \$ | Hold eye open and rinse slowly and gently with water for 15-20 minutes. |
| | \$ | Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. |
| | \$ | Call a poison control center or doctor for treatment advice. |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-800-892-0099** for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as natural rubber \geq 14 mils, socks and shoes.

USER SAFETY RECOMMENDATIONS

Users should:

- X Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- X Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- X Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water by disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, chemical resistant gloves made of any waterproof material such as natural rubber \geq 14 mils, socks and shoes.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, greenhouses or sodfarms.

Keep all unprotected persons out of operating areas or vicinity where there may be drift.
Do not enter treated areas without protective clothing until sprays have dried.

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the extent consistent with applicable law Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements, Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is later, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law, if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, consistent with applicable law. Read and follow the entire label of each product to be used in the tank mix with this product.

TABLE OF CONTENTS

GENERAL INFORMATION

General Restrictions and Limitations

Mode of Action

Resistance Management.....

Rainfastness.....

Jar Test to Determine Compatibility of Adjuvants and V-10161 4 SC.....

APPLICATION INSTRUCTIONS – General Ground & Air, Air and Chemigation:

Sprayer Preparation

Mixing Instructions.....

Sprayer Cleanup

Application Equipment.....

Carrier Volume

Aerial Application.....

Chemigation

ROTATIONAL RESTRICTIONS

CROP SPECIFIC RECOMMENDATIONS

TURFGRASS

ORNAMENTAL PLANTS

STORAGE AND DISPOSAL

GENERAL INFORMATION

V-10161 4 SC fungicide is formulated as 4 lbs ai/gal suspendable concentrate (SC). The active ingredient in V-10161 4 SC is fluopicolide. V-10161 4 SC exhibits protective, curative, eradicated and antispore activity. Fluopicolide is locally systemic and translaminar and also moves systemically via xylem tissue.

Optimal disease control is achieved when V-10161 4 SC is applied in a regularly scheduled spray program used in combination and/or rotation with other effective fungicides that have different modes of action (i.e., non group 43 fungicides).

GENERAL RESTRICTIONS AND LIMITATIONS

- A tank mix partner must be used with V-10161 4 SC for resistance management.
- Do not apply to greenhouse food crops.
- Do not apply this product when weather conditions favor spray drift from treated areas.
- When applying by air, observe drift management restrictions and precautions listed under "AERIAL APPLICATION". Best control is achieved through ground application.
- Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making decisions. Where states have more stringent regulations, they should be observed.

MODE OF ACTION

V-10161 4 SC is active against selective oomycetes. It exhibits the typical mode of action of specific systemic fungicides against oomycete fungi affecting several steps in the reproduction cycle.

The biochemical mode of action of fluopicolide is specific. Biochemical studies have shown that fluopicolide has an effect on spectrin-like proteins, believed to play a role in maintaining the membrane stability in ascomycete fungi or oomycetes, especially during hyphal tip extension. Microscopy studies demonstrate that fluopicolide induces a quick redistribution of these proteins from the membrane to the cytoplasm in both hyphae and zoospores.

RESISTANCE MANAGEMENT

V-10161 4 SC contains fluopicolide, a group 43 Fungicide. The target site of action of group 43 fungicides is specific but the detailed, biochemical mode of action is not known. Fungal isolates with acquired resistance to fluopicolide may eventually dominate the fungal population if fluopicolide is used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by fluopicolide. To maintain the performance of V-10161 4 SC in the field, do not exceed the total number of sequential applications of V-10161 4 SC and the total number of applications of V-10161 4 SC per season stated in the "CROP SPECIFIC REQUIREMENTS". Adhere to the label instructions regarding the consecutive use of V-10161 4 SC. The following recommendations may be considered to delay the development of fungicide resistance:

- **Tank Mixtures:** V-10161 4 SC must be used in a tank mixture with a fungicide from a different target site of action group that are registered for the same use and that is effective against the pathogens of concern. Valent recommends using at least the minimum labeled rates of each fungicide in the tank mix.

- **Integrated Pest Management (IPM):** V-10161 4 SC must be integrated into a multi-faceted disease and pest management program. Cultural practices known to reduce disease development should be followed. Consult your local extension specialist, certified crop advisor and/or Valent representative for additional IPM strategies established for your area. V-10161 4 SC may be used in Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.
- **Monitoring:** Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development.
- **Reporting:** If a Group 43 target site fungicide appears to be less effective or no longer effective against a pathogen that it previously controlled or suppressed, contact a Valent representative, local extension specialist or certified crop advisor to assist in determining the cause of reduced performance.

RAINFASTNESS

Conventional foliar spray applications of V-10161 4 SC are mainly for downy mildew and other foliar disease control and are rainfast 2 hours after application. Applications for foliar disease control must not be made if measurable rain is expected within 2 hours of application or foliar disease control may be reduced.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND V-10161 4 SC

A jar test should be performed before mixing commercial quantities of V-10161 4 SC, when using V-10161 4 SC for the first time, or when a new water source is being used.

1. Add 1 pt of water plus adjuvant to a quart jar. The water should be from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add 2.5 ml (1/2 tsp) of V-10161 4 SC to the quart jar, gently mix until product goes into suspension.
3. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
4. An ideal tank mix combination will be uniform and free of suspended particles. If any of the following conditions are observed the choice of adjuvant should be questioned:
 - a) Layer of oil or globules on the mixture's surface.
 - b) Flocculation: fine particles in suspension or as a layer on the bottom of the jar.
 - c) Clabbering: thickening texture (coagulated) like gelatin.

APPLICATION INSTRUCTIONS – General Ground & Air, Air and Chemigation:

SPRAYER PREPARATION

Before applying V-10161 4 SC, start with clean, well maintained application equipment. The spray tank, hoses and booms must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply V-10161 4 SC. If two or more products were tank mixed prior to V-10161 4 SC application, the most restrictive cleanup procedure should be followed.

MIXING INSTRUCTIONS

1. Fill clean spray or nurse tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, slowly add the V-10161 4 SC to the tank. Agitation should create a rippling or rolling action on the water surface.
3. When tank mixing V-10161 4 SC with other labeled pesticides, add water soluble packets first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions.
4. Add any required adjuvants.
5. Fill tank to desired level with water. Agitation should continue until all spray solution has been used or applied.

SPRAYER CLEANUP

Spray equipment must be cleaned following application of V-10161 4 SC. After V-10161 4 SC is applied, the following steps must be used to clean the spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
3. Drain tank completely.
4. Remove all nozzles and screens and rinse them in clean water.
5. The rinsate solution may be applied to the crops recommended on this label. Do not exceed the maximum labeled use rate. If cleaners are used, consult the cleaner label for disposal instructions. If no instructions are given, dispose of rinsate at an approved waste disposal facility.

APPLICATION EQUIPMENT

Application equipment should be clean and in good repair. Nozzles should be frequently checked for accuracy.

CARRIER VOLUME

Apply V-10161 4 SC in sufficient water to ensure thorough coverage of foliage, bloom and fruit. Thorough coverage is required for optimal disease control. For ground application, apply a minimum of 20 gallons of spray mixture per acre and for aerial application, do not apply less than 5 gallons of spray mixture per acre to assure uniform coverage. Follow individual "CROP SPECIFIC REQUIREMENTS" for appropriate spray volumes.

AERIAL APPLICATION

To minimize drift, apply the largest droplet size consistent with uniform coverage and satisfactory disease control. To obtain satisfactory application and avoid drift, the following directions must be observed:

Do not apply during low-level inversion conditions, when winds are gusty or under other conditions that favor drift. Do not spray when wind velocity is less than 2 mph or more than 10 mph.

- **Carrier Volume and Spray Pressure:** Application at less than 5 gals. per acre may provide inadequate coverage and control. The higher gallonage applications generally afford more consistent disease control. Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Nozzle Selection and Orientation:** Formation of very small drops may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray pressure. Use nozzles that produce flat fan or cone spray patterns. Use non-drip type nozzles, such as diaphragm type nozzles, to avoid unwanted discharge of spray solution. The nozzles must be directed toward the rear of the aircraft, producing a spray discharge at an angle between 0 and 15° downward. Do not place nozzles on the outer 25% of the wings or rotors.
- **Adjuvants and Drift Control Additives:** Refer to tank mix partner's label for adjuvant recommendation. Drift control additives may be used although it is better to obtain coarser sprays through appropriate nozzle selection and use wherever possible. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label, and be certain of compatibility with the tank mix and nozzle types being used before selecting any adjuvant types.

CHEMIGATION

V-10161 4 SC may be applied through sprinkler irrigation systems mainly for soil born infections. Follow all label requirements for these crops regarding application rates, timing of application, special instructions and precautions.

For chemigation applications apply this product only through center pivot, solid set, hand move irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of efficacy or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

The system must be properly calibrated (with water only) to ensure that the amount of V-10161 4 SC applied corresponds to the required rate on this label for the crop being chemigated.

Apply V-10161 4 SC in 1/2 to 3/4 inches of water during the first sprinkler set. Allow time for all lines to flush the fungicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining fungicide, a dye indicator may be injected into the lines to mark the end of the application period.

If you have any questions about calibration, you should contact your State Extension Specialist, equipment manufacturers or other experts.

Special Precautions for Chemigation

1. Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
2. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
3. The system must be free of leaks and clogged nozzles.
4. The pesticide must be supplied continuously for the duration of the aqueous application. An uneven application may cause injury to the crop or poor control.
5. Agitation must be maintained in the nurse tank.
6. The sprinkler chemigation system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
7. The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in the case where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with the pesticides being used and capable of being fitted with a system interlock.
12. Do not apply when wind speed favors drift beyond the area intended for treatment.

Chemigation Systems Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption, if such a system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to the public water system must contain a functional, reduced pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. All chemigation systems connected to the public water system must also follow restrictions listed in the preceding section titled "Special Precautions for Chemigation".

ROTATIONAL RESTRICTIONS

The following rotational crops may be planted after applying V-10161 4 SC at the labeled rate. Planting earlier than the recommended rotational interval is not allowed.

CROPS	ROTATIONAL INTERVAL
Cucurbit Vegetables Fruiting Vegetables Leafy Vegetables Tuber Vegetables	Immediately
All Other Crops	18 months

SPECIFIC REQUIREMENTS – All Food Crops

When to Apply: Begin application when crop and/or environmental conditions favor disease development. Applications must be made on a 7 to 14 day interval, depending on disease pressure. Under severe disease pressure use the higher labeled rate and shortest interval. Apply as a foliar spray in sufficient water to obtain thorough coverage.

Resistance Management: V-10161 4 SC must be used as part of an integrated pest management (IPM) program. Apply as a foliar spray in a tank mix with another labeled fungicide product with a different mode of action on the target pathogen in sufficient water to obtain thorough coverage.

Restrictions and Limitations

- The maximum V-10161 4 SC application rate is 4 fl oz per acre per application and 12 fl oz per acre per season.
- Apply no more than 2 sequential applications of V-10161 4 SC before alternating with an effective fungicide from a different resistance management group.
- Do not make more than 4 applications of V-10161 4 SC per acre per season.

Crops	Diseases	Application Rates		Minimum Time from Last Application to Harvest (PHI)	Specific Use Instructions
		fl oz/A	GPA		
Cucurbit Vegetables: Acorn Squash; Balsam Apple; Balsam Pear; Bittermelon; Butternut Squash; Calabaza; Cantaloupe; Chayote, Fruit; Chinese Cucumber; Chinese Okra; Chinese Preserving Melon; Chinese Waxgourd; Citron Melon; Cucumber; Cucuzza; Gherkin; Gourd, Edible; Hechima; Hubbard Squash; Hyotan; <i>Momordica</i> spp; Muskmelon; Pumpkin; Spaghetti Squash; Summer Squash; Watermelon; Winter Squash	Downy Mildew (<i>Pseudoperonospora cubensis</i>) Phytophthora (<i>Phytophthora cactorum</i>) Phytophthora Blight (<i>Phytophthora capsici</i>)	3 to 4 (0.09 to 0.125 lb ai/A)	Ground: 20 to 50 Aerial Minimum: 5	2 days	A tank mix including a labeled rate of another labeled fungicide product with a different mode of action on the target pathogen must be used with V-10161 4 SC for resistance management. Recommended spray interval: 10 days
	Late Blight (<i>Phytophthora infestans</i>) Phytophthora (<i>Phytophthora parasitica</i>) Phytophthora Blight (<i>Phytophthora capsici</i>)				

continued

SPECIFIC REQUIREMENTS – All Food Crops (continued)

Crops	Diseases	Application Rates		Minimum Time from Last Application to Harvest (PHI)	Specific Use Instructions
		fl oz/A	GPA		
Grapes	Downy Mildew (<i>Plasmopara viticola</i>)			21 days	A tank mix including a labeled rate of another labeled fungicide product with a different mode of action on the target pathogen must be used with V-10161 4 SC for resistance management. Recommended spray interval: 12 to 14 days
Leafy Vegetables (except Brassica Vegetables): Amaranth, Chinese Spinach; Arugula, Roquette; Cardoon; Celery; Celtuce; Chinese Celery; Chervil; Chrysanthemum, Edible-leaved; Chrysanthemum, Garland; Corn Salad; Cress, Garden; Cress, Upland; Dandelion; Dock, Sorrel; Endive, Escarole; Fennel; Florence; Lettuce, Head and Leaf; Orach; Parsley; Purslane, Garden; Purslane, Winter; Radicchio, Red Chicory; Rhubarb; Spinach; Spinach, New Zealand; Spinach, Vine; Swiss Chard	Downy Mildew (<i>Bremia lactucae</i>)	3 to 4 (0.09 to 0.125 lb ai/A)	Ground: 20 to 50 Aerial Minimum: 5	2 days	A tank mix including a labeled rate of another labeled fungicide product with a different mode of action on the target pathogen must be used with V-10161 4 SC for resistance management. Recommended spray interval: 10 days
Sweet Potato	Late Blight (<i>Phytophthora infestans</i>) Pink Rot (<i>Phytophthora erythroseptica</i>)			7 days	A tank mix including a labeled rate of another labeled fungicide product with a different mode of action on the target pathogen must be used with V-10161 4 SC for resistance management. Recommended spray interval: 10 days

TURFGRASS

V-10161 4 SC is effective at preventing and controlling damping off, root rot and stem diseases caused by pythium blight infesting turfgrass. Pythium blight is sometimes referred to as greasy spot or cottony blight and is in a group of fungi known as water molds. This disease is common in turfgrass areas with poor soil drainage and a wet turfgrass canopy. High humidity and high temperatures provide an ideal environment for development of pythium blight. Pythium blight occurs in all cool season grasses, but is especially prevalent in creeping bentgrass, annual bluegrass and perennial ryegrass.

DIRECTIONS FOR USE ON TURFGRASS

Turfgrass	Diseases	Application Rate	Special Instructions
<p>Established and Overseeded Turfgrass: Golf Course Turf Residential Turf Commercial Turf Sod Farms</p>	<p>Pythium Blight Pythium Damping Off</p>	<p>0.2 fl oz per 1000 sq ft (8.6 fl oz/A)</p>	<p>Do not apply more than a total of 0.4 fl oz per 1000 sq ft per year (0.54 lb ai/A per year).</p> <p>Apply in a minimum volume of 2 gals per 1000 sq ft</p> <p>A tank mix including a labeled rate of another labeled fungicide product with a different mode of action on the target pathogen must be used with V-10161 4 SC for resistance management.</p> <p>Overseeded Turf: Apply after seed germination to prevent pythium damping off.</p> <p>Established Turf: Apply when conditions favor disease development. Favorable conditions included high temperatures and high humidity.</p> <p>If necessary, make an additional application on a 14 day interval.</p>

ORNAMENTAL PLANTS

V-10161 4 SC is effective at preventing water mold, root rot, stem rot, crown rot and damping off caused by pythium and phytophthora diseases in ornamental plants. V-10161 4 SC is mixed with water and applied as a soil drench at the time of seeding or transplanting. V-10161 4 SC may be used on container, bench, or bed grown ornamentals in greenhouses or outdoor nurseries, interiorscapes, conifers including Christmas trees and for use on ornamentals grown for outdoor landscaping.

IMPORTANT: The large number of existing ornamental species and their varieties and cultivars coupled with the constant introduction of new varieties makes it impossible to field test V-10161 4 SC in every locale where it is sold or in all of the combinations of location and plant varieties. Further differences include the soil or media type; pH; moisture or fertility; environmental conditions such as temperature, lighting or degree days; horticultural practice; the manner of use and application of this product. To ensure that V-10161 4 SC is compatible with the ornamental plant variety or cultivar under your specific conditions, test this product on a limited scale and observe for phytotoxicity or other unintended effects for two weeks before making large scale applications.

DIRECTIONS FOR USE ON ORNAMENTAL PLANTS

Ornamentals	Diseases	Product Rates	Special Instructions														
Bedding Plants Conifers Flowering Plants Foliage Plants Ground Covers Non-Bearing Fruit Nut Trees Ornamentals Ornamental Trees Shrubs Vines	Phytophthora Pythium Blight	4 to 8 fl oz per 100 gals	<p>Application should be made prior to disease development. Use the higher rate with high potential for disease development. Make applications as necessary, but no more than once every 14 days.</p> <p>A tank mix must be used with V-10161 4 SC for resistance management.</p> <p>Do not apply more than two (2) applications of V-10161 4 SC per cropping cycle.</p> <p>Sequential Application: Do not apply sequentially, alternate with another fungicide registered for use on ornamental plants.</p> <p>Foliar Application: For foliar sprays, 100 gallons of spray mix will treat approximately 20,000 sq ft of area. Apply the spray mixture to all plant surfaces and to the point of runoff.</p> <p>Drench Application: Use enough solution to wet the root zones of plants. Use the information below as a guide for drench application.</p> <p>Container grown plants: Apply in volume as listed. For pot sizes other than those listed, adjust volume accordingly.</p> <table border="1" data-bbox="927 1325 1450 1556"> <thead> <tr> <th>Pot diameter</th> <th>fl oz per pot</th> </tr> </thead> <tbody> <tr><td>4</td><td>3</td></tr> <tr><td>5</td><td>4</td></tr> <tr><td>6</td><td>6</td></tr> <tr><td>8</td><td>10</td></tr> <tr><td>10</td><td>20</td></tr> <tr><td>12</td><td>30</td></tr> </tbody> </table> <p>Plants grown in flats or beds: Apply at a volume of 1 to 2 pints per square foot (5 to 10 qts per 10 sq ft).</p>	Pot diameter	fl oz per pot	4	3	5	4	6	6	8	10	10	20	12	30
Pot diameter	fl oz per pot																
4	3																
5	4																
6	6																
8	10																
10	20																
12	30																
<p>NOTE: Since ornamental varieties are numerous, constantly changing and may react differently to V-10161 4 SC and tank mixtures including V-10161 4 SC, test the product(s) on a small scale before making large scale applications.</p>																	

RATE CONVERSION CHART

Rate of V-10161 4 SC per 100 gals (fl oz)	Amount of V-10161 4 SC to add to water to achieve desired volume (fl oz)									
	1 gallon		5 gallons		10 gallons		25 gallons		50 gallons	
	fl oz	ml	fl oz	ml	fl oz	ml	fl oz	ml	fl oz	ml
4	0.04	1.2	0.2	6	0.4	12	1	30	2	60
6	0.06	1.8	0.3	9	0.6	18	1.5	45	3	90
8	0.08	2.4	0.4	12	0.8	24	2	60	4	120

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Keep pesticide in original container.
 Do not put concentrate or dilute into food or drink containers.
 Store in a cool dry place.
 Do not contaminate water, food or feed by storage and disposal.
 Do not store or transport near feed or food.
 Do not store at temperature below 32°F. If the product is exposed to temperatures below 32°F, thaw at 50°F or higher and shake gently to unify the product.
 For help with any spill, leak, fire or exposure involving this material, call day or night 800-892-0099.

PESTICIDE DISPOSAL

Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Do not reuse container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Copyright© 2007 by Valent U.S.A. Corporation

Manufactured for:
Valent U.S.A. Corporation
 P.O. Box 8025
 Walnut Creek, CA 94596-8025
 www.valent.com

Made in U.S.A.

EPA Reg. No. 59639-140
 EPA Est. No. _____

059639-00140.200712-28.V101614SC.doc.