

59639-144

10/18/2007

1/18



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

59639-144

Date of Issuance:

OCT 18 2007

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

V-10166 VPP Fungicide

Name and Address of Registrant (include ZIP Code):

Valent Corporation
1600 Riviera Ave., Suite 200
Walnut Creek, CA 94596-8025

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(C) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following change to the label:
 - a. Change the product registration number to "EPA Reg. No. 59639-144".

Page 1 of 2

Signature of Approving Official:

Mary L. Waller

Mary L. Waller, Product Manager (21)
Fungicide Branch, Registration Division (7505P)

Date:

10/18/2007

2/18

b. Revise the Non-Agricultural Use Requirements box by deleting the first sentence and revising the second sentence to read as follows: "Do not enter or allow others to enter until sprays have dried."

c. Revise the last sentence on page 8 to read as follows: "Where states have more stringent regulations, they must be observed."

d. Revise the fourth bullet on page 13 to read as follows: "All applications must be made according to the following use directions. Failure to follow. . .".

e. Revise the tank mix statement on page 5 to read as follows: "Tank mixing or use of this product shall be the exclusive risk of user, applicator and/or application advisor to the extent allowed by applicable law."


3. Submit the following studies within 18 months of the date of this product registration:

- a. Storage Stability – OPPTS 830.6317
- b. Corrosion Characteristics – OPPTS 830.6320

4. Submit one copy of the revised final printed label for the record before the product is released for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.



Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure: Label stamped "Accepted with Comments"

3/18



**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

10/18/2007

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

59639-144

GROUP 3 FUNGICIDE

V-10116 VPP Fungicide

Turfgrass Fungicide

A SYSTEMIC FUNGICIDE FOR CONTROL OF CERTAIN TURFGRASS DISEASES

Ornamental Fungicide

A SYSTEMIC FUNGICIDE FOR CONTROL OF CERTAIN ORNAMENTAL DISEASES

Active Ingredient	By Wt.
*Metconazole.....	50%
Other Ingredients.....	50%
Total	100%

*5-[(4-chlorophenyl)methyl]-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol


V-10161 VPP Fungicide is a water dispersible granule containing 50% active ingredient.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NET CONTENT _____



4/18

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID

- If swallowed:**
- Call a poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- If in eyes:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long-sleeved shirt and long pants, socks and shoes.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

7/18

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label; under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements, Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law, if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

8/18

GENERAL INFORMATION

V-10116 VPP Fungicide is formulated as 50% water dispersible granular (WDG). The active ingredient in *V-10116 VPP* is metconazole, a broad-spectrum triazole fungicide that works by inhibiting demethylation and other processes in sterol biosynthesis. *V-10116 VPP* is systemic and is quickly absorbed into plant tissue. *V-10116 VPP* displays preventative and curative properties.

V-10116 VPP may be used for disease control in turfgrass on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, athletic fields and sod farms.

V-10116 VPP may be used for disease control on ornamental plants in commercial indoor and outdoor nurseries, greenhouses and on ornamental plants found in landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Optimal disease control is achieved when *V-10116 VPP* is applied in a regularly scheduled spray program used in combination and/or rotation with other effective fungicides that have different modes of action (i.e., non Group 3 fungicides).

MODE OF ACTION

The active ingredient in *V-10116 VPP*, metconazole, belongs to the sterol biosynthesis inhibitor group of fungicides as classified by the U.S. EPA and Canada PMRA as a target site of action Group 3 fungicide.

RESISTANCE MANAGEMENT

V-10116 VPP contains metconazole a Group 3 fungicide (sterol biosynthesis inhibitors). Metconazole is effective against pathogens resistant to fungicides with modes of action different from those of target site Group 3 Fungicides (e.g., dicarboximides, strobilurins, benzimidazoles or phenylamides). Fungal isolates with acquired resistance to metconazole may eventually dominate the fungal population if metconazole is used repeatedly at the same site or in successive years as the primary method of control for the targeted pathogen species. Selection for resistance may be particularly rapid if resistance to Group 3 fungicides is already present in the pathogen population. This may result in reduced disease control by *V-10116 VPP* or other Group 3 fungicides. The following recommendations may be considered to delay the development of fungicide resistance:

1. **Tank mixtures:** If *V-10116 VPP* is used in tank mixtures with fungicides from different target site of action Groups that are registered and/or permitted for the same use and that are effective against the pathogens of concern, Valent recommends using at least the minimum labeled rates of each fungicide in the tank mix.

9/18

2. **Integrated Pest Management (IPM):** *V-10116 VPP* should be integrated into an overall disease and pest management program. Cultural practices known to reduce disease development should be followed. Consult your local extension specialist, certified advisor and/or Valent representative for additional IPM strategies established for your area. *V-10116 VPP* may be used in advisory (disease forecasting) programs, which recommend application timing based on environmental factors favorable for disease development.
3. **Monitoring:** Monitor efficacy of all fungicides used in the disease management program against the targeted pathogen and record other factors that may influence fungicide performance and/or disease development.
4. **Reporting:** If a Group 3 target site fungicide appears to be less or no longer effective against a pathogen that it previously controlled or suppressed, contact a Valent representative, local extension specialist or certified advisor to assist in determining the cause of reduced performance.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND *V-10116 VPP*

A jar test should be performed before mixing commercial quantities of *V-10116 VPP*, when using *V-10116 VPP* for the first time, when using new adjuvants, when using new tank mixes or when a new water source is being used.

1. Add 1 pt. of the water to a quart jar. The water should be from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add about 2 g of *V-10116 VPP* to the quart jar; gently mix until product goes into suspension.
3. Add appropriate amount of each tank mix adjuvant or partner.
4. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
5. An ideal tank mix combination will be uniform and free of large suspended particles. If any of the following conditions are observed the choice of adjuvant should be questioned:
 - a) Layer of oil or globules on the mixture's surface.
 - b) Flocculation: fine particles in suspension or as a layer on the bottom of the jar.
 - c) Clabbering: Thickening texture (coagulated) like gelatin.

TANK MIX PARTNERS

V-10116 VPP is compatible with most herbicides, insecticides, fungicides and other types of tank mix components. If tank mixes are used, follow all applicable restrictions and precautions on all labels. When evaluating potential tank mix partners, always apply to a small area of target plants and then monitor for adverse affects for at least two weeks before making large-scale applications.

10/18

SPRAYER PREPARATION

Before applying *V-10116 VPP*, start with clean, well maintained application equipment. The spray tank hoses and booms must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply *V-10116 VPP*. Nozzles should be frequently checked for accuracy. If two or more products were tank mixed prior to *V-10116 VPP* application, the most restrictive cleanup procedure should be followed.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, slowly add the *V-10116 VPP* to the spray tank. Agitation should create a rippling or rolling action on the water surface.
3. If tank mixing *V-10116 VPP* with other labeled pesticides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions.
4. Add any required adjuvants.
5. Fill spray tank to desired level with water. Agitation should continue until all spray solution has been applied.
6. Mix only the amount of spray solution that can be applied the day of mixing. *V-10116 VPP* should be applied within 12 hours of mixing.

SPRAYER CLEANUP

Spray equipment should be cleaned each day following *V-10116 VPP* application. After *V-10116 VPP* is applied, the following steps should be used to clean the spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
3. Drain tank completely.
4. Remove all nozzles and screens and rinse them in clean water.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. Do not apply this product when weather conditions favor spray drift from treated areas. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making decisions. Where states have more stringent regulations, they should be observed.

TURFGRASS

- *V-10116 VPP* may be used for disease control on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, athletic fields and sod farms.
- *V-10116 VPP* should be used in conjunction with turf management practices that promote good plant health and optimum disease control. Proper identification of disease infestations is essential for providing control. Consult diagnostic guides, kits or other means of identification to determine the best control measures.
- *V-10116 VPP* is more effective when used preventatively or early in disease development. Length of disease control will vary according to local environmental conditions, disease pressure and turfgrass management practices.
- Apply to dry turf or after mowing. For control of foliar diseases, allow turf to dry before irrigation. For control of soil born diseases, *V-10116 VPP* can be watered in after application.
- *V-10116 VPP* may be tank mixed with other registered turfgrass fungicides. Perform a jar test as describe above to determine compatibility before making applications. Directions for use of *V-10116 VPP* have been developed using no adjuvant in the spray solution.
- Applications of *V-10116 VPP* can injure golf course greens that are under high heat (temperatures exceeding 90°F) and/or drought stress. Under high heat and/or drought stress conditions (mid to late summer), limit applications to no more than 16 oz per acre (0.37 oz per 1,000 sq ft) every 30 days.
- Important: Under conditions of optimal disease development (such as high humidity and temperature), use higher rates and shorter application intervals. When applications are made curatively, use higher rates of *V-10116 VPP* and short application intervals. Use higher application rate in areas with a history of severe disease pressure.
- All applications should be made according to the following use directions. Failure to follow label directions may result in turfgrass injury and or inferior disease control.

APPLICATION DIRECTIONS:

Apply *V-10116 VPP* using standard, low-pressure spray equipment designed to deliver coarse to medium size spray droplets (nozzles designed to deliver fine droplets should be avoided). Apply in a sufficient volume of water to provide thorough spray coverage and a uniform spray pattern. Spray boom height should be no higher than 20 inches. For foliar disease, use a minimum of 2 gals of water per 1,000 sq ft For crown and root diseases, use 2 to 4 gals of water per 1,000 sq ft For spot treatment, use 0.2 oz per 1 to 2 gals of water. Applications may be made by ground only.

TURFGRASS TOLERANCE

Due to the large number of turfgrass types and possible tank mixes and application techniques, it has not been determined if *V-10116 VPP* can be safely used on all turfgrasses under all conditions. It is recommended that the user determine safety under

12/18

local conditions by making an application to a small test area. Monitor the test area for at least 2 weeks to determine safety under local growing conditions.

RECOMMENDED TURFGRASS SPECIES

- Annual Bluegrass
- Bentgrass
- Centipedegrass
- Fescue
- Kentucky Bluegrass
- Perennial Ryegrass
- St. Augustinegrass
- Seashore Paspalum
- Zoysia

RESTRICTIONS AND LIMITATIONS

- Do not apply more than 4 lbs (2 lbs ai) of V-10116 VPP per acre per year.
- Do not use on bermudagrass.
- Do not apply through any type of irrigation system.
- Do not apply by aerial application.
- Do not graze animals on treated areas.
- Do not feed clippings from treated areas to livestock or poultry.

13/18

Table 1. TURFGRASS – Specific Diseases, Rates, Timings and Application Information

Diseases (Pathogens)	Application Rates		Application Interval/ Timing	Special Instructions
	oz/1,000 sq ft	oz/A		
Dollar Spot (<i>Sclerotinia homeocarpa</i>)	0.18 to 0.37	8 to 16	14 to 21 days	Use preventatively. Begin application when conditions favor disease development.
Brown Patch (<i>Rhizoctonia solani</i>)	0.28 to 0.37	12 to 16	14 to 21 days	Begin applications when conditions favor disease development.
Anthracnose (<i>Colletotrichum graminicola</i>)	0.28 to 0.37	12 to 16	14 to 21 days	Begin applications when conditions favor disease development.
Gray Leaf Spot (<i>Pyricularia grisea</i>) Red Thread (<i>Laetisaria fuciformis</i>) Rhizoctonia Large Patch (<i>Rhizoctonia solani</i>) Rust Diseases (<i>Puccinia</i> spp.) Summer Patch (<i>Magnaporthe poae</i>) Zoysia Patch (<i>Rhizoctonia solani</i>)	0.37	16	14 days	Begin applications when conditions favor disease development or when the disease first appears.

continued

14/18

Table 1. TURFGRASS – Specific Diseases, Rates, Timings and Application Information

Diseases (Pathogens)	Application Rates		Application Interval/ Timing	Special Instructions
	oz/1,000 sq ft	oz/A		
Fairy Ring (Various <i>basidiomycetes</i>)	0.37	16	21 days	Apply as soon as possible after first symptoms appear. Apply in a volume of 4 gals per 1,000 sq ft For optimal control reapply after 21 days. Symptoms may take several weeks to disappear following application.
Pink Snow Mold Fusarium Patch (<i>Microdochium nivale</i>) Gray Snow Mold (<i>Typhula</i> spp.) Yellow Patch (<i>Rhizoctonia cerealis</i>)	0.37 to 0.44	16 to 19.2	Late Fall	Make 1 to 2 applications in combination with PCNB, chlorothalonil or other fungicides active on snow mold in late fall before snow cover and if possible, one application during a mid-winter thaw.

V-10116 VPP Rate Conversion Chart for Turfgrass

oz product/ 1000 sq ft	oz product/acre	lbs product/acre	lbs ai/acre
0.18	8	0.5	0.25
0.28	12	0.75	0.38
0.37	16	1.0	0.50
0.44	19.2	1.2	0.60

15/18

ORNAMENTAL PLANTS

- *V-10116 VPP* may be used for disease control on ornamental plants in commercial indoor and outdoor nurseries, greenhouses and ornamental plants found in landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.
- Use *V-10116 VPP* as part of a preventative disease control program.
- *V-10116 VPP* is more effective when used preventatively or early in disease development. Length of disease control will vary according to local environmental conditions, disease pressure and management practices.
- All applications should be made according to the following use directions. Failure to follow label directions may result in injury and or inferior disease control.

ORNAMENTAL PLANT TOLERANCE

IMPORTANT: *V-10116 VPP* may cause growth regulation to some ornamental plant species. The large number of existing ornamental species and their varieties and cultivars coupled with the constant introduction of new varieties makes it impossible to field test *V-10116 VPP* in every locale where it is sold or in all of the combinations of location and plant variety. Further differences include the soil or media type, pH, moisture or fertility, environmental conditions such as temperature, lighting or degree-days and horticultural practice and the manner of use and application of this product. To ensure that *V-10116 VPP* is compatible with the ornamental plant variety or cultivar under your specific conditions, test this product on a small test area and observe for phytotoxicity or other unintended effects for two weeks before making large-scale applications.

APPLICATION DIRECTIONS

Foliar applications

- Apply *V-10116 VPP* at rates of 2 to 8 oz of product per 100 gals of water for disease control in ornamental plants.
- Apply as a full coverage spray to the point of drip.
- Apply as needed on a 14 to 28 day interval.
- Under conditions of optimal disease development (such as high humidity and temperature), use higher rates and shorter application intervals.
- When applications are made curatively, use higher rates of *V-10116 VPP* and shorter application intervals. Use higher application rate in areas with a history of severe disease pressure.
- For general disease control in landscapes, apply 4 to 8 oz of product per 100 gals of water on a 21 day interval.

16/18

Drench applications

- To control soil born, seedling and crown diseases, apply *V-10116 VPP* as a preventative drench treatment prior to disease development.
- Apply at 1 to 4 oz per 100 gals of water in a volume of 1 to 2 pts of solution per square foot of surface area on a 7 to 28 day interval.
- Application to healthy roots is necessary to ensure systemic uptake, translocation and disease protection.

COMPATIBILITY

V-10116 VPP is compatible with most commonly used insecticides, fungicides and spray adjuvants used in the production of ornamental plants. When using *V-10116 VPP* in tank mixes with other pesticides, observe all directions for use and precautions on the respective tank mix label. When making an application of a tank mix for the first time, it is recommended that physical compatibility be checked and that a few plants be treated and observed for phytotoxicity or other unintended effects for 2 to 4 weeks before making large scale applications.

APPLICATION EQUIPMENT

Apply *V-10116 VPP* dispersed in water using suitable hand or power operated application equipment such as portable, pump-up, backpack, hydraulic or boom sprayers in a manner to provide complete and uniform coverage.

RESTRICTIONS AND LIMITATIONS

- Do not apply more than 4 lbs (2 lbs ai) of product per acre per year.
- Do not apply through any type of irrigation system.
- Do not apply by aerial application.
- Do not graze animals on treated areas.
- Do not feed plants from treated areas to livestock or poultry.

17/18

Table 2. DIRECTIONS FOR USE ON ORNAMENTALS

Diseases	Pathogen	Special Instructions
Conifer Blight <i>Phomopsis Blight</i>	<i>Phomopsis juniperovora</i>	Apply 2 to 8 oz/100 gals on a 14 to 21 day interval.
Flower Blight <i>Anthracnose</i>	<i>Colletotrichum</i> spp.	Apply 2 to 8 oz/100 gals on a 14 to 21 day interval.
Leaf Blights/Spots <i>Anthracnose</i>	<i>Colletotrichum</i> spp.	Apply 2 to 8 oz/100 gals on a 14 to 21 day interval.
Downy Mildew	<i>Colletotrichum</i> spp.	Apply 2 to 8 oz/100 gals on a 14 to 21 day interval.
Powdery Mildew <i>Podosphaera xanthii</i>	<i>Microspora azaleae</i>	Apply 2 to 8 oz/100 gals on a 14 to 21 day interval.
Rusts	<i>Puccinia</i> spp.	Apply 2 to 8 oz/100 gals on a 14 to 21 day interval.
Soil Born Disease (Drench) <i>Rhizoctonia solani</i> <i>Sclerotium rolfsii</i>	<i>Fusarium</i> spp.	Apply 1 to 4 oz/100 gals of water in a volume of 1 to 2 pts of solution per sq ft of surface area on a 7 to 28 day interval.

Table 3. ORNAMENTAL PLANT LIST

Flowering and Foliage Plants	Herbaceous Ornamental Plants	Woody Ornamental Plants	Non-Bearing Tree, Fruit, Nut and Vine
Begonia Geranium Pansy Petunia Poinsettia Snapdragon	Carnation Chrysanthemum Dianthus Iris Marigold Phlox Verbena Zinnia	Ash Azalea Crabapple Crape Myrtle Dogwood Juniper Lilac Oaks Pines Poplars Photinia Rhododendron Rose	Apple Cherry Citrus Peach Pecan Plum Walnut

18/18

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

PESTICIDE STORAGE

Store in a cool dry place.

Keep pesticide in original container.

Keep container closed when not in use.

Do not put concentrate or dilute into food or drink containers.

Not for use or storage in or around the home.

Do not store at temperature below 32°F. If the product is exposed to temperatures below 32°F, thaw at 50°F or higher and shake gently to unify the product.

For help with any spill, leak, fire or exposure involving this material, call day or night 800-892-0099.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Do not reuse container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Copyright© 2007 by Valent U.S.A. Corporation

Manufactured for:

Valent U.S.A. Corporation

P.O. Box 8025

Walnut Creek, CA 94596-8025

www.valentpro.com

Made in U.S.A.

EPA Reg. No. 59639-RUU

EPA Est. No. _____