



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 4, 2019

Zenna Burke
Senior Regulatory Manager
Valent U.S.A. LLC
1600 Riviera Ave., Suite 200
Walnut Creek, CA 94596

Subject: Label Amendment – California use restriction on ornamentals; changes according to PRN 2017-1
Product Name: V-10116 VPP Fungicide
EPA Registration Number: 59639-144
Application Date: March 14, 2019
Decision Number: 549601

Dear Ms. Burke:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.


A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 59639-144
Decision No. 549601

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact me by phone at (703) 305-5410, or via email at johnson.hope@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to be 'HJ', written in a cursive style.

Hope Johnson, Product Manager 21
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure



METCONAZOLE GROUP 3 FUNGICIDE

V-10116 VPP Fungicide

Turfgrass Fungicide

A SYSTEMIC FUNGICIDE FOR CONTROL OF THE LISTED TURFGRASS DISEASES

Ornamental Fungicide

A SYSTEMIC FUNGICIDE FOR CONTROL OF THE LISTED ORNAMENTAL DISEASES

Active Ingredient	By Wt
Metconazole*.....	50%
Other Ingredients.....	50%
Total	100%

*5-[(4-chlorophenyl)methyl]-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl)cyclopentanol

V-10116 VPP Fungicide is a water dispersible granule containing 50% active ingredient.


KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE NEXT PAGE[PANEL][BOOKLET] FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA Reg. No. 59639-144
EPA Est.

NET WEIGHT 5 POUNDS



FIRST AID	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 800-892-0099 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long-sleeved shirt and long pants, socks and shoes.

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS
<p>Users should:</p> <ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to shrimp, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposing of equipment washwaters or rinsate. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas.

Groundwater Advisory: This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils or soils with shallow water tables are more prone to runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features, such as ponds, streams and springs, will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is coveralls, chemical-resistant gloves made of any waterproof material, socks and shoes.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to the uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, greenhouses or sodfarms.

Keep all unprotected persons out of operating areas or vicinity where there may be drift.
Do not enter or allow others to enter treated areas until sprays have dried.

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests and resistance of the target pest to this product. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. To the extent allowed by law Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label **and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY

LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements, Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law, if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability**, which may not be modified by any oral or written agreement.

PRODUCT INFORMATION

V-10116 VPP Fungicide is formulated as 50% water dispersible granular (WDG). The active ingredient in V-10116 VPP Fungicide is metconazole, a broad-spectrum triazole fungicide that works by inhibiting demethylation and other processes in sterol biosynthesis. V-10116 VPP Fungicide is systemic and is quickly absorbed into plant tissue. V-10116 VPP Fungicide displays preventive and curative properties.

V-10116 VPP Fungicide may be used for disease control in turfgrass on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, athletic fields and sod farms.

V-10116 VPP Fungicide may be used for disease control on ornamental plants in commercial indoor and outdoor nurseries, greenhouses and on ornamental plants found in landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

V-10116 VPP Fungicide provides optimal disease control when applied in combination or as part of a rotation in a regularly scheduled spray program with other effective fungicides that have different modes of action (i.e., non Group 3 fungicides).

MODE OF ACTION

The active ingredient in V-10116 VPP Fungicide, metconazole, belongs to the sterol biosynthesis inhibitor group of fungicides as classified by the U.S. EPA and Canada PMRA as a target site of action Group 3 fungicide.

Resistance Management

For resistance management, V-10116 VPP Fungicide contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to V-10116 VPP Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of V-10116 VPP Fungicide or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Avoid application of more than the maximum number of applications listed in the directions for use and follow label instructions regarding sequential applications of V-10116 VPP Fungicide or other fungicides in the same groups in a season.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Valent U.S.A. LLC at 800-89-VALENT (898-2536). You can also contact your pesticide distributor or university extension specialist to report resistance.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND V-10116 VPP

Perform a jar test before mixing commercial quantities of V-10116 VPP Fungicide, when using V-10116 VPP Fungicide for the first time, when using new adjuvants, when using new tank mixes or when a new water source is being used.

1. Add 1 pt of the water to a quart jar. Use water from the same source and temperature as water that will be used in the spray tank mixing operation.
2. Add about 1 teaspoon (2.4 grams) of V-10116 VPP Fungicide to the quart jar; gently mix until product goes into suspension.
3. Add appropriate amount of each tank mix adjuvant or partner.
4. Place cap on jar, invert 10 times, let stand for 15 minutes and evaluate.
5. An ideal tank mix combination will be uniform and free of large suspended particles. If any of the following conditions are observed, reevaluate the choice of the adjuvant:
 - a) Layer of oil or globules on the mixture's surface.
 - b) Flocculation: fine particles in suspension or as a layer on the bottom of the jar.
 - c) Clabbering: Thickening texture (coagulated) like gelatin.

TANK MIX PARTNERS

V-10116 VPP Fungicide is compatible with most herbicides, insecticides, fungicides and other types of tank mix components. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Do not tank mix with any product which has a prohibition on tank mixing. When evaluating potential tank mix partners, always apply to a small area of target plants and then monitor for adverse affects for at least two weeks before making large-scale applications.

SPRAYER PREPARATION

Before applying V-10116 VPP Fungicide, start with clean, well maintained application equipment. The spray tank hoses and booms must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply V-10116 VPP Fungicide. Check nozzles frequently for accuracy. If two or more products were tank mixed prior to V-10116 VPP Fungicide application, follow the most restrictive cleanup procedure.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, slowly add the V-10116 VPP Fungicide to the spray tank. A rippling or rolling action on the water surface will occur as the result of the agitation.
3. If tank mixing V-10116 VPP Fungicide with other labeled pesticides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions.
4. Add any required adjuvants.
5. Fill spray tank to desired level with water. Continue to agitate until all spray solution has been applied.
6. Mix only the amount of spray solution that can be applied the day of mixing. Apply V-10116 VPP Fungicide within 12 hours of mixing.

SPRAYER CLEANUP

Clean the spray equipment each day following V-10116 VPP Fungicide application. After V-10116 VPP Fungicide is applied, follow the steps below to clean the spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.

3. Drain tank completely.
4. Remove all nozzles and screens and rinse them in clean water.

SPRAY DRIFT MANAGEMENT

It is the responsibility of the applicator to ensure that spray drift does not occur. Do not apply this product when weather conditions favor spray drift from treated areas. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making decisions. Where states have more stringent regulations, they must be observed.

TURFGRASS

- Use V-10116 VPP Fungicide for disease control on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, athletic fields and sod farms.
- Use V-10116 VPP Fungicide in conjunction with turf management practices that promote good plant health and optimum disease control. Proper identification of disease infestations is essential for providing control. Consult diagnostic guides, kits or other means of identification to determine the best control measures.
- V-10116 VPP Fungicide is more effective when used preventively or early in disease development. Length of disease control will vary according to local environmental conditions, disease pressure and turfgrass management practices.
- Apply to dry turf or after mowing. For control of foliar diseases, allow turf to dry before irrigation. For control of soilborne diseases, V-10116 VPP Fungicide can be watered in after application.
- V-10116 VPP Fungicide may be tank mixed with other registered turfgrass fungicides. Perform a jar test as describe above to determine compatibility before making applications. Directions for use of V-10116 VPP Fungicide have been developed using no adjuvant in the spray solution.
- V-10116 VPP Fungicide can injure golf course greens that are under high heat (temperatures exceeding 90°F) and/or drought stress. Under high heat and/or drought stress conditions (mid to late summer), do not apply more than 16 oz per acre (0.37 oz per 1,000 sq ft) every 30 days.
- Important: Under conditions of optimal disease development (such as high humidity and temperature), use higher labeled rates and shorter application intervals. When applications are made curatively, use higher specified rates of V-10116 VPP Fungicide and shorter specified application intervals. Use the higher specified application rate in areas with a history of severe disease pressure.
- Failure to follow label directions may result in turfgrass injury and or inferior disease control.

APPLICATION DIRECTIONS

Apply V-10116 VPP Fungicide using standard, low-pressure spray equipment designed to deliver coarse to medium size spray droplets (do not use nozzles designed to deliver fine droplets). Apply in a sufficient volume of water to provide thorough spray coverage and a uniform spray pattern. Do not position the spray boom more than 20 inches above the turf. For foliar disease, apply 2 gallons of spray solution per 1,000 sq ft. For crown and root diseases, apply 2 to 4 gallons of spray solution per 1,000 sq ft. For spot treatment, mix 0.1 to 0.2 oz of V-10116 VPP Fungicide per gallon of water.

TURFGRASS TOLERANCE

Due to the large number of turfgrass types, possible tank mixes and application techniques, it has not been determined if V-10116 VPP Fungicide can be safely used on all turfgrasses under all conditions. Before making a large scale application, determine safety in a small test area. Monitor the test area for at least 2 weeks to determine safety under local growing conditions.

TURFGRASS SPECIES

Annual Bluegrass
Bentgrass
Bermudagrass
Centipedegrass
Fescue
Kentucky Bluegrass
Perennial Ryegrass
St. Augustinegrass
Seashore Paspalum
Zoysia

TURFGRASS RESTRICTIONS

- Do not apply more than 4 lb (2 lb ai) of V-10116 VPP Fungicide per acre per year.
- Do not apply directly to bermudagrass greens.
- Do not apply through any type of irrigation system.
- Do not apply by air.
- Do not graze animals on treated areas for one year after application.
- Do not feed clippings from treated areas to livestock or poultry.
- Do not exceed 0.2 oz product per gallon of water for spot treatment.

Table 1. TURFGRASS – Specific Diseases, Rates, Timings and Application Information

Diseases (Pathogens)	Application Rates		Application Interval/ Timing	Special Instructions
	oz/1,000 sq ft	oz/A		
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	0.18 to 0.37	8 to 16	14 to 21 days	Use preventively. Begin application when conditions favor disease development.
Brown Patch (<i>Rhizoctonia solani</i>) Anthracnose (<i>Colletotrichum graminicola</i>)	0.28 to 0.37	12 to 16	14 to 21 days	Begin applications when conditions favor disease development.
Gray Leaf Spot (<i>Pyricularia grisea</i>) Necrotic Ring Spot (<i>Ophiosphaerella korrae</i>) Red Thread (<i>Laetisaria fuciformis</i>) Rhizoctonia Large Patch (<i>Rhizoctonia solani</i>) Rust Diseases (<i>Puccinia</i> spp.) Summer Patch (<i>Magnaporthe poae</i>) Take-all Patch (<i>Gaeumannomyces graminis</i>) Waitea Patch Brown Ring Patch (<i>Waitea circinata</i>) Zoysia Patch (<i>Rhizoctonia solani</i>)	0.37	16	14 days	Begin applications when conditions favor disease development or when the disease first appears.
Fairy Ring (<i>Various Basidiomycetes</i>)	0.37	16	21 days	Apply as soon as possible after first symptoms appear. Apply in a volume of 4 gal per 1,000 sq ft. For optimal control reapply after 21 days. Symptoms may take several weeks to disappear following application.
Pink Snow Mold Fusarium Patch (<i>Microdochium nivale</i>) Gray Snow Mold (<i>Typhula</i> spp.) Yellow Patch (<i>Rhizoctonia cerealis</i>)	0.37 to 0.44	16 to 19.2	Late Fall	Make 1 to 2 applications in combination with PCNB, chlorothalonil or other fungicides active on snow mold in late fall before snow cover and if possible, one application during a mid-winter thaw.

V-10116 VPP Rate Conversion Chart for Turfgrass

oz product/1000 sq ft	oz product/acre	lb product/acre	lb ai/acre
0.18	8	0.5	0.25
0.28	12	0.75	0.38
0.37	16	1.0	0.50
0.44	19.2	1.2	0.60

1 level teaspoon contains 2.4 grams, and one 8 fl oz cup contains 4.1 oz of V-10116VPP Fungicide

ORNAMENTAL PLANTS

[Do not use in California]

- V-10116 VPP Fungicide provides disease control on ornamental plants in commercial outdoor nurseries and ornamental plants found in landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.
- V-10116 VPP Fungicide provides disease control on ornamental plants in indoor greenhouses and nurseries
- Use V-10116 VPP Fungicide as part of a preventive disease control program.
- V-10116 VPP Fungicide is more effective when used preventively or early in disease development. Length of disease control will vary according to local environmental conditions, disease pressure and management practices.
- Follow all use directions on this label. Failure to follow label directions may result in injury and or inferior disease control.

ORNAMENTAL PLANT TOLERANCE

IMPORTANT: V-10116 VPP Fungicide may cause growth regulation to some ornamental plant species. The large number of existing ornamental species and their varieties and cultivars coupled with the constant introduction of new varieties makes it impossible to field test V-10116 VPP Fungicide in every locale where it is sold or in all of the combinations of location and plant variety. Further differences include the soil or media type, pH, moisture or fertility, environmental conditions such as temperature, lighting or degree-days and horticultural practice and the manner of use and application of this product. To ensure that V-10116 VPP Fungicide is compatible with the ornamental plant variety or cultivar under your specific conditions, test this product on a small test area and observe for phytotoxicity or other unintended effects for at least two weeks before making large-scale applications.

APPLICATION DIRECTIONS

Foliar applications

- Apply V-10116 VPP Fungicide at 1 to 4 oz of product per 100 gallon of water for disease control in ornamental plants.
- Apply as a full coverage spray to the point of run-off.
- Apply as needed on a 14 to 28 day interval.
- Under conditions of optimal disease development (such as high humidity and temperature), use higher specified rates and shorter specified application intervals.
- When applications are made curatively, use higher specified rates of V-10116 VPP Fungicide and shorter specified application intervals. Use the higher specified application rate in areas with a history of severe disease pressure.
- For general disease control in landscapes, apply 1 to 4 oz of product per 100 gallon of water on a 21- to 28-day interval.

Drench applications

- To control soilborne, seedling and crown diseases, apply V-10116 VPP Fungicide as a preventive drench treatment prior to disease development.
- Apply at 1 to 4 oz per 100 gallon of water in a volume of 1 to 2 pt of solution per square foot of surface area on a 14 to 28 day interval.
- Roots must be healthy to ensure systemic uptake, translocation and disease protection.

COMPATIBILITY

V-10116 VPP Fungicide is compatible with most commonly used insecticides, fungicides and spray adjuvants used in the production of ornamental plants. When using V-10116 VPP Fungicide in tank mixes with other pesticides, observe all directions for use and precautions on the label of tank mix partner. Before applying a tank mix for the first time, check physical compatibility of the mixture. Do not tank mix with any product which contains a prohibition on tank mixing.

APPLICATION EQUIPMENT

Apply V-10116 VPP Fungicide using suitable hand or power operated application equipment such as portable, pump-up, backpack, hydraulic or boom sprayers in a manner to provide complete and uniform coverage.

ORNAMENTAL RESTRICTIONS

- Do not apply more than 4 lb (2 lb ai) of V-10116 VPP Fungicide per acre per year.
- Do not apply through any type of irrigation system.
- Do not apply by air.
- Do not graze animals on treated areas of one year after application.
- Do not feed plants from treated areas to livestock or poultry.
- [Do not use in California.]

Table 2. DIRECTIONS FOR USE ON ORNAMENTALS

Diseases*	Pathogen	Application Rates (By Weight)	Special Instructions
Anthracnose	<i>Apiognomonina</i> spp. <i>Colletotrichum</i> spp. <i>Cryptocline</i> spp. <i>Discula</i> spp. <i>Gleosporium</i> spp.	1 to 4 oz/100 gal on a 14- to 28-day interval	When using a typical high volume sprayer use 100 gal of spray mix to treat 20,000 sq ft of area. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
Botrytis	<i>Botrytis</i> spp.		
Blossom Rot/Brown Rot	<i>Monilinia</i> spp.		
Boxwood Blight	<i>Calonectria pseudonaviculatum</i> <i>Cylindrocladium buxicola</i> <i>Cylindrocladium pseudonaviculatum</i>	4 oz/100 gal on a 14- to 28-day interval	
Conifer Blight	<i>Diplodia pinea</i> <i>Phomopsis</i> spp. <i>Rhabdocline</i> spp. <i>Rhizosphaera</i> spp. <i>Sphaeropsis</i> spp.	2 to 4 oz/100 gal on a 14- to 28-day interval	
Damping-Off/Stem and Root Rots	<i>Rhizoctonia solani</i>	1 to 4 oz/100 gal on a 14- to 28-day interval	
Eastern Filbert Blight	<i>Anisogramma anomola</i>		
Fusarium Root Rot and Damping Off	<i>Fusarium oxysporum</i> <i>Fusarium solani</i>	4 oz/100 gal on a 14- to 28-day interval	
Leaf Spots	<i>Alternaria</i> spp. <i>Cercospora</i> spp. <i>Diplocarpon</i> spp. <i>Mycosphaerella</i> spp. <i>Myrothecium</i> spp. <i>Phyllosticta</i> spp. <i>Septoria</i> spp.		
Powdery Mildew	<i>Erysiphe</i> spp. <i>Microsphaera</i> spp. <i>Oidium</i> spp. <i>Phyllactinia</i> spp. <i>Podosphaera</i> spp. <i>Sphaerotheca</i> spp.	1 to 4 oz/100 gal on a 14- to 28-day interval	
Rusts	<i>Chrysomyxa</i> spp. <i>Coleosporium</i> spp. <i>Gymnosporangium</i> spp. <i>Puccinia</i> spp. <i>Uredo</i> spp. <i>Uromyces</i> spp.		

*If symptoms are visible of a labeled disease, use a registered contact, protectant fungicide.

Table 2. DIRECTIONS FOR USE ON ORNAMENTALS (continued)

Diseases*	Pathogen	Application Rates (By Weight)	Special Instructions
Scab	<i>Cladosporium</i> spp. <i>Sphaceloma</i> spp. <i>Venturia</i> spp.		When using a typical high volume sprayer use 100 gal of spray mix to treat 20,000 sq ft of area. If using a low volume sprayer, adjust concentration to apply the same amount of product per unit area.
Shot Hole	<i>Wilsonomyces carpophilus</i> <i>Blumeriella</i> spp.		
White Mold	<i>Sclerotinia sclerotium</i>	2 to 4 oz/100 gal on a 14- to 28-day interval	

*If symptoms are visible of a labeled disease, use a registered contact, protectant fungicide.

Table 3. ORNAMENTAL PLANT LIST

Note: V-10116 VPP Fungicide has been tested on the following plants with no crop injury.

Flowering and Foliage Plants	Herbaceous Ornamental Plants	Woody Ornamental Plants	Non-Bearing Tree, Fruit, Nut and Vine
Geranium Petunia Poinsettia Snapdragon	Carnation Chrysanthemum Dalily Dianthus Hydrangea Iris Lantana Liriope Marigold Phlox Verbena	Amelanchier Ash Azalea Bayberry Beech Boxwood Birch Camellia Conifers Cotoneaster Crabapple Crape Myrtle Dogwood Elm Euonymus Fir Hawthorn Hemlock Holly Juniper Laurel Lilac Linden London Plane Magnolia Maple Mountain Ash Mountain Laurel Oaks Pine Poplar Photinia Pyracantha Rhododendron Rose Sassafrass Spirea Spruce Sweetgum Sycamore	Apple Apricot Cherry Citrus Filbert Grape Peach Pear Pecan Pistachio Plum Walnut

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in a cool dry secure place.

Keep pesticide in original container.

Keep container closed when not in use.

Do not put concentrate or dilute into food or drink containers.

Do not store in or around the home.

For help with any spill, leak, fire or exposure involving this material, call day or night
800-892-0099.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

©2019 Valent U.S.A. LLC

Manufactured for:

Valent U.S.A. LLC

P.O. Box 8025

Walnut Creek, CA 94596-8025



Nufarm

Grow a better tomorrow.

Distributed by:

Nufarm Americas Inc.

11901 S. Austin Ave.,

Alsip, IL 60803

Made in U.S.A.

EPA Reg. No. 59639-144

EPA Est. No. _____

059639-00144.20190927.V10116VPP.AMEND.Clean