59639-151 7.21-2010



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Sue-Chi Shen Valent U.S.A. Corporation 1600 Riviera Avenue, Suite 200 Walnut Creek, CA 94596

JUL 2 1 2010

Dear Ms. Shen:

Subject: Supplemental Label; Associated with Master Label Stamped on July 20, 2010 V-10170 5 FS Insecticide EPA Registration No. 59639-151 Date Submitted: July 21, 2010


The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records. Please submit one (1) final printed copy for the above mentioned label before releasing the product for shipment. If you have any questions regarding this label, please contact Kable Bo Davis at (703) 306-0415 or

davis.kable@epa.gov.

Sincerely yours,

Venus Eagle Product Manager (01) Insecticide-Rodenticide Branch Registration Division 7505P

Enclosed: Stamped Supplemental Label



UPDATED LABELING OF TREATED SEED

This supplemental label expires on January 20, 2012 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR V-10170 5FS INSECTICIDE BEFORE APPLYING. USE OF V-10170 5 FS INSECTICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR V-10170 5 FS INSECTICIDE.

LABELING OF TREATED SEED

The Federal Seed Act requires that the container of seed treated with *V*-10170 5 FS Insecticide must be labeled with the following information:

- This seed has been treated with V-10170 5 FS Insecticide which containsclothianidin.
- Do not use treated seed for feed, food, or oil processing.

In addition, the U.S. Environmental Protection Agency requires the following statements on the container of seed treated with *V*-10170 5 FS Insecticide:

- Regardless of the type of application (seed treatment, soil or foliar) do not apply more than a total of 6 oz clothianidin (active ingredient) per acre per cropping cycle except for soybean. Regardless of the type of application (seed treatment, soil or foliar) do not apply more than a
- total of 3.2 oz clothianidin (active ingredient) per acre per cropping cycle for soybean.
- Rapeseed greens and seed grown or harvested from V-10170 5 FS Insecticide treated seed must not be used for feed or human consumption. Rapeseed grown and harvested from V-10170 5 FS Insecticide treated seed is only for industrial uses and cannot be used for edible oil or any human/feed consumption.
- Store away from feeds and other foodstuffs.
- Wear long sleeved shirt, long pants and waterproof gloves when handling treated seed.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.

- Do not graze or feed soybean forage and hay to livestock.
- Areas planted with treated seed of cereal grains (except rice, sorghum, sweet corn and wild rice), field corn, popcorn, sugar beet and soybean may be replanted immediately with bulb vegetables, canola, cereal grains (except rice), cotton, corn (field, pop and sweet), fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), rapeseed, root and tuber vegetables, sorghum, soybean and sugar beet. These areas may also be replanted after 30 days with dry beans, grasses, non-grass animal feeds and rice. These areas may also be replanted after 8 months with sugarcane. Do not plant any other crop in the treated area for at least one year after these treated seeds are planted.
- Areas planted with treated seed of canola, rapeseed, sorghum and sweet corn may be replanted immediately with bulb vegetables, canola, cereal grains (except rice), cotton, corn (field, pop and sweet), fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), rapeseed, root and tuber vegetables, sorghum, soybean and sugar beet. These areas may also be replanted after 30 days with dry beans, grasses, non-grass animal feeds and rice. Do not plant any other crops in the treated area for at least 4 months after these treated seeds are planted.

Cereal Grains Except Corns (Field, Pop and Sweetcorn), Rice, Sorghum and Wild Rice		
Insect Pests	V10170 5 FS Insecticide Use Rates fl oz/ 100 lbs seed	Further Use Instructions
Wireworm	0.25 – 1.79	For wireworm seed protection, use a 0.25 fl oz/100 lbs seed rate or higher, depending on field history and site populations.
Aphid	0.75 – 1.79	
Wireworm	0.25 – 1.79	For wireworm seed protection, use a 0.25 fl oz/100 lbs seed rate or higher, depending on field history and site populations.
Aphid	0.75 – 1.79	
Hessian Fly	1.79	For Hessian Fly protection to wheat seedlings in the Fall season, use 1.79 fl oz/100 lbs seed.
	Rice Insect Pests Wireworm Aphid Wireworm Aphid Aphid	Rice, Sorghum and WildInsect PestsV10170 5 FS Insecticide Use Rates fl oz/ 100 lbs seedWireworm0.25 – 1.79Aphid0.75 – 1.79Wireworm0.25 – 1.79

• Regardless of the type of application (seed treatment, soil, or foliar) do not apply more than a total of 6 oz clothianidin (active ingredient) per acre per cropping cycle.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Copyright © 2010 by Valent U.S.A. Corporation

Manufactured for: Valent U.S.A. Corporation P. O. Box 8025 Walnut Creek, CA 94596-8025

Made in U.S.A.

059639-00152.20100719.V101705FS.Supp