



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

October 14, 2025

Zenna Scarlett
zenna.scarlett@valent.com
VALENT U.S.A. LLC

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment - Update
resistance management language and other minor edits.
Product Name: ARENA 0.25 G INSECTICIDE
Admin Number: 59639-157
EPA Receipt Date: 04/25/2019
Action Case Number: 00605490

Dear Zenna Scarlett:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have questions, please contact Ralph Narain by telephone at 202-566-2853 or via email at narain.ralph@epa.gov.

Sincerely,

Ralph Narain

Ralph Narain, Entomologist/Risk Manager
IVB3, RD
Office of Pesticide Programs



CLOTHIANIDIN	GROUP	4A	INSECTICIDE
--------------	-------	----	-------------

ARENA[®] 0.25 G

INSECTICIDE

FOR SYSTEMIC INSECT CONTROL IN TURFGRASS, SOD FARMS, LANDSCAPE ORNAMENTALS, INTERIOR PLANTSCAPES AND NON-BEARING FRUITS AND NUTS (IN LANDSCAPES).

Active Ingredient	By wt.
*Clothianidin.....	0.25%
Other Ingredients	99.75%
Total	100.00%

*(E)-1-(2-chloro-1,3-thiazol-5-ylmethyl)-3-methyl-2-nitroguanidine.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE NEXT PAGE [BACK] [PANEL][BOOKLET] FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA Reg. No. 59639-157

EPA Est. No. _____

Manufactured for:

Valent U.S.A. LLC

P.O. Box 8025

Walnut Creek CA 94596-8025

ACCEPTED

Oct 14, 2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg. No. 59639-157

Net Contents:

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 800-892-0099 for emergency medical treatment information.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, chemical-resistant gloves made of any waterproof material including polyethylene or polyvinyl chloride.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If there are no instructions for washables available, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

This product is toxic to bees exposed to direct treatment or residues on blooming crops. Do not apply during bloom or when bees are present.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH
PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE AND WITH
APPLICABLE STATE AND FEDERAL REGULATIONS.**

Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, socks and shoes, chemical-resistant gloves made of any water proof material.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in the box apply to use of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPA applies when this product is used to produce agricultural plants on farms, forests, nurseries and commercial greenhouses.

Keep children and pets off treated area until the dust has settled.

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability**, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law. Read and follow the entire label of each product to be used in the tank mix with this product.

RESISTANCE MANAGEMENT

For resistance management, *Arena* 0.25G Insecticide contains a Group 4A insecticide. Any insect population may contain individuals naturally resistant to *Arena* 0.25G Insecticide and other Group 4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of *Arena* 0.25G Insecticide or other Group 4A insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues between the individual components of the mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee.
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests that the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance contact Valent U.S.A. LLC at 1-800-89-VALENT (898-2536).

PRODUCT INFORMATION

Arena 0.25 G Insecticide is a broad spectrum insecticide, for control of listed insects infesting turfgrass, sod farms, landscape ornamentals, interior plantscapes and non-bearing fruit trees (including apple and pear) and nut trees (in landscapes). When applied as directed, *Arena* 0.25 G Insecticide provides excellent and long residual insect control.

RESTRICTIONS

- Do not use a foliar application of clothianidin (*Arena* 0.25 G Insecticide) following a soil application of clothianidin.
- Do not apply more than a total of 160 lb of *Arena* 0.25 G Insecticide (0.4 lb ai/A) per acre per season.
- If the maximum season limit of clothianidin as mentioned above has been applied and pest populations require additional treatments, use another registered pesticide that is not in the neonicotinoid class of chemistry.
- Do not apply this product, by any application method, to linden, basswood or other *Tilia* species.

PLANT TOLERANCE

Neither the manufacturer nor the seller has determined whether or not *Arena* 0.25 G can be used safely on all cultivars of plants listed on the label. *Arena* 0.25 G Insecticide has been tested on many cultivars with no phytotoxicity observed at label rates. Since all plant species and their varieties and cultivars have not been tested for tolerance, it is recommended that a small number of plants be treated to make certain that no phytotoxicity occurs, prior to any large scale application to plants. The end user assumes all risks arising from application of *Arena* 0.25 G Insecticide in a manner inconsistent with its labeling. *Arena* 0.25G can be blended just prior to application or co-applied with other crop management tools. However, all plant species and their varieties and cultivars have not been tested with possible blended or co-applied combinations or sequential pesticide treatments. The user must conduct a small scale compatibility and phytotoxicity trial under local conditions to ensure compatibility prior to any large-scale use.

COMMERCIAL AND RESIDENTIAL LANDSCAPE USES

Arena 0.25 G Insecticide is for use on landscape ornamentals, interior plantscapes and non-bearing fruit trees (including apple and pear) and nut trees (in landscapes).

Apply *Arena* 0.25 G Insecticide when threshold populations are observed. Do not wait until large insect populations have been established.

Apply to established plants. For control of pests on established plants, irrigate the treated area thoroughly.

Choose lower rates for light infestations and/or smaller plants and higher rates for heavy infestations and/or larger plants.

Application Methods:

Arena 0.25 G Insecticide is a systemic and soil active insecticide when applied to the soil.

Residential Landscapes and In-ground Plants: For flower beds, broadcast *Arena* 0.25 G Insecticide evenly over the top of plants. For treatment of individual shrubs or trees, broadcast *Arena* 0.25 G Insecticide evenly on soil around base of plant within 18 inches of the trunk. Pull back mulch prior to application if mulch is more than 1 inch thick. Irrigate after application with enough water to dissolve granules and move product into root zone.

RESTRICTIONS

- Do not apply more than a total of 160 lb *Arena* 0.25 G Insecticide (0.4 lb active ingredient clothianidin) per acre per season.

Pests Controlled by <i>Arena</i> 0.25 G Insecticide in Landscapes Ornamentals, Interior Plantscapes and Non-bearing Fruit Trees (Including Apple and Pear) and Nut Trees (in Landscapes)	
Adelgids including Hemlock Woolly Fungus Gnats (larvae) Glassy-Winged Sharpshooter Lacebugs including Azalea Leaf Beetles Leafhoppers Leafminers including Boxwood Serpentine Mealybugs including Citrus Long-tailed Madeira Obscure Pink Hibiscus Root	Psyllids including Asian Citrus Root Weevils (larvae and adult) including Black Vine Roundheaded Borers Scales (soft) *Thrips including Chilli Western Flower (suppression) Whiteflies including Giant Greenhouse Silverleaf Sweetpotato White Grubs including Oriental Beetle

*Not for use in California

SOIL APPLICATION TO INTERIORSCAPES, LANDSCAPES, NON-BEARING FRUIT AND NUT TREES		
For systemic insect control on ornamental plants grown in-ground and outdoor landscapes (commercial, industrial, recreational and residential)		
Crops	<i>Arena</i> 0.25 G Insecticide Rates	Special Instructions
Ornamental plants Including Shrubs Bedding Plants Flowering Plants Foliage Plants Ground Covers Evergreens Ornamental Trees Non-bearing Fruit Trees Non-bearing Nut Trees Non-bearing Vines	<p>Shrubs 0.5 to 1.0 lb/ft of height</p> <p>5 to 10 lb per 10 cumulative ft of height</p> <p>Trees 0.5 to 1.0 lb/inch of trunk diameter of breast height (DBH)</p> <p>For multi-stem trees, base rate on cumulative inches of diameter of all stems at breast height.</p>	<p>Only apply to moist soil media. Do not apply to dry or saturated soil.</p> <p>Do not apply to saturated or frozen soil, or when plants are not actively taking up water from soil.</p> <p>Heavy rainfall or inadequate irrigation immediately following application may decrease performance.</p> <p>For optimal control, apply early in the shrub or tree's annual growing cycle.</p>
	<p>Broadcast Application in Herbaceous Flower Beds and Ground Covers</p> <p>Apply 3.6 lb/1,000 sq ft of bed</p>	<p>Broadcast application may not be effective in trees and shrubs.</p>
<p><i>Arena</i> 0.25 G Insecticide is a systemic insecticide and must move to the area of the plant where the target pest is feeding. Control of pests may be delayed following an application for one or more weeks as <i>Arena</i> 0.25 G Insecticide moves to the feeding site. Factors such as plant size, soil type and water demands will impact the time between application and obvious insect control.</p> <ul style="list-style-type: none"> Do not apply more than a total of 160 lb <i>Arena</i> 0.25 G Insecticide (0.4 lb ai/A) per acre per season. Non-bearing fruit and nut trees in landscapes are only for ornamental use. Do not harvest fruit and nuts from treated trees within one year of application. This product may be blended just prior to application or co-applied with other compatible insecticides. 		

TURFGRASS

Arena 0.25 G Insecticide can be used as directed on grass grown in industrial, commercial and residential landscapes such as, airports, athletic fields, cemeteries, commercial sod farms, golf courses, home lawns, parks and playgrounds.

Arena 0.25 G Insecticide has sufficient residual activity to provide high levels of control when applications are made preceding or during the egg laying activity of the target pests. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Follow applications with sufficient irrigation or rainfall to move the active ingredient through the thatch.

Arena 0.25 G Insecticide can be applied to turf at 80 to 160 lb per acre. The rate is dependent on the target pest(s), their stage of development and the desired of control. Multiple applications can be made but do not exceed the maximum amount per year (160 lb/A). Consult your local State Agricultural Experiment Station, State Extension Turf Specialists or other turf experts for specific information concerning the timing of application(s). When targeting multiple pests, make application to earliest egg lay or adult flight. The high end of the rate range may be needed if extended control of secondary pests are desired.

Apply *Arena* 0.25 G Insecticide as a broadcast application to turfgrass for control of targeted pests. Use a calibrated spreader to uniformly broadcast *Arena* 0.25 G Insecticide over the treatment area. Do not mow turfgrass until after irrigation or rainfall has occurred to allow for the uptake and distribution of *Arena* 0.25 G Insecticide in soil and turfgrass

APPLICATION ON TURGRASS		
PESTS	ARENA 0.25 G INSECTICIDE APPLICATION RATES	TIMING INSTRUCTIONS
Annual Bluegrass Weevil (larval stages) Bill Bugs Black Turfgrass Ataenius (larval stages) <i>Phyllophaga</i> spp. (May or June Beetle) Spittlebug White Grubs including Asiatic Garden Beetle European Chafer Green June Beetle Japanese Beetle Northern Masked Chafer Oriental Beetle Southern Masked Chafer	80 to 160 lb/A (0.2 to 0.4 lb ai/A)	Preventative applications: Make applications through peak egg hatch of target species. Use the upper end of the rate range if application is made 60 days or more prior to peak adult flight and/or egg lay. Applications: For applications after egg hatch or when obvious turf damage has occurred use the upper end of the rate range.
Armyworms Chinch Bugs Crane Fly including American European Cutworms Nuisance Ants Sod webworms	120 to 160 lb/A (0.3 to 0.4 lb ai/A)	Cranefly: Apply in the spring, when larvae are mature but prior to pupation or in the fall prior to egg hatch. Other pests: Apply when pest first appear.
Mole Crickets (suppression)	160 lb/A (0.4 lb ai/A)	Time application to peak egg lay or early instars.
<ul style="list-style-type: none"> Do not apply more than a total of 160 lb of Arena 0.25 G Insecticide (0.4 lb ai/A) per acre per season. 		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in a dry place away from excessive heat.

Do not store near food or feed.

Store in original container only.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty the bag into granular application equipment. Then dispose of the empty bag by depositing in a sanitary landfill, or by other procedures allowed by state and local ordinances.

©2019 Valent U.S.A. LLC.

Arena is a registered trademark of Sumitomo Chemical Company, Ltd.

Manufactured for:
Valent U.S.A. LLC
P.O. Box 8025
Walnut Creek CA 94596-8025



Grow a better tomorrow.

Distributed by:
Nufarm Americas Inc.
11901 S. Austin Ave.,
Alsip, IL 60803

Made in U.S.A.

EPA Reg. No. 59639-157
EPA Est. No.

059639-00157.20190425.Arena_0.25G_AMEND.VPP.Clean