

59639-160

07/17/2012

1/42



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

OFFICE OF
PREVENTION PESTICIDES AND
TOXIC SUBSTANCES

JUL 17 2012

Robert L Hamilton, Ph D
Senior Regulatory Scientist
Registration & Regulatory Affairs
Valent U S A Corp
1101 14th Street, N W , Suite 150
Washington, DC 20005


Dear Dr Hamilton,

Subject Fast Track Label Amendments
Knack Insect Growth Regulator
EPA Reg No 59639-95
Esteem 35 WP Insect Growth Regulator
EPA Reg No 59639-115
Pyriproxyfen 0.86 EC Ag Insect Growth Regulator
EPA Reg No 59639-160
Submissions dated May 31, 2012

The label amendments above, submitted in connection with the Federal Insecticide, Rodenticide, and Fungicide Act, as amended, are acceptable. A stamped copy of each product's label is enclosed for your records.

Submit three (3) copies of your final printed labeling before releasing the product for shipment. Please also submit revised copies of each product's labeling to Andrew Ertman in support of the pending IR-4 tolerance petitions. Please contact Kevin Sweeney at (703) 305-5063 if you have any questions. Thank you for your cooperation in this matter.

Sincerely,


Richard Gebken

Product Manager 10


Insecticide Branch

Registration Division (7505P)



GROUP	7C	INSECTICIDE
-------	----	-------------

PYRIPROXYFEN 0.86 EC Ag
INSECT GROWTH REGULATOR



Active Ingredient	By Wt
*Pyriproxyfen	11.23%
Other Ingredients	88.77%
Total	100.00%

*(2-[1 methyl-2-(4-phenoxyphenoxy)ethoxy]pyridine)

Contains 0.86 pound ai per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT _____

...
...
...
...
...
...

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin Harmful if swallowed Avoid contact with skin eyes or clothing Causes moderate eye irritation Wash thoroughly with soap and water after handling and before eating drinking chewing gum or using tobacco Remove and wash contaminated clothing before reuse

FIRST AID

- | | |
|-------------------------------|--|
| If on skin or clothing | <ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice |
| If swallowed | <ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person |
| If in eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice |
| If inhaled | <ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth-to-mouth if possible • Call a poison control center or doctor for treatment advice |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact **800-892-0099** for emergency medical treatment information

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below If you want more options follow the instructions for selection category D on an EPA chemical resistance category selection chart

Applicators and other handlers must wear long-sleeved shirt and long pants socks shoes and chemical-resistant gloves (such as barrier laminate or butyl rubber ≥ 14 mils)

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables use detergent and hot water Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS

Users should

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing
- Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product s concentrate Do not reuse them

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by disposing of equipment washwaters or rinsate. Avoid direct application and/or spray drift to bee hives.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, chemical-resistant gloves such as Barrier Laminate or Butyl Rubber ≥ 14 mils, shoes plus socks.

**DISCLAIMER RISKS OF USING THIS PRODUCT
LIMITED WARRANTY AND LIMITATION OF LIABILITY**

IMPORTANT Read the entire Label including this Disclaimer Risks of Using this Product Limited Warranty and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT rather return the unopened product within 15 days of purchase for a refund of the purchase price

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as Buyer) of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label under average use conditions when used strictly in accordance with the label and **subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. **TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements, Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty one days from date of planting, or twenty one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law, if Buyer does not notify Valent of any claims in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product and Buyer accepts it, subject to the foregoing **Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability**, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent consistent with applicable law.

Read and follow the entire label of each product to be used in the tank mix with this product.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management purposes Pyriproxyfen 0.86 EC Ag Insect Growth Regulator is a Group 7C insecticide. Any insect population may contain individuals naturally resistant to Pyriproxyfen 0.86 EC Ag and other Group 7C insecticides. The resistant individuals dominate the insect population if these insecticides are used repeatedly. These resistant insects may not be controlled by Pyriproxyfen 0.86 EC Ag or other Group 7C insecticides, although local experts should be consulted for local resistance recommendations. The classification scheme is based on mode of action. It is recognized that resistance of insects and mites to insecticides and acaricides can also result from enhanced metabolism, reduced penetration or behavioral changes that are not linked to any site of action classification but are specific for individual chemicals or chemical groupings. Despite this, alternation of compounds from different chemical classes remains a viable management technique.

To delay insecticide resistance:

- Avoid exclusive repeated use of insecticides from the same chemical subgroup
- Integrate other control methods (chemical, cultural, biological) into insect control programs

For further information contact your local distributor.

GENERAL INFORMATION

- Do not apply this product through any type of irrigation system.

Pyriproxyfen Insect Growth Regulator is intended for use in Insect Pest Management (IPM) or Insect Resistance Management (IRM) programs. Pyriproxyfen will not control insect adults and it is recommended to be used in combination and/or rotation with other IPM or IRM materials. Contact your local state extension service for details.

Pyriproxyfen acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

ARTICHOKE (GLOBE)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Artichoke Globe	Artichoke Plum Moth Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	8 to 10 fl oz	Apply with ground equipment in adequate water for uniform coverage (10 to 50 gals/A)
			HIGHER WATER VOLUMES MAY PROVIDE IMPROVED INSECT CONTROL Begin applications when whiteflies reach threshold levels per University/Extension recommendations (e g 3 to 5 adults per leaf and/or equivalent levels of immatures) Under severe pressure use the higher recommended rates Only whole fields should be treated
Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs Do not apply within 7 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 20 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.134 lb pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses treated areas sooner than 30 days after the last application Do not apply this product with an adjuvant			

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides or insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 8 fl oz per acre for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

ASPARAGUS

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Asparagus	Whitefly	8 to 10 fl oz	<p>Apply with ground equipment in adequate water for uniform coverage (10 to 50 gals/A)</p> <p>HIGHER WATER VOLUMES MAY PROVIDE IMPROVED INSECT CONTROL</p> <p>Begin applications when whiteflies reach threshold levels per University/Extension recommendations (e.g. 3 to 5 adults per leaf and/or equivalent levels of immatures)</p> <p>Under severe pressure, use the higher recommended rates</p> <p>Only whole fields should be treated</p>
	<p>Do not apply within 7 days of harvest</p> <p>Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season</p> <p>Do not exceed 20 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season</p> <p>Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment</p> <p>Regardless of formulation, do not apply more than 0.134 lb pyriproxyfen per acre per season</p> <p>Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application</p> <p>Do not apply this product with an adjuvant</p>		

MANAGING WHITEFLY RESISTANCE Repeated use of the same class of insecticides or insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 8 fl oz per acre for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local state and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

BANANA AND PLANTAIN

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement, which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly, since one of the major obstacles is getting the product to the underside of the leaves where whiteflies are present.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Banana Plantain	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	13 to 16 fl oz	<p>Apply with air or ground equipment in adequate water for uniform coverage</p> <p>By Air 3 to 10 gals/A By Ground 10 to 50 gals/A</p> <p>HIGHER WATER VOLUMES MAY PROVIDE IMPROVED INSECT CONTROL</p> <p>Begin applications when whiteflies reach threshold levels per University/Extension recommendations (e.g. 3 to 5 adults per leaf and/or equivalent levels of immatures)</p> <p>Only whole fields should be treated</p>
	<p>Do not apply within 14 days of harvest</p> <p>Do not make more than 3 Pyriproxyfen 0.86 EC Ag applications per growing season</p> <p>Do not exceed 48 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season</p> <p>Regardless of formulation, do not apply more than 0.327 lb pyriproxyfen per acre per season</p> <p>Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment</p> <p>Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application</p>		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides or insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than three applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 13 fl oz per acre for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local, state, and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

BERRIES (Crop Group 13)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROPS	PESTS	PRODUCT RATE PER ACRE	SPECIAL INSTRUCTIONS
Berries Bingleberry Blackberry including Bingleberry Black Satin Berry Boysenberry Cherokee Blackberry Chesterberry Cheyenne Blackberry Coryberry Darrowberry Dewberry Dirksen thornless berry Himalayaberry Hullberry Lavacaberry Lowberry Lucretiaberry Mommoth Blackberry Marionberry Nectorberry Ollalieberry Oregon Evergreen Berry Phenomenalberry Rangeberry Ravenberry Rossberry Shawnee Blackberry Youngberry Blueberry Currant Elderberry Gooseberry Huckleberry Juneberry Lingonberry Loganberry Raspberry (black and red) Salal	Cherry Fruitworm Cranberry Fruitworm Lecanium Scale	16 fl oz	<p>Apply with ground or air equipment as a full coverage spray (minimum of 5 gals/A by air or 50 gals/A by ground). Thorough coverage is critical for adequate control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Fruitworms Apply when egg laying begins and again at petal fall. Additional applications of another insecticide at 10 – 14 day intervals may be needed under high population pressure or sustained moth flight.</p> <p>Lecanium Scale Apply at either dormant, delayed dormant or an in season cover spray. At dormant or delayed dormant use a spray oil at the manufacturer's recommended oil rate. In a cover spray make applications when scale crawlers first emerge. Follow local recommendations regarding in season use and rates of oil in your area.</p>
Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.			
Do not apply within 7 days of harvest. Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season. Do not exceed 32 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season. Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment. Regardless of formulation do not apply more than 0.218 lb pyriproxyfen per acre per season.			

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

BRASSICA LEAFY VEGETABLES (Crop Group 5)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement, which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly, since one of the major obstacles is getting the product to the underside of the leaves where whiteflies are present.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Brassica Leafy Vegetables Broccoli Broccoli Raab (rapini) Brussels Sprouts Cabbage Cauliflower Cavalo Broccolo Chinese Broccoli (gai lan) Chinese Cabbage (bok choy and napa) Chinese Mustard (gai choy) Collards Kale Kohlrabi Mizuna Mustard Greens Mustard Spinach Rape Greens Turnip Greens	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	8 to 10 fl oz	Apply with ground equipment in adequate water for uniform coverage (10 to 50 gals/A) HIGHER WATER VOLUMES MAY PROVIDE IMPROVED INSECT CONTROL Begin applications when whiteflies reach threshold levels per University/Extension recommendations (e.g. 3 to 5 adults per leaf and/or equivalent levels of immatures) Under severe pressure, use the higher recommended rates Only whole fields should be treated
	Do not apply within 7 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 20 fl oz Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation, do not apply more than 0.134 lb pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application		

MANAGING WHITEFLY RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 8 fl oz/A for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local, state, and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

BULB VEGETABLES (Crop Group 3)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Bulb Vegetables Garlic Garlic Great Headed Leek Onion Dry Bulb Onion Green Onion Welsh Shallot	Onion Thrips Western Flower Thrips	8 fl oz	FOR SUPPRESSION OF THRIPS Apply with ground equipment in adequate water for uniform coverage (20 to 50 gals/A). Begin applications when thrips reach threshold levels per University/Extension. Thorough coverage is critical for adequate water volume to ensure thorough coverage.
	Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs. <p>Do not apply within 3 days of harvest.</p> <p>Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season.</p> <p>Do not exceed 16 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season.</p> <p>Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment.</p> <p>Regardless of formulation, do not apply more than 0.134 lb pyriproxyfen per acre per season.</p> <p>Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application.</p>		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

CITRUS
(Including Ugli Fruit and White Sapote)
(Crop Group 10)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Citrus Ugli Fruit White Sapote	Black Scale Brown Soft Scale California Red Scale Florida Red Scale Snow Scale	16 fl oz	<p>Apply with ground equipment in 200 to 1500 gals/A. Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply anytime prior to fruit becoming infested when scale is in the crawler stage. Application at the 2nd or 3rd generation of scale may result in excessive dead scale remaining on the fruit at harvest. Thorough coverage is essential for good control.</p> <p>Pyriproxyfen 0.86 EC Ag may be applied with superior spray oils at 1/2 to 1%. Follow the manufacturer's directions and recommendations on timing and rates.</p> <p>NOTE: For scale, apply only one application during emergence of scale crawlers per calendar year.</p>
	<p>Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p> <p>Do not apply within 1 day of harvest. Do not make more than 1 Pyriproxyfen 0.86 EC Ag application per growing season. Do not exceed 16 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season. Regardless of formulation, do not apply more than 0.134 lb pyriproxyfen per acre per season.</p>		

continued

CITRUS (Including Ugli Fruit and White Sapote) (Crop Group 10) continued

Citrus Ugli Fruit White Sapote	Citrus Black Fly Citrus Leafminer Citrus Whitefly	10 fl oz	<p>Apply with ground application in 200 to 1500 gals/A. Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply when pests are just beginning to build and before infestation has occurred. Thorough coverage is essential for good control.</p>
	<p>Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p> <p>Do not apply within 1 day of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 20 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 21 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.134 lb pyriproxyfen per acre per season</p>		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

COTTON

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whiteflies are present.

CROP	PESTS	PRODUCT RATE PER ACRE	SPECIAL INSTRUCTIONS
Cotton	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	8 to 10 fl oz Do not apply less than 8 fl oz/A	Apply with air or ground equipment in adequate water for uniform coverage By Air 3 to 10 gals/A By Ground 10 to 50 gals/A HIGHER WATER VOLUMES MAY PROVIDE IMPROVED INSECT CONTROL Make only one application per growing season Begin application when whiteflies reach threshold levels per University/Extension recommendations (e.g. 3 to 5 adults per leaf and/or equivalent levels of immatures) Under severe pressure, use the higher recommended rates Only whole fields should be treated
	Do not apply within 28 days of harvest Do not make more than 1 Pyriproxyfen 0.86 EC Ag application per growing season Do not exceed 10 fl oz Pyriproxyfen 0.86 EC Ag per acre per season Regardless of formulation do not apply more than 0.067 lb pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after last application		

MANAGING WHITEFLY RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to one application per growing season. Pyriproxyfen 0.86 EC Ag Insect Growth Regulator may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 8 fl oz per acre for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local, state and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

CRANBERRY

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Cranberry	Cherry Fruitworm Cranberry Fruitworm Lecanium Scale	16 fl oz	<p>Apply with ground or air equipment as a full coverage spray (minimum of 5 gals/A by air or at least 20 gals/A by ground). Thorough coverage is critical for adequate control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Fruitworms Apply when egg laying begins and again at petal fall. Additional applications of another insecticide at 10 to 14 day intervals may be needed under high population pressure or sustained moth flight.</p> <p>Lecanium Scale Apply at either dormant, delayed dormant or an in season cover spray. At dormant or delayed dormant use a spray oil at the manufacturer's recommended oil rate. In a cover spray make applications when scale crawlers first emerge. Follow local recommendations regarding in season use and rates of oil in your area.</p>
	<p>Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p> <p>Do not apply within 7 days of harvest. Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season. Do not exceed 32 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season. Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment. Regardless of formulation do not apply more than 0.218 lb pyriproxyfen per acre per season.</p>		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

CUCURBITS (Crop Group 9)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Cucurbits Balsam Apple Balsam Pear Bitter Melon Cantaloupe Chayote Chinese Cucumber Chinese Waxgourd Citron Melon Cucumber Edible Gourd Gherkin <i>Momordica</i> spp Muskmelon Pumpkin Summer Squash Watermelon Winter Squash	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	8 to 10 fl oz Do not apply less than 8 fl oz/A	Apply with air or ground equipment in adequate water for uniform coverage By Air 3 to 10 gals/A By Ground 10 to 50 gals/A HIGHER WATER VOLUMES PROVIDE IMPROVED INSECT CONTROL Begin application when whiteflies reach threshold levels per University/Extension recommendations (e.g. 3 to 5 adults per leaf and/or equivalent levels of immatures) Under severe pressure, use the higher recommended rates Only whole fields should be treated
Do not apply within 7 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 20 fl oz Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation, do not apply more than 0.134 lb pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application			

MANAGING WHITEFLY RESISTANCE Repeated use of the same class of insecticides or insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 8 fl oz per acre for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local, state and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

FRUITING VEGETABLES (Crop Group 8)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Fruiting Vegetables Eggplant Ground Cherry Okra Pepino Pepper Tomatillo Tomato	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	8 to 10 fl oz Do not apply less than 8 fl oz/A of Pyriproxyfen when used as a stand alone application	Apply with air or ground equipment in adequate water for uniform coverage By Air 3 to 10 gals /A By Ground 10 to 50 gals/A HIGHER WATER VOLUMES PROVIDE IMPROVED INSECT CONTROL Begin application when whiteflies reach threshold levels per University/Extension recommendations (e.g. 3 to 5 adults per leaf and/or equivalent levels of immatures). Under severe pressure, use the higher recommended rates. Only whole fields should be treated.
	Banded Wing Whitefly Cabbage Looper Green Peach Aphid Greenhouse Whitefly (immatures & adults) Silverleaf Whitefly Sweetpotato Whitefly Tobacco Hornworm	6 to 8 fl oz	Apply as a tank mix with ground equipment for uniform coverage (25 to 150 gals/A by ground). Make application when eggs or insects first appear. If control is needed during this period, use an alternate insecticide registered for the target pest. Pyriproxyfen 0.86 EC Ag may also be mixed or alternated with commonly used insecticides to comply with local IPM programs. Comply with all applicable directions, restrictions, and precautions on the registered tank mix partner labels.
	Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs. Do not apply within 1 day of harvest. Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season. Do not exceed 16 fl oz Pyriproxyfen 0.86 EC Ag per acre per season. Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment. Regardless of formulation, do not apply more than 0.109 lb pyriproxyfen per acre per season. Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application.		

continued

FRUITING VEGETABLES (Crop Group 8) - continued

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Fruiting Vegetables Tomato	Banded Wing Whitefly Beet Armyworm Fruitworm Greenhouse Whitefly (immatures & adults) Potato Aphid Silverleaf Whitefly Sweetpotato Whitefly Thrips Tomato Pinworm Yellowstriped Armyworm	6 to 8 fl oz + Danitol ^o 2.4 EC Spray 10 2/3 fl oz	EARLY SEASON WHITEFLY CONTROL PROGRAM FOR REDUCTION OF GEMINIVIRUS Apply as a tank mix with ground equipment for uniform coverage (10 to 120 gals/A) Make application when scouting first indicates adult whiteflies are on young plants NOTE This may result in treatment immediately following transplanting MID TO LATE SEASON WHITEFLY CONTROL PROGRAM FOR PREVENTION OF IRREGULAR FRUIT RIPENING Apply as a tank mix with ground equipment for uniform coverage (20 to 120 gals/A by ground) Make application when scouting indicates whiteflies are present Pyriproxyfen 0.86 EC Ag may also be mixed or alternated with commonly used insecticides to comply with local IPM programs Comply with all applicable directions restrictions and precautions on the registered label for <i>Danitol 2.4 EC Spray</i>
	Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs Do not apply within 14 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 16 fl oz Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.109 lb pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

GRAPE

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an insect growth regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATE PER ACRE	SPECIAL INSTRUCTIONS
Grape	Glassy Winged Sharpshooter Grape Berry Moth Lecanium Scale	16 fl oz	<p>Apply with ground equipment as a full coverage spray (50 to 200 gals/A). Thorough coverage is critical for adequate control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Lecanium Scale Apply at either dormant, delayed dormant or an in season cover spray. At dormant, delayed dormant you may use a spray oil at the manufacturer's recommended oil rate. Do not use oil in an in season spray. In a cover spray, make applications when scale crawlers first emerge.</p> <p>Follow local recommendations regarding in season use and rates of oil in your area.</p>
			<p>Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p> <p>Do not apply within 21 days of harvest. Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season. Do not exceed 32 fl oz Pyriproxyfen 0.86 EC Ag per acre per season. Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment. Regardless of formulation, do not apply more than 0.218 lb active of pyriproxyfen per acre per season.</p>

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

LEAFY VEGETABLES (EXCEPT BRASSICA VEGETABLES) (Crop Group 4)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement, which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly, since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Leafy Vegetables (Except Brassica Vegetables) Amaranth (Chinese Spinach) Arugula (Rocket) Cardoon Celery Celery Chinese Celeriac Chervil Chrysanthemum Edible Leaved Chrysanthemum Garland Corn Salad Cress Garden Cress Upland Dandelion Dock (Sorrel) Endive (Escarole) Fennel Florence Lettuce Head and Leaf Orach Parsley Purslane Garden Purslane Winter Radicchio (Red Chicory) Rhubarb Spinach New Zealand Spinach Vine Spinach Swiss Chard	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	8 to 10 fl oz	Apply with ground equipment in adequate water for uniform coverage (10 to 50 gals/A) HIGHER WATER VOLUMES MAY PROVIDE IMPROVED INSECT CONTROL Begin applications when whiteflies reach threshold levels per University/Extension recommendations Under severe pressure, use the higher recommended rates Only whole fields should be treated
	<ul style="list-style-type: none"> Do not apply within 14 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 20 fl oz Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation, do not apply more than 0.134 lb pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application Do not apply this product with an adjuvant 		

MANAGING WHITEFLY RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 8 fl oz/A for whitefly control, because reduced rates can lead to a greater potential for resistance development. Follow local, state, and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

LEAVES OF ROOT & TUBER VEGETABLES (Crop Group 2)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Leaves of Root & Tuber Vegetables Beet Garden Burdock Edible Carrot Cassava Bitter Cassava Sweet Celeriac Chervil Turnip Rooted Chicory Dasheen (Taro) Parsnip Radish Radish Oriental (Daikon) Rutabaga Salsify Black Sugar Beet Sweet Potato Tanier Turnip Yam True	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	8 to 10 fl oz	Apply with ground equipment in adequate water for uniform coverage (10 to 50 gals/A) HIGHER WATER VOLUMES MAY PROVIDE IMPROVED INSECT CONTROL Begin applications when whiteflies reach threshold levels per University/Extension recommendations (e.g. 3 to 5 adults per leaf and/or equivalent levels of immatures) Under severe pressure use the higher recommended rates Only whole fields should be treated
	Do not apply within 7 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 16 fl oz Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.109 lb pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application Do not allow livestock to graze on treated areas Do not apply this product with an adjuvant		

MANAGING WHITEFLY RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 8 fl oz/A for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local, state and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

LEGUME VEGETABLES (Succulent or Dried)
(Crop Group 6)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

continued

LEGUME VEGETABLES (Crop Group 6) – continued

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Legume Vegetables (any cultivar of Bean (<i>Phaseolus</i>) Field Pea (<i>Pisum</i>) and Soybean) Beans (<i>Lupinus</i>) Grain Lupin Sweet Lupin White Lupin White Sweet Lupin Beans (<i>Phaseolus</i>) Field Bean Kidney Bean Lima Bean Navy Bean Pinto Bean Runner Bean Snap Bean Tepary Bean Wax Bean Bean (<i>Vigna</i>) Adzuki Bean Asparagus Bean Blackeyed Pea Catjang Chinese Longbean Cowpea Crowder Pea Moth Bean Mung Bean Rice Bean Southern Pea Urd Bean Yardlong Bean Broad Bean (fava) Chickpea (garbanzo) Guar Jackbean Lablab Bean Lentil Pea (<i>Pisum</i>) Dwarf Pea Edible podded Pea English Pea Field Pea Garden Pea Green Pea Snowpea Sugar Snap Pea Pigeon Pea Soybean Sword Bean	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	8 to 10 fl oz	EARLY SEASON WHITEFLY PROGRAM Apply with ground equipment in adequate water for uniform coverage (10 to 50 gals/A Thorough coverage is critical for adequate control It is essential to use adequate water volume to ensure thorough coverage Begin applications when whiteflies reach threshold levels per University/Extension recommendations (e g 3 to 5 adults per leaf and/or equivalent levels of immatures) Under severe pressure use the higher recommended rates Only whole fields should be treated
	Do not apply within 7 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season • Do not exceed 20 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.134 lb pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application Do not allow livestock to graze treated areas Do not use on legume vegetables varieties grown for livestock feed		

MANAGING WHITEFLY RESISTANCE Repeated use of the same class of insecticides or insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 8 fl oz per acre for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local, state and Federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

OLIVE

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Olive	Black Scale San Jose Scale	13 to 16 fl oz	<p>Apply with ground equipment in 100 to 400 gals/A. Apply in sufficient water volume for tree size. Thorough coverage is critical for adequate control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply at either dormant, delayed dormant, or an in-season cover spray. At dormant, delayed dormant, use a spray oil at the manufacturer's recommended oil rate. In a cover spray, make applications when scale crawlers first emerge. Follow local recommendations regarding in-season use and rates of oil in your area.</p> <p>Under heavy scale pressure, use the higher rate of Pyriproxyfen 0.86 EC Ag.</p>
	<p>Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p> <ul style="list-style-type: none"> Do not apply within 7 days of harvest. Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season. Do not exceed 32 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season. Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment. <p>Regardless of formulation, do not apply more than 0.218 lb pyriproxyfen per acre per season.</p>		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

PASTURE GRASS AND RANGELAND

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an insect growth regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Pasture Grass and Rangeland	Cricket Grasshopper Locust	8 to 10 fl oz	<p>Apply with air or ground equipment in adequate water for uniform coverage</p> <p>By Air 3 to 10 gals/A By Ground 10 to 50 gals/A</p> <p>HIGHER WATER VOLUMES PROVIDE IMPROVED INSECT CONTROL</p> <p>Begin application when pests reach threshold levels per University/Extension recommendations</p> <p>Under severe pressure, use the higher recommended rates</p>
	<p>Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Insect Pest Management (IPM) and Insect Resistance Management (IRM) programs</p> <p>Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season</p> <p>Do not exceed 20 fl oz Pyriproxyfen 0.86 EC Ag per acre per season</p> <p>Regardless of formulation, do not apply more than 0.134 lb pyriproxyfen per acre per season</p> <p>Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment</p>		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

PAWPAW

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Pawpaw	Scale Whitefly	13 to 16 fl oz	<p>Apply 100 to 400 gals/A with ground equipment Coverage is essential for good control Apply in sufficient water volume for tree size Use of higher water volume will assure better coverage</p> <p>Scale Apply at either dormant delayed dormant or an in season cover spray At dormant or delayed dormant use a spray oil at the manufacturer s recommended oil rate Do not use oil in an in season spray</p> <p>In a cover spray make applications when scale crawlers first emerge</p> <p>Under heavy scale pressure use the higher rate of Pyriproxyfen</p>
<p>In season sprays of Pyriproxyfen 0.86 EC Ag may not protect fruit from blemishes by scale</p> <p>Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs</p> <p>Do not apply within 14 days of harvest</p> <p>Do not make more than 3 Pyriproxyfen 0.86 EC Ag applications per growing season</p> <p>Do not exceed 48 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season</p> <p>Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment</p> <p>Regardless of formulation do not apply more than 0.327 lb pyriproxyfen per acre per season</p>			

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than three applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

PINEAPPLE

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Pineapple	Black Scale Brown Soft Scale California Red Scale Florida Red Scale Snow Scale	16 fl oz	<p>Apply with ground equipment in 20 to 50 gals/A. Apply in sufficient water volume for plant size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply anytime prior to fruit becoming infested when scale is in the crawler stage. Application at the 2nd or 3rd generation of scale may result in excessive dead scale remaining on the fruit at harvest. Thorough coverage is essential for good control.</p> <p>Pyriproxyfen 0.86 EC Ag may be applied with superior spray oils at 1/2 to 1%. Follow the manufacturer's directions and recommendations on timing and rates.</p>
	<p>Pyriproxyfen may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p> <p>Do not apply within 1 day of harvest. Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per calendar year. Do not exceed 16 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season. Do not apply earlier than 21 days after last Pyriproxyfen 0.86 EC Ag treatment. Regardless of formulation, do not apply more than 0.134 lb pyriproxyfen per acre per season.</p>		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per calendar year. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

POME FRUIT (Crop Group 11)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Pome Fruit Apple Crabapple Loquat Mayhaw Quince	Codling Moth Suppression of Pandemis Leafroller Obliquebanded Leafroller	13 to 16 fl oz	<p>Apply by ground with airblast equipment in a minimum of 50 gals/A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>1st Generation Codling Moth Apply at Biofix plus 100 degree days (this is usually close to petal fall). Make a second application 14 to 21 days later or timed to coincide with peak moth flights. For subsequent applications, if needed, apply other insecticides registered for Codling Moth control. The addition of oil has been shown to increase Codling Moth activity of Pyriproxyfen 0.86 EC Ag. Follow local recommendations regarding in season use and rates of oil in your area.</p> <p>Obliquebanded and Pandemis Leafroller Apply to the overwintering larvae at the pink to petal fall tree growth stage. Pyriproxyfen 0.86 EC Ag does not directly control the larvae but inhibits normal pupation and adult emergence.</p>
	Florida Red Scale Lecanium Scale San Jose Scale	13 to 16 fl oz	<p>Apply by ground with airblast equipment in a minimum of 50 gals/A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply at delayed dormant through pink. For delayed dormant use a superior spray oil at the manufacturer's recommended rate of oil. Under heavy scale pressure, use the higher rate of Pyriproxyfen 0.86 EC Ag.</p> <p>Scale For control in season, make application in cover sprays when scale crawlers first emerge. The use of oil at the crawler emergence stage has been shown to improve control. Follow local recommendations regarding in season use and rates of oil in your area.</p>

continued

POME FRUIT (Crop Group 11) - continued

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Pome Fruit Apple Crabapple Loquat Mayhaw Quince	Green Apple Aphid Rosy Apple Aphid Spirea Aphid Spotted Tentiform Leafminer Western Tentiform Leafminer	10 to 16 fl oz	<p>Apply by ground with airblast equipment in a minimum of 50 gals/A. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Leafminers Apply at peak moth flight for control of 1st or 2nd generation. This usually occurs at pre pink to pink apple stage for 1st generation and 2nd cover for 2nd generation.</p> <p>Rosy Apple Aphid Apply at 1/2 green apple growth stage.</p> <p>Green Apple & Spirea Aphid Apply when aphids first appear but before aphids are colonized.</p> <p>TIMING IS CRITICAL FOR ACCEPTABLE APHID CONTROL</p>
	Pear Psylla	13 to 16 fl oz	<p>Apply by ground with airblast equipment in 75 to 400 gals/A with a superior spray oil at manufacturer's recommended rates for the stage of growth. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply a single application during the delayed dormant to pink stage of growth at 16 fl oz/A or make two applications from delayed dormant through petal fall stage of growth at 13 to 16 fl oz/A. Make application just before sustained egg lay is anticipated.</p> <p>DO NOT USE SKIP ROWS DURING APPLICATION</p> <p>For in season control of pear psylla use an insecticide recommended for this use.</p>
	Lecanium Scale San Jose Scale	13 to 16 fl oz	<p>Apply by ground with airblast equipment in 100 to 400 gals/A at delayed dormant with a superior spray oil at the manufacturer's recommended rates. Under heavy scale pressure use the higher rate.</p> <p>Apply in sufficient water volume for tree size. Thorough coverage is critical for control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>For control of scale in season make application in cover sprays when scale crawlers first emerge. The use of oil at the crawler emergence stage has been shown to improve performance. Follow local recommendations regarding in season use and rates of oil in your area.</p>

continued

POME FRUIT (Crop Group 11) - continued

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Pome Fruit Pear Oriental Pear	Codling Moth	13 to 16 fl oz	Apply by ground with airblast equipment in 100 to 400 gals/A
	Suppression of Obliquebanded Leafroller Pandemis Leafroller		1st Generation Codling Moth Apply at Biofix plus 100 degree days Make a second application 14 to 21 days later or timed to coincide with peak moth flights For subsequent applications if needed apply other insecticides registered for Codling Moth control The addition of oil has been shown to increase Codling Moth activity of Pyriproxyfen 0.86 EC Ag Follow local recommendations regarding in season use and rates of oil in your area For additional generations apply other insecticides registered for Codling Moth Control
	Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs Do not apply within 45 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 32 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.218 lb pyriproxyfen per acre per season		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains To optimize resistance management practices Valent U S A Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator no more than two applications per growing season Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides Follow local state and federal IPM and IRM recommendations Read and follow all product labels before applying any insecticide

POMEGRANATE

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Pomegranate	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	13 to 16 fl oz	Apply with air or ground equipment in adequate water for uniform coverage By Air 3 to 10 gals/A By Ground 10 to 50 gals/A HIGHER WATER VOLUMES MAY PROVIDE IMPROVED INSECT CONTROL Begin applications when whiteflies reach threshold levels per University/Extension recommendations (e g 3 to 5 adults per leaf and/or equivalent levels of immatures) Only whole fields should be treated
	Do not apply within 14 days of harvest Do not make more than 3 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 48 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.327 lb pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides or insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than three applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 13 fl oz per acre for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local, state and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

ROOT AND TUBER VEGETABLES (Crop Group 1)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag does exhibit translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whiteflies are present.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Root & Tuber Vegetables Arracacha Arrowroot Artichoke Chinese Artichoke Jerusalem Beet Garden Beet Sugar Burdock Edible Canna Edible Carrot Cassava (bitter and sweet) Celeriac Chaoyote (root) Chervil Turnip rooted Chicory Chufa Dasheen (taro) Ginger Ginseng Horseradish Leren Parsley Turnip rooted Parsnip Potato Radish Radish Oriental (dai kon) Rutabaga Salsify Salsify Black Salsify Spanish Skirret Sweet Potato Tanier Turmeric Turnip Yam Bean Yam True	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	8 fl oz	Apply with ground equipment in adequate water for uniform coverage (20 to 50 gals/A) HIGHER WATER VOLUMES MAY PROVIDE IMPROVED INSECT CONTROL Begin applications when whiteflies reach threshold levels per University/Extension recommendations (e.g. 3 to 5 adults per leaf and/or equivalent levels of immatures) Under severe pressure use the higher recommended rates Only whole fields should be treated
	Do not apply within 7 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 16 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.109 lb pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application Do not allow livestock to graze treated areas		

continued

ROOT AND TUBER VEGETABLES (Crop Group 1) continued

MANAGING WHITEFLY RESISTANCE Repeated use of the same class of insecticides or insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 8 fl oz per acre for whitefly control because reduced rates can lead to a greater potential for resistance development. Follow local, state and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

SMALL FRUIT VINE CLIMBING CROPS (EXCEPT GRAPE) (Crop Group 13 07E)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROPS	PESTS	PRODUCT RATE PER ACRE	SPECIAL INSTRUCTIONS
Small Fruit Vine Climbing Crops Berry Schisandra Gooseberry Grape Amur River Kiwifruit Fuzzy Kiwifruit Hardy Maypop and cultivars, varieties and/or hybrids of these	Scale	16 fl oz	<p>Apply with ground or air equipment as a full coverage spray (minimum of 5 gals/A by air or 50 gals/A by ground). Thorough coverage is critical for adequate control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply at either dormant, delayed dormant, or an in-season cover spray. At dormant or delayed dormant, use a spray oil at the manufacturer's recommended oil rate. In a cover spray, make applications when scale crawlers first emerge. Follow local recommendations regarding in-season use and rates of oil in your area.</p>
	<p>Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.</p> <p>Do not apply within 30 days of harvest. Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season. Do not exceed 32 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season. Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment. Regardless of formulation, do not apply more than 0.218 lb pyriproxyfen per acre per season. Do not apply this product with an adjuvant.</p>		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

STONE FRUIT (Crop Group 12)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Stone Fruit Apricot Cherry Chickasaw Plum Damson Plum Fresh Prune Japanese Plum Nectarine Peach Plum Plumcot Prune	Lecanium Scale San Jose Scale Suppression of Peach Twig Borer Oriental Fruit Moth	13 to 16 fl oz	Apply in 100 to 400 gals/A with ground equipment Coverage is essential for good control Apply in sufficient water volume for tree size Use of higher water volume will assure better coverage Scale Apply at dormant delayed dormant or as an in season cover spray At dormant delayed dormant use a spray oil at the manufacturer s recommended oil rate Under heavy scale pressure use the higher rate of Pyriproxyfen Peach Twig Borer Pyriproxyfen 0.86 EC Ag has provided good suppression of Peach Twig Borer when applied delayed dormant through bloom sprays Use a spray oil at the manufacturer s recommended rate for the stage of crop growth
In season sprays of Pyriproxyfen 0.86 EC Ag may not protect fruit from blemishes by scale Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management IPM and Insect Resistance Management (IRM) programs Do not apply within 14 days of harvest Do not make more than 3 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 48 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.327 lb pyriproxyfen per acre per season			

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than three applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

STRAWBERRY

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an insect growth regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag exhibits translaminar movement which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly since one of the major obstacles is getting the product to the underside of the leaves where whitefly are present.

CROP	PESTS	PRODUCT RATE PER ACRE	SPECIAL INSTRUCTIONS
Strawberry	Banded Wing Whitefly Greenhouse Whitefly Silverleaf Whitefly Sweetpotato Whitefly	10 fl oz	Apply with ground equipment as a full coverage spray (100 to 400 gals/A) Thorough coverage is critical for adequate control It is essential to use adequate water volume to ensure thorough coverage Begin applications when whiteflies reach threshold levels per University/Extension recommendations (e g 3 to 5 adults per leaf and/or equivalent levels of immatures) Under severe pressure use the higher recommended rates Only whole fields should be treated
			Do not apply within 2 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 20 fl oz Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 30 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.134 lb of pyriproxyfen per acre per season Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application

MANAGING WHITEFLY RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

TREE NUTS (Crop Group 14)

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROP	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Tree Nuts Almond	Black Scale San Jose Scale Suppression of Peach Twig Borer	13 to 16 fl oz	Apply with ground equipment in 100 to 400 gals/A. Coverage is essential for good control. Apply in sufficient water volume for tree size. Use of higher water volume will assure better coverage. Apply at either dormant, delayed dormant or an in-season cover spray. At dormant, delayed dormant use a spray oil at the manufacturer's recommended oil rate. In a cover spray, make applications when scale crawlers first emerge. Follow local recommendations regarding in-season use and rates of oil in your area. Pyriproxyfen 0.86 EC Ag has provided good suppression of Peach Twig Borer when applied delayed dormant through bloom sprays.
	Almond Do not apply within 21 days of harvest Do not make more than 1 Pyriproxyfen 0.86 EC Ag application per growing season Do not exceed 16 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season Regardless of formulation do not apply more than 0.109 lb pyriproxyfen per acre per		

continued

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to one application per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state and federal Integrated Pest Management (IPM) and Insect Resistance Management (IRM) recommendations. Read and follow all product labels before applying any insecticide.

TREE NUTS (Crop Group 14) continued

CROPS	PESTS	PRODUCT RATES PER ACRE	SPECIAL INSTRUCTIONS
Tree Nuts Beechnut Brazil Butternut Cashew Chestnut Chinquapin Filbert (Hazelnut) Hickory Macadamia Pecan Pistachio Walnut	Codling Moth Dusky Veined Aphid Walnut Aphid	16 fl oz	Apply with ground equipment in 100 to 400 gals/A Apply in sufficient water volume for tree size Thorough coverage is critical for control It is essential to use adequate water volume to ensure thorough coverage 1st Generation Codling Moth Apply at Biofix plus 100 degree days Make a second application 14 to 21 days later or timed to coincide with peak moth flights For subsequent applications if needed apply other insecticides registered for Codling Moth control The addition of oil has been shown to increase Codling Moth activity of Pyriproxyfen Follow local recommendations regarding in season use and rates of oil in your area
	Black Scale Brown Soft Scale Italian Scale Lecanium Scale San Jose Scale Walnut Scale Suppression of Peach Twig Borer	13 to 16 fl oz	Apply with ground equipment in 100 to 400 gals/A at either dormant delayed dormant or an in season cover spray The addition of a superior spray oil at dormant or delayed dormant may improve scale control However do not use a spray oil in oil sensitive crops such as Walnut Follow manufacturer s recommendations regarding use and rates of oil for your area In a cover spray make applications when scale crawlers first emerge Follow local recommendations regarding in season use and rates of oil in your area Under heavy scale pressure use the higher rate Coverage is essential for good control Apply in sufficient water volume for tree size Use of higher water volume will assure better coverage
	Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs Do not apply within 21 days of harvest Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season Do not exceed 32 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment Regardless of formulation do not apply more than 0.218 lb pyriproxyfen per acre per season		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant insect strains To optimize resistance management practices Valent U S A Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator no more than two applications per growing season Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides Follow local state and federal IPM and IRM recommendations Read and follow all product labels before applying any insecticide

TROPICAL & SUBTROPICAL FRUIT

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an Insect Growth Regulator (IGR) and activity depends on the insect's development, evidence of activity will be slower than typical contact insecticides.

CROPS	PEST	PRODUCT RATE PER ACRE	SPECIAL INSTRUCTIONS
Tropical & Subtropical Fruit Acerola Atemoya Avocado Biriba Black Sapote Canistel Cherimoya Custard Apple Feijoa Guava Ilama Jaboticaba Longan Lychee Mamey Sapote Mango Passionfruit Papaya Pulasan Rambutan Sapodilla Soursop Spanish Lime Star Apple Starfruit Sugar Apple Wax Jambu	Scale	16 fl oz	<p>Apply with ground or air equipment as a full coverage spray (minimum of 5 gals/A by air or 50 gals/A by ground). Thorough coverage is critical for adequate control. It is essential to use adequate water volume to ensure thorough coverage.</p> <p>Apply at either dormant, delayed dormant, or an in-season cover spray. At dormant or delayed dormant, use a spray oil at the manufacturer's recommended oil rate. In a cover spray, make applications when scale crawlers first emerge. Follow local recommendations regarding in-season use and rates of oil in your area.</p>
	Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs.		
	<p>Do not apply within 14 days of harvest.</p> <p>Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season.</p> <p>Do not exceed 32 fl oz of Pyriproxyfen 0.86 EC Ag per acre per season.</p> <p>Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment.</p> <p>Regardless of formulation, do not apply more than 0.218 lb pyriproxyfen per acre per season.</p>		

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant Pyriproxyfen 0.86 EC Ag insect strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

WATERCRESS

GENERAL Pyriproxyfen 0.86 EC Ag Insect Growth Regulator acts by suppressing embryogenesis within the insect egg and by inhibiting metamorphosis and adult emergence of target insects. Pyriproxyfen 0.86 EC Ag has no activity on adult insects, but hatching of eggs laid by treated adults will be suppressed. Since Pyriproxyfen 0.86 EC Ag is an insect growth regulator (IGR) and activity depends on the insects' development, evidence of activity will be slower than typical contact insecticides.

Pyriproxyfen 0.86 EC Ag exhibits translaminar movement, which means that Pyriproxyfen 0.86 EC Ag applied to the upper surfaces of the leaves will move to the lower surface of the leaf. This translaminar movement is very important to chemical control of whitefly, since one of the major obstacles is getting the product to the underside of the leaves where whiteflies are present.

CROP	PESTS	PRODUCT RATE PER ACRE	SPECIAL INSTRUCTIONS
Watercress	Sharpshooters Whitefly	8 to 10 fl oz	<p>Apply with ground equipment as a full coverage spray (10 to 50 gals/A) Thorough coverage is critical for adequate control It is essential to use adequate water volume to ensure thorough coverage</p> <p>Begin applications when whiteflies reach threshold levels per University/Extension recommendations</p> <p>Under severe pressure use the higher recommended rates</p> <p>Only whole fields should be treated</p>
<p>Pyriproxyfen 0.86 EC Ag may be mixed and/or alternated with commonly used insecticides to comply with local Integrated Pest Management (IPM) and Insect Resistance Management (IRM) programs</p> <ul style="list-style-type: none">• Do not apply within 7 days of harvest• Do not make more than 2 Pyriproxyfen 0.86 EC Ag applications per growing season• Do not exceed 20 fl oz Pyriproxyfen 0.86 EC Ag per acre per season• Do not apply earlier than 14 days after last Pyriproxyfen 0.86 EC Ag treatment• Regardless of formulation do not apply more than 0.134 lb of pyriproxyfen per acre per season• Do not plant any crop other than those with registered pyriproxyfen uses in treated areas sooner than 30 days after the last application• Do not apply this product with an adjuvant			

MANAGING INSECT RESISTANCE Repeated use of the same class of insecticides with similar modes of action can lead to the buildup of resistant whitefly strains. To optimize resistance management practices, Valent U.S.A. Corporation is restricting Pyriproxyfen 0.86 EC Ag Insect Growth Regulator to no more than two applications per growing season. Pyriproxyfen 0.86 EC Ag may be used in alternation with other IGR materials possessing dissimilar modes of action and/or with other chemical classes of insecticides. Valent also does not recommend using rates below 8 fl oz per acre for whitefly control, because reduced rates can lead to a greater potential for resistance development. Follow local, state, and federal IPM and IRM recommendations. Read and follow all product labels before applying any insecticide.

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Store in a cool dry place

Keep pesticide in original container

Keep container closed when not in use

Do not put concentrate or dilute into food or drink containers

Not for use or storage in or around the home

Do not contaminate water food or feed by storage disposal or cleaning of equipment

For help with any spill leak fire or exposure involving this material call day or night 800 892 0099

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

CONTAINER DISPOSAL

Non refillable container Do not reuse or refill this container Triple rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

Copyright© 2012 by Valent U S A Corporation

Danitol is a registered trademark of Sumitomo Chemical Company Ltd

***Danitol* is a restricted use product**

Manufactured for

Valent U S A Corporation

P O Box 8025

Walnut Creek CA 94596 8025

www.valent.com

Made in U S A

EPA Reg No 59639 160

EPA Est _____

059639 00160 20120531 PYRI 0 86 EC Ag
20081204