



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

February 20, 2025

Ana Vida Reyes
anavida.reyes@valent.com
VALENT U.S.A. LLC

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment - To update label to current EPA standards and to comply with the Metconazole and Metalaxyl/Mefenoxam Interim Registration Review Decisions
Product Name: NIPSIT SUITE CANOLA SEED PROTECTANT
Admin Number: 59639-184
EPA Receipt Date: 10/22/2020 & 01/28/2022
Action Case Number: 00473645 & 00472233

Dear Ana Vida Reyes:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all of the information submitted with your application to support the Registration Review of the above referenced product in connection with the Metconazole and Metalaxyl/Mefenoxam Interim Decisions, and has concluded that your submissions are acceptable.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. The next label printing of this product must use this labeling unless subsequent changes have been approved. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process.

Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. If you have questions, please contact Ralph Narain by telephone at 202-566-2853 or via email at narain.ralph@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Ralph Narain for".

Loren LaPointe, Acting Product Manager 01
IVB3, RD
Office of Pesticide Programs

Enclosure: Stamped label



CLOTHIANIDIN	GROUP	4A	INSECTICIDE
METALAXYL	GROUP	4	FUNGICIDE
METCONAZOLE	GROUP	3	FUNGICIDE

NipsIt® SUITE Canola Seed Protectant

A SEED TREATMENT FORMULATION PROVIDING SYSTEMIC SEED AND SEEDLING PROTECTION AGAINST LISTED EARLY SEASON CANOLA DISEASES AND FLEA BEETLES

Active Ingredients	By Wt.
Clothianidin ¹	22.1500%
Metalaxyl ²	0.4150 %
Metconazole ³	0.0825%
Other Ingredients.....	77.3525%
Total	100.00%

¹(E)-1-[(2-chlorothiazol-5-yl)methyl]-3-methyl-2-nitroguanidine


² methyl N-(2,6-dimethylphenyl)-N-(methoxyacetyl)-DL-alaninate

³(1RS,5RS;1RS,5SR)-5-(4-chlorobenzyl)-2,2-dimethyl-1-(1H-1,2,4-triazol-1-ylmethyl) cyclopentanol

One gallon of NipsIt SUITE Canola Seed Protectant contains 2.325 lbs. clothianidin, 0.0436 lbs. metalaxyl, and 0.0087 lbs. metconazole.

KEEP OUT OF REACH OF CHILDREN

CAUTION



SEE NEXT PAGE FOR PRECAUTIONARY STATEMENTS

NET CONTENTS

EPA Reg. No. 59639-184
EPA Est. No.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing.

FIRST AID	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have a person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN	
Treatment is supportive and symptomatic.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 800-892-0099 for emergency medical treatment information.	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All mixers, loaders, applicators and other handlers must wear: Long-sleeved shirt and long pants, shoes plus socks and chemical-resistant gloves made out of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, Viton® ≥14 mils. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, and chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment wash water or rinsate. Do not apply where runoff is likely to occur. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Apply this product only as specified on label.

Treated seed exposed on soil surface may be hazardous to birds. Cover or collect spilled seeds.

Metalaxyl and Metconazole have properties and characteristics associated with chemicals detected in ground water. These chemicals may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore, contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

Surface Water Advisory

This product may impact surface water quality through spray and runoff of rain water. This product has a high potential for runoff for several months or more after application. Poorly draining soils or soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

For use only in commercial seed treatment equipment. Not for use in hopper box, planter box, slurry box, or other on-farm seed treatment applications. Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

The maximum application rate for mefenoxam and metalaxyl containing products (including seed treatments, foliar applications, soil applications) is 12.3 lbs. AI per acre per calendar year. **Do not** apply more than 12.3 lbs. mefenoxam and metalaxyl ai per acre per calendar year.

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

ASSUMPTION OF RISK

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests to this product. Such risks of crop injury, non-performance, or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, **THEN THIS PRODUCT SHOULD NOT BE APPLIED.** By applying this product Buyer acknowledges and accepts these inherent unintended risks **AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.**

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label **and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law AND AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability**, which may not be modified by any oral or written agreement.

COMMERCIAL TANK MIXES

NOTICE: Commercial tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the commercial tank mix with this product.

PRECAUTIONS AND RESTRICTIONS

NipsIt SUITE Canola Seed Protectant must be used only in accordance with specifications on this label utilizing commercial liquid or commercial slurry seed treatment equipment to secure proper application. Always mix product thoroughly before use. Not for use in hopper box, planter box, slurry box or other on-farm seed treatment applications.

Regardless of type of application (seed treatment or foliar) do not apply more than 0.2 lb (90.7 grams) of active ingredient clothianidin per acre per season.

Regardless of type of application, **do not** apply more than 12.3 lb mefenoxam and metalaxyl ai per acre per calendar year.

As stated by regulations specified in 40 CFR 153.155, all seed treated with *NipsIt* SUITE Canola Seed Protectant must be colored with an EPA-approved dye or colorant to visually distinguish seed that is treated with a pesticide. This product contains such dye or colorant. This regulation is necessary to prevent treated seed from accidental use as a food for man or feed for animals.

Seed Quality and Storage of Treated Seed

Seed treatments applied to low vigor or poor quality seed or to mechanically damaged seed may result in loss of seed germination or reduced seedling vigor. To verify seed safety, treat a small portion of seed with the desired slurry mixture (*NipsIt* SUITE Canola Seed Protectant alone or in combination with other commercial seed treatment products) prior to treatment of an entire seed lot. Utilize standard seed germination methodologies to determine and verify seed safety. Proper storage conditions of treated seed are recommended. Due to seed quality and seed storage conditions beyond the control of Valent U.S.A. LLC, no claims are made to guarantee the germination of carry-over treated seed.

PRODUCT INFORMATION

NipsIt SUITE Canola Seed Protectant is a systemic insecticide and fungicide seed protectant formulation for use on canola and rapeseed for early season control of flea beetle, seed-borne blackleg and the seedling disease complex (damping-off, seedling blight, seed rot, and root rot) caused by *Pythium spp.*, *Fusarium spp.* and *Rhizoctonia solani*. Clothianidin, the insecticide component within *NipsIt* SUITE Canola Seed Protectant, provides seedling protection against early season flea beetle feeding and damage. The fungicide components of *NipsIt* SUITE Canola Seed Protectant are metalaxyl and metconazole. Metalaxyl protects seed and seedlings against seed rot, damping-off, and seedling blight and root rot associated with *Pythium spp.* Metconazole fungicide provides protection against seed rot, damping-off, seedling blight and root rot associated with *Fusarium spp.* and *Rhizoctonia solani*, as well as seed-borne blackleg (*Leptosphaeria maculans*).

Accurate application of *NipsIt* SUITE Canola Seed Protectant to seed is required for control of listed pests. Application to seed by commercial seed treatment equipment utilizing standard liquid or slurry treaters, is necessary. Mix *NipsIt* SUITE Canola Seed Protectant prior to application. *NipsIt* SUITE Canola Seed Protectant may be applied direct to seed without adding water.

Pre-test tank mixtures with other seed treatment products to verify formulation compatibility and to assure proper physical compatibility of products. Be aware of limitations or cautions documented on labels of all products used in a seed treatment mixture.

MODES OF ACTION

Metalaxyl fungicide contained in *NipsIt* INSIDE Canola Seed Protectant is a member of the acylalanine chemical group (FRAC Code 4) and interrupts fungal nucleic acid synthesis. Metconazole belongs to the sterol biosynthesis inhibitor group of fungicides (FRAC Code 3). Clothianidin is a neonicotinoid insecticide (IRAC MoA Group 4A) acting as a nicotine acetylcholine receptor agonist/antagonist in insect metabolism.

RESISTANCE MANAGEMENT

For insecticide resistance management, please note that *NipsIt SUITE* Canola Seed Protectant contains a Group 4A insecticide: clothianidin. Any insect population may contain individuals naturally resistant to *NipsIt SUITE* Canola Seed Protectant and other Group 4A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

- To delay insecticide/acaricide resistance, take the following steps:
- Rotate the use of *NipsIt SUITE* Canola Seed Protectant or other Group 4A insecticides/acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides/acaricides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other
- chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For fungicide resistance management, please note that *NipsIt SUITE* Canola Seed Protectant contains a Group 3 fungicide/metconazole and a Group 4 fungicide/metalaxyl. Any fungal population may contain individuals naturally resistant to *NipsIt SUITE* Canola Seed Protectant and other Group 3, or Group 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay fungicide/bactericide resistance, take one or more of the following steps:

- Rotate the use of *NipsIt SUITE* Canola Seed Protectant or other Group 3 or 4, fungicides/bactericides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pathogens problems in your area.

For further information or to report suspected resistance contact Valent U.S.A. LLC at 800-6-VALENT (682-5368).

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with *NipsIt* SUITE Canola Seed Protectant must be labeled with the following statements:

- This seed has been treated with *NipsIt* SUITE Canola Seed Protectant which contains clothianidin, metalaxyl and metconazole.
- Do not use treated seed for feed, food or oil processing.

In addition, the U.S. Environmental Protection Agency requires the following statements on the container of seed treated with *NipsIt* SUITE Canola Seed Protectant:

- Regardless of the type of application (seed treatment or foliar), do not apply more than a total of 0.2 lb (90.7 gram) clothianidin (active ingredient) per acre per season. (Based on the seed size of 150,000 seeds/lb, at the label rate, the mg of clothianidin will be equal to 0.01208 mg /seed).
- Regardless of type of application, **do not** apply more than 12.3 lb mefenoxam and metalaxyl ai per acre per calendar year.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Store away from feeds and other foodstuffs.
- Wear long sleeved shirt, long pants and chemical- resistant made out of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, Viton® ≥14 mils when handling treated seed.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- The compound is toxic to birds and mammals. Treated clothianidin seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect seeds spilled during loading.
- Treated seeds must be planted at a minimum depth of 1 inch.
- This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- This product contains metalaxyl and metconazole that is known to leach through soil into groundwater under certain conditions as result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- Do not use rapeseed greens and seed grown or harvested from *NipsIt* Suite Canola Seed Protectant treated seed for feed or human consumption. Rapeseed grown and harvested from *NipsIt* Suite Canola Seed Protectant treated seed is only for industrial uses and cannot be used for edible oil or any human/feed consumption.
- Areas planted with treated seed of canola and rapeseed may be replanted immediately with bulb vegetables, canola, cereal grains (except rice), cotton, fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), rapeseed, root and tuber vegetables, soybean and sugar beet. These areas may also be replanted after 30 days with dry beans, grasses, non-grass animal feeds and rice. Do not plant any other crops in the treated area for at least 4 months after these treated seeds are planted.

- After seeds have been planted, do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 24 hours. Exception: Once seeds are planted and covered by soil or other planting media, the Worker Protection Standard (WPS) allows workers to enter the treated area without restriction.
- Do not re-use bags from treated seed to handle food or feed products.
- Do not use augers used for handling treated seed to move seed for feed, food, or oil processing.

APPLICATION DIRECTIONS

NipsIt SUITE Canola Seed Protectant is a ready-to-use formulation, which includes blue colorant. No additional colorant, dyes, binders, polymers, or water are required.

Calibrate seed treatment equipment prior to application of *NipsIt* SUITE Canola Seed Protectant to assure proper rate of application is achieved. Consult with the seed treatment equipment manufacturer for guidance on the operation and proper calibration of the treating equipment, as non-uniform, inaccurate seed coverage may result in suboptimal insect or disease protection to canola seedlings.

Apply 22 fl oz/100 lbs of seed of *NipsIt* SUITE Canola Seed Protectant using standard commercial seed treatment equipment to provide uniform seed coverage. Allow treated seed to dry prior to bagging.

The following table provides use rate information of *NipsIt* SUITE Canola Seed Protectant for disease and insect protection guidelines.

Canola and Rapeseed			
SEED/SEEDLING DISEASES	INSECT PESTS	NIPSIT SUITE CANOLA SEED PROTECTANT USE RATES	
		fl oz/100 lbs seed	¹ mg active ingredient per seed
Seed Borne Black Leg <i>(Leptosphaeria maculans)</i> Seedling Disease Complex (damping off and seedling blight, seed rot and root rot) <i>Pythium</i> spp. <i>Rhizoctonia</i> spp. <i>Fusarium</i> spp.	Early Season Flea Beetle feeding and damage	22.0	Clothianidin 0.01208 mg Metalaxyl 0.000227 mg Metconazole 0.0000452 mg
<ul style="list-style-type: none"> Regardless of type of application (seed treatment or foliar), do not apply more than a total of 0.2 lbs (90.7 gm) active ingredient clothianidin per acre per season.” Do not use rapeseed greens and seed grown or harvested from NipsIt Suite Canola Seed Protectant treated seed for feed or human consumption. Rapeseed grown and harvested from NipsIt Suite Canola Seed Protectant treated seed is only for industrial uses and cannot be used for edible oil or any human/feed consumption. Regardless of type of application, do not apply more than 12.3 lb mefenoxam and metalaxyl ai per acre per calendar year. 			

¹mg active ingredient per seed is based on 150,000 seeds per pound.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE

Store in a dry place away from excessive heat.

Do not store near food or feed.

Store in original container only.

To close package, replace and tighten cap to form an airtight seal.

PESTICIDE DISPOSAL

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or other procedures allowed by State and local authorities.

RECYCLING

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at **877-952-2272** (toll free) or www.acrecycle.org.

Copyright© 2020 by Valent U.S.A. LLC

NipsIt is a registered trademark of Valent U.S.A. LLC

Manufactured For:

Valent U.S.A. LLC

P.O. Box 5075

San Ramon CA 94583

(800)-682-5368

Made in U.S.A.

059639-00184.20241210.Metalaxyl.ID.Clean