



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

July 29, 2025

Ana Vida Reyes
Senior Regulatory Manager
Valent U.S.A. LLC
4600 Norris Canyon Road
San Ramon, CA 94583

Subject: Label Amendment - Registration Review Mitigation for Metalaxyl/Mefenoxam and Metconazole
Product Name: Metlock CT Fungicide
EPA Registration Number: 59639-191
Application Date: January 28, 2022 and June 28, 2022
Decision Number: 581548 and 585545
Case Number: 00476855 and 00478117

Dear Ana Vida Reyes:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Metalaxyl/Mefenoxam and Metconazole Interim Decisions, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


If you have any questions about this letter, please contact Concepción Rodríguez by phone at 202-566-0820, or via email at rodriguez.concepcion@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'M. K. Muhammad-Perch', with a long, sweeping horizontal line extending to the right.

Maryam K. Muhammad-Perch, Team Lead
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label



| | | | |
|-------------|-------|---|-----------|
| METCONAZOLE | GROUP | 3 | FUNGICIDE |
| METALAXYL | GROUP | 4 | FUNGICIDE |

METLOCK™ CT Fungicide

A FUNGICIDE SEED TREATMENT PRODUCT PROVIDING SYSTEMIC SEED AND SEEDLING PROTECTION AGAINST LISTED EARLY SEASON SEEDLING DISEASES

| Active Ingredient | By Wt. |
|--------------------------------|---------|
| Metalaxyl ¹ | 4.51% |
| Metconazole ² | 2.25% |
| Other Ingredients | 93.24 % |
| Total | 100.00% |

¹N-(2,6-dimethylphenyl)-N-(methoxyacetyl)alanine methyl ester

²5-[(4-chlorophenyl)methyl]-2,2-dimethyl-1-(1*H*-1,2,4-triazol-1-ylmethyl)cyclopentanol

METLOCK™ CT Fungicide contains 0.387 lbs metalaxyl and 0.193 lbs metconazole per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE NEXT PAGE FOR PRECAUTIONARY STATEMENTS AND FIRST AID.

NET CONTENTS

EPA Reg. No. 59639-191
EPA Est. No.

| FIRST AID | |
|--|---|
| If swallowed: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person. |
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice. |
| NOTE TO PHYSICIAN No specific antidote. Supportive care. Treatment based on judgment of the physician in response to reactions of the patient. | |
| HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 800-892-0099 for emergency medical treatment information. | |

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: Long-sleeved shirt and long pants, socks and shoes, and chemical resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride or Viton ≥ 14 mils.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated area.

Metaxyl and Metconazole have properties and characteristics associated with chemicals detected in ground water. These chemicals may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

Metconazole may impact surface water quality through runoff of rainwater. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Surface Water Advisory

This product may impact surface water quality through spray and runoff of rainwater. This product has a high potential for runoff for several months or more after application. Poorly draining soils or soils with shallow water tables are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: Once the seeds are planted in soil, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area without restriction if there will be no worker contact with the soil subsurface or treated seed.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: long-sleeved shirt and long pants; shoes plus socks and chemical-resistant gloves ; made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride or Viton \geq 14 mils.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: Do not apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests, resistance of the target pest to this product. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN DO NOT APPLY PRODUCT. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND AGREES THAT TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

To the extent consistent with applicable law, Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. To the extent consistent with applicable law, Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, EXCEPT AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the extent consistent with applicable law, Valent or Seller shall not be liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law, Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law, if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

APPLICATION INSTRUCTIONS

Read entire label. Use strictly in accordance with precautionary statements and directions and with applicable state and federal regulations.

RESTRICTIONS

METLOCK CT Fungicide must be used only in accordance with use instructions on this label utilizing mechanical, slurry or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. This product is for commercial application only. Always mix product thoroughly before use.

As stated by regulations specified in 40 CFR 153.155, all seed treated with *METLOCK CT* Fungicide must be colored with an EPA-approved dye or colorant to visually distinguish seed is treated with a pesticide. It is mandatory that such dye or colorant be added into the slurry mixture containing *METLOCK CT* Fungicide, prior to application. This regulation is necessary to prevent treated seed from accidental use as a food for man or feed for animals.

Seed Quality and Storage of Treated Seed

Seed treatments applied to low vigor or poor quality seed or to mechanically damaged seed may result in loss of seed germination or reduced seedling vigor. To verify seed safety, treat a small portion of seed with the desired slurry mixture (*METLOCK CT* Fungicide alone or in combination with other commercial seed treatment products) prior to treatment of an entire seed lot. Utilize standard seed germination methodologies to determine and verify seed safety. Proper storage conditions of treated seed are recommended. Due to seed quality and seed storage conditions beyond the control of Valent U.S.A. LLC, no claims are made to guarantee the germination of carry-over treated seed.

Seed Bag Label Requirement of Treated Seed

The Federal Seed Act requires that bags containing treated seed be labeled or tagged with the following information:

- This seed has been treated with *METLOCK CT* Fungicide containing metconazole and metalaxyl.
- Do not use for feed, food or oil processing.
- Store away from feeds and other foodstuffs.
- Wear long sleeved shirt, long pants and chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride or Viton ≥ 14 mils when handling treated seed.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.

- No plant back restriction for Barley, Canola, Corn (including field, sweet and pop) Oat, Peanut, Rye, Soybean, Sugar Beet, Triticale and Wheat. A 30-day plant back interval for Brassica Leafy Vegetables and Leafy Vegetables. A 120-day plant back interval for all other crops.

USE INFORMATION

METLOCK CT Fungicide is a systemic fungicide formulated for seed treatment use to provide protection against seed-borne and soil-borne diseases, enhancing plant health and plant stand establishment of seed/seedlings. The fungicide active ingredients in *METLOCK* CT Fungicide are metconazole, a broad-spectrum, systemic, triazole fungicide and metalaxyl, an oomycete-active, systemic acylalanine fungicide. The fungicide components provide activity against *Pythium* spp., *Rhizoctonia solani*, *Fusarium* spp. and other Deuteromycete and Basidiomycete (smut and bunt) fungi.

MODE OF ACTION

Metalaxyl fungicide contained in *METLOCK* CT Fungicide is a member of the acylalanine chemical group (FRAC Code 4) and interrupts fungal nucleic acid synthesis. Metconazole belongs to the sterol biosynthesis inhibitor group of fungicides (FRAC Code 3).

RESISTANCE MANAGEMENT

For fungicide resistance management, please note that *METLOCK* CT Fungicide contains a Group 3 fungicide/metconazole and a Group 4 fungicide/metalaxyl. Any fungal population may contain individuals naturally resistant to *METLOCK* CT Fungicide and other Group 3 or Group 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of *METLOCK* CT Fungicide or other Group 3 or 4 fungicides within a growing season sequence with different groups that control the same pathogens.
- Avoid application of more than maximum number of applications per year as listed in the Crop Specific Directions and maximum number of consecutive sprays of *METLOCK* CT Fungicide or other fungicides in the same group in a season
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pathogens problems in your area.

For further information or to report suspected resistance contact Valent U.S.A. LLC at 800-6-VALENT (682-5368). You can also contact your pesticide distributor or university extension specialist to report resistance.

APPLICATION DIRECTIONS

Accurate application of *METLOCK* CT Fungicide to seed is required for best performance. Application to seed by commercial seed treatment equipment utilizing standard liquid or slurry treaters, is necessary. Thoroughly mix *METLOCK* CT Fungicide in the desired seed slurry mixture, prior to application to seed.

Always pre-test tank mixtures with other seed treatment products to evaluate formulation compatibility and to ensure proper physical compatibility of products. Follow more restrictive limitations or cautions documented on labels of all products used in a seed treatment mixture. Do not tank mix with any products which contain a prohibition on tank mixing.

ROTATIONAL RESTRICTIONS

- No plant back interval for Barley, Canola, Corn (including field, sweet and pop), Oat, Peanut, Rye, Soybean, Sugar Beet, Triticale and Wheat.
- A 30-day plant back interval for Brassica Leafy Vegetables and Leafy Vegetables.
- A 120-day plant back interval for all other crops.

The following tables provide use rates of *METLOCK* CT Fungicide by labeled crop, with disease protection guidelines:

Table 1.

| BARLEY | | | |
|--|--|--|---|
| Seed/Seedling Diseases | <i>METLOCK</i> CT Fungicide Use Rates | | |
| | fl oz/100 lbs seed | Active Ingredient (mg ai/seed)³ | Total Active Ingredient (mg ai/seed)³ |
| Covered Smut Early Season Common Root Rot Early Season Fusarium Seedling Dieback Early Season Rhizoctonia Root Rot Fusarium Seed Scab Loose Smut Pythium Seed Rot and Seedling Dieback Seed Decay Fungi | 1.0–1.5 | Metalaxyl (0.001-0.0015) Metconazole (0.0005-0.00075) | 0.0015–0.00225 |

³mg active ingredient per seed is based on 13,605 seeds per pound.

Table 2.

| CORN (FIELD, POP, SWEET, includes seed production) | | | | |
|--|--|----------------------------------|--|---|
| Seed/Seedling Diseases | <i>METLOCK</i> CT Fungicide Use Rates | | | |
| | fl oz/100 lbs seed | fl oz/80,000 Kernal Count | Active Ingredient (mg ai/seed)⁴ | Total Active Ingredient (mg ai/seed)⁴ |
| Fusarium Seed/Seedling Dieback Pythium Seed Rot and Seedling Dieback Rhizoctonia Damping-off Seed Decay Fungi | 1.0–1.5 | 0.49-0.74 | Metalaxyl (0.0084-0.0126) Metconazole (0.0042-0.0063) | 0.0126–0.0189 |

⁴calculation is based on 1,630 seeds per pound.

Table 3.

| COTTON | | | |
|---|--|--|---|
| Seed/Seedling Diseases | <i>METLOCK</i> CT Fungicide Use Rates | | |
| | fl oz/100 lbs seed | Active Ingredient (mg ai/seed)⁵ | Total Active Ingredient (mg ai/seed)⁵ |
| Fusarium Seed/seedling Dieback Pythium Seed Rot and Seedling Decay Rhizoctonia Damping-off Seed Decay Fungi Thielaviopsis Black Root Rot | 5.0–6.5 | Metalaxyl (0.015235-0.0198) Metconazole (0.007598-0.009877) | 0.022833-0.029677 |

⁵mg active ingredient per seed is based on 4,500 seeds per pound.

Table 4.

| OATS | | | |
|--|--|--|---|
| Seed/Seedling Diseases | <i>METLOCK</i> CT Fungicide Use Rates | | |
| | fl oz/100 lbs seed | Active Ingredient (mg ai/seed)⁶ | Total Active Ingredient (mg ai/seed)⁶ |
| Early Season Common Root Rot Early Season Fusarium Seedling Dieback Early Season Rhizoctonia Root Rot Fusarium Seed Scab Loose Smut Pythium Seed Rot and Seedling Dieback Seed Decay Fungi | 1.0–1.5 | Metalaxyl (0.00072-0.00107) Metconazole (0.00036-0.00054) | 0.00108–0.00161 |

⁶mg active ingredient per seed is based on 19,047 seeds per pound

Table 5.

| WHEAT | | | |
|--|--|--|---|
| Seed/Seedling Diseases | <i>METLOCK CT</i> Fungicide Use Rates | | |
| | fl oz/100 lbs seed | Active Ingredient (mg ai/seed)⁷ | Total Active Ingredient (mg ai/seed)⁷ |
| Common Bunt Early Season Common Root Rot Early Season Fusarium Seedling Dieback Early Season Rhizoctonia Root Rot Early Season Leaf and Stripe Rust (Suppression) Flag Smut Fusarium Seed Scab Loose Smut Pythium Seed Rot and Seedling Dieback Seed Decay Fungi | 1.0–1.5 | Metalaxyl (0.0012-0.0018) Metconazole (0.0006-0.0009) | 0.0018–0.0027 |

⁷mg active ingredient per seed is based on 11,338 seeds per pound.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE

Store in a dry place away from excessive heat.

Do not store near food or feed.

Store in original container only.

To close package, replace and tighten cap to form an airtight seal.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

[Use the following statement for containers equal to or less than 5 gallons]

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

[Use the following statement for containers greater than 5 gallons]

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or incinerator.

[Use the following statement for all formulation types/all refillable container types]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or incineration.

©2025 Valent U.S.A. LLC

METLOCK is a trademark of Valent U.S.A. LLC

Manufactured For:

Valent U.S.A. LLC

P. O. Box 5075

San Ramon CA 94583-0975

(800)-682-5368

Made in U.S.A.

EPA Reg. No. 59639-191

EPA Est. No.

059639-00191.20250618.METLOCKCT_AMEND_Metalaxyl_ID_Clean