



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

November 07, 2024

Robert Hamilton  
robert.hamilton@valent.com  
VALENT U.S.A. LLC

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment - Add an old supplemental label and other minor revisions  
Product Name: CERTAINTY TURF HERBICIDE  
Admin Number: 59639-226  
EPA Receipt Date: 05/16/2023  
Action Case Number: 00480623

Dear Robert Hamilton:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

The label submitted with the application has been stamped "Accepted Only Indicated Revisions Reviewed" and is enclosed for your records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have questions, please contact Francisco Llarena-Arias via email at [llarena-arias.francisco@epa.gov](mailto:llarena-arias.francisco@epa.gov).

Sincerely,

A handwritten signature in black ink, reading "Francisco Llarena-Arias". The script is elegant and cursive, with the first letters of each word being capitalized and prominent.

Francisco Llarena-Arias, Environmental Protection Specialist  
FHB, RD  
Office of Pesticide Programs

# MASTER LABEL FOR EPA REG. NO. 59639-226

SULFOSULFURON GROUP 2 HERBICIDE

**REDLINE COPY**

## Certainty® Turf Herbicide

***Bold italicized text is information for the reader and is not part of the label.***

[Bracketed Text is Optional]

**ACCEPTED**

ONLY INDICATED  
REVISIONS REVIEWED

11/07/2024

Under the Federal Insecticide, Fungicide and  
Rodenticide Act as amended, for the pesticide  
registered under EPA Reg. No.

59639-226

No label revisions other than those indicated were  
reported to the Agency.

### PRODUCT LOGO

**Certainty Turf Herbicide is a selective herbicide for control of annual and perennial grass and broadleaf weeds in highly managed turf sites.**

### Complete Directions For Use

Read the entire label before using this product.

Use only according to label instructions.

Not all products recommended on this label are registered for use in California. Check the registration status of each product in California before using.

Read the "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

[THIS IS AN END-USE PRODUCT. VALENT U.S.A. LLC DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING.]

### INGREDIENTS

ACTIVE INGREDIENT:	By Wt
Sulfosulfuron*, .....	75.0%
OTHER INGREDIENTS: .....	25.0%
	100.0%

\* N-[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl-2-(ethylsulfonyl)imidazo[1,2-a]pyridine-3-sulfonamide N-[(4,6-dimethoxypyrimidin-2-yl)carbamoyl]-2-(ethylsulfonyl)imidazo[1,2-a]pyridine-3-sulfonamide

**SEE [NEXT] [PAGE] [PANEL] [BOOKLET] FOR [PRECAUTIONARY STATEMENTS]  
[AND] [DIRECTIONS FOR USE]**

EPA Reg. No. 59639-226

**Certainty Turf Herbicide** is formulated as a water dispersible granule (WDG).

## **TABLE OF CONTENTS**

<b>INGREDIENTS.....</b>	<b>1</b>
<b>IMPORTANT PHONE NUMBERS .....</b>	<b>77</b>
<b>PRECAUTIONARY STATEMENTS.....</b>	<b>77</b>
Hazards to Humans and Domestic Animals.....	77
Environmental Hazards .....	77
Personal Protective Equipment (PPE) .....	88
<b>DIRECTIONS FOR USE .....</b>	<b>99</b>
<b>STORAGE AND DISPOSAL .....</b>	<b>1144</b>
<b>PRODUCT INFORMATION .....</b>	<b>1144</b>
<b>Weed Resistance Management.....</b>	<b>1313</b>
<b>MIXING .....</b>	<b>1444</b>
Mixing with Water .....	1444
Mixing for Hand-Held Sprayers.....	1444
Tank-Mixing Procedure .....	1545
Surfactants and Adjuvants.....	1545
Colorants and Dyes .....	1545
<b>APPLICATION EQUIPMENT AND TECHNIQUES .....</b>	<b>1545</b>
Ground Broadcast Equipment .....	1545
Hand-Held Sprayers.....	1747
Equipment Cleaning .....	1747
<b>MANDATORY SPRAY DRIFT MANAGEMENT .....</b>	<b>1848</b>
SPRAY DRIFT ADVISORIES .....	1848
IMPORTANCE OF DROPLET SIZE .....	1848
BOOM HEIGHT.....	1949
SHIELDED SPRAYERS .....	1949
TEMPERATURE AND HUMIDITY.....	1949
TEMPERATURE INVERSIONS .....	1949
WIND .....	1949
<b>WARM-SEASON TURFGRASSES .....</b>	<b>2020</b>
Sedge Control .....	2124
Tall Fescue Control .....	2124
Dallisgrass Suppression in Bermudagrass .....	2222
Virginia Buttonweed Suppression.....	2222
Annual Bluegrass ( <i>Poa annua</i> ) Control in Non-Overseeded Turf.....	2222

Annual Bluegrass ( <i>Poa annua</i> ) Control Prior to Overseeding Turf with Perennial Ryegrass.....	2222
Removal of Overseeded Perennial Ryegrass (Transition Aid) .....	2424
Additional Weeds Controlled .....	2525
<b>COOL-SEASON TURFGRASSES.....</b>	<b>2626</b>
Weed Control in Established Creeping Bentgrass .....	2626
Roughstalk Bluegrass ( <i>Poa trivialis</i> ) Control.....	2626
Weed Control in Kentucky Bluegrass .....	2727
Sedge Control .....	2727
Roughstalk Bluegrass ( <i>Poa trivialis</i> ) Control.....	2727
Tall Fescue Control .....	2727
Additional Weeds Controlled .....	2828
Weed Control in Mixed Stands of Kentucky Bluegrass .....	2828
Sedge Suppression .....	2929
Roughstalk Bluegrass ( <i>Poa trivialis</i> ) Control.....	2929
Additional Weeds Controlled .....	3030
<b>ORNAMENTALS .....</b>	<b>3131</b>
Directed-Spray Applications .....	3232
Over-the-Top Applications .....	3232
Pre-Plant Applications .....	3232
<b>NATIVE GRASSES.....</b>	<b>3434</b>
Sedge Control .....	3434
Tall Fescue Control .....	3434
Additional Weeds Controlled .....	3535
<b>RISKS OF USING THIS PRODUCT, LIMITED WARRANTY AND DISCLAIMER, AND LIMITATION OF LIABILITY.....</b>	<b>3737</b>
<b>INGREDIENTS .....</b>	
<b>IMPORTANT PHONE NUMBERS .....</b>	
<b>PRECAUTIONARY STATEMENTS.....</b>	
Hazards to Humans and Domestic Animals.....	
Environmental Hazards .....	
<b>STORAGE AND DISPOSAL .....</b>	
<b>PRODUCT INFORMATION .....</b>	
<b>MIXING .....</b>	
Mixing with Water.....	
Mixing for Hand-Held Sprayers .....	

<b>INGREDIENTS</b>	<b>1</b>
<b>IMPORTANT PHONE NUMBERS</b>	<b>4</b>
<b>PRECAUTIONARY STATEMENTS</b>	<b>4</b>
Hazards to Humans and Domestic Animals	4
Environmental Hazards	4
Personal Protective Equipment (PPE)	5
<b>DIRECTIONS FOR USE</b>	<b>6</b>
NON-TARGET ORGANISM ADVISORY STATEMENT	6
WINDBLOWN SOIL PARTICLES	6
<b>STORAGE AND DISPOSAL</b>	<b>8</b>
<b>PRODUCT INFORMATION</b>	<b>8</b>
<b>Weed Resistance Management</b>	<b>10</b>
<b>MIXING 11</b>	
Mixing with Water	11
Mixing for Hand-Held Sprayers	11
Tank-Mixing Procedure	12
Surfactants and Adjuvants	12
Colorants and Dyes	12
<b>APPLICATION EQUIPMENT AND TECHNIQUES</b>	<b>12</b>
Ground Broadcast Equipment	12
Hand-Held Sprayers	14
Equipment Cleaning	14
<b>MANDATORY SPRAY DRIFT MANAGEMENT</b>	<b>15</b>
SPRAY DRIFT ADVISORIES	15
IMPORTANCE OF DROPLET SIZE	15
BOOM HEIGHT	16
SHIELDED SPRAYERS	16
TEMPERATURE AND HUMIDITY	16
TEMPERATURE INVERSIONS	16
WIND	16
<b>WARM-SEASON TURFGRASSES</b>	<b>17</b>
Sedge Control	18
Tall Fescue Control	18
Dallisgrass Suppression in Bermudagrass	19
Buttonweed Suppression	19

Annual Bluegrass (Poa annua) Control in Non-Overseeded Turf.....	19
Annual Bluegrass (Poa annua) Control Prior to Overseeding Turf with Perennial Ryegrass.....	19
Removal of Overseeded Perennial Ryegrass (Transition Aid) .....	21
Additional Weeds Controlled .....	21
<b>COOL-SEASON TURFGRASSES.....</b>	<b>23</b>
Weed Control in Established Creeping Bentgrass .....	23
Roughstalk Bluegrass (Poa trivialis) Control.....	23
Weed Control in Kentucky Bluegrass .....	24
Sedge Control .....	24
Roughstalk Bluegrass (Poa trivialis) Control.....	24
Tall Fescue Control .....	24
Additional Weeds Controlled .....	25
<b>1 Suppression or partial control only.....</b>	<b>25</b>
Weed Control in Mixed Stands of Kentucky Bluegrass .....	25
Sedge Suppression .....	26
Roughstalk Bluegrass (Poa trivialis) Control.....	26
Additional Weeds Controlled .....	27
<b>ORNAMENTALS .....</b>	<b>28</b>
Directed Spray Applications .....	29
Over the Top Applications .....	29
Pre-Plant Applications .....	29
<b>NATIVE GRASSES.....</b>	<b>31</b>
Sedge Control .....	31
Tall Fescue Control .....	31
Additional Weeds Controlled .....	32
<b>1 Suppression or partial control only].....</b>	<b>33</b>
<b>RISKS OF USING THIS PRODUCT, LIMITED WARRANTY AND DISCLAIMER, AND LIMITATION OF LIABILITY.....</b>	<b>34</b>
<b>Certainty is a registered trademark of Valent U.S.A. LLC.....</b>	<b>35</b>
<b>©2022 Valent U.S.A. LLC .....</b>	<b>35</b>
e.....	
Surfactants and Adjuvants .....	
Colorants and Dyes .....	
<b>APPLICATION EQUIPMENT AND TECHNIQUES.....</b>	
Mandatory Spray Drift Management.....	
Ground Broadcast Equipment.....	
Hand Held Sprayers .....	
Equipment Cleaning.....	

**WARM SEASON TURFGRASSES**

Sedge Control

Tall Fescue Control

Dallisgrass Suppression in Bermudagrass

Buttonweed Suppression

Annual Bluegrass (*Poa annua*) Control in Non-Overseeded Turf

Annual Bluegrass (*Poa annua*) Control Prior to Overseeding Turf with Perennial Ryegrass

Removal of Overseeded Perennial Ryegrass (Transition Aid)

Additional Weeds Controlled

**COOL SEASON TURFGRASSES**

Weed Control in Established Creeping Bentgrass

Roughstalk Bluegrass (*Poa trivialis*) Control

Weed Control in Kentucky Bluegrass

Sedge Control

Roughstalk Bluegrass (*Poa trivialis*) Control

Tall Fescue Control

Additional Weeds Controlled

Weed Control in Mixed Stands of Kentucky Bluegrass

Sedge Suppression

Roughstalk Bluegrass (*Poa trivialis*) Control

Additional Weeds Controlled

**NATIVE GRASSES**

Sedge Control

Tall Fescue Control

Additional Weeds Controlled

**WEEDS CONTROLLED [Optional Section]**

**LIMIT OF WARRANTY AND LIABILITY**



## IMPORTANT PHONE NUMBERS

1. FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE, **800-89-VALENT (898-2536)**
2. IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT, OR FOR MEDICAL ASSISTANCE, CALL COLLECT, DAY OR NIGHT, **(800) 892-0099**

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

Keep out of reach of children.

## CAUTION!

CAUSES MODERATE EYE IRRITATION.

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.</li><li>• Call a poison control center or physician for treatment advice.</li></ul>
<ul style="list-style-type: none"><li>• Have the product container or label with you when calling a poison control center or physician or going for treatment.</li><li>• In case of an emergency involving this product, Call Collect, day or night, (800) 892-0099.</li><li>• This product is identified as <b><i>Certainty</i> Turf Herbicide, EPA Reg. No. 59639-226.</b></li></ul>	

### Environmental Hazards

#### Groundwater Label Advisory

This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

#### Surface Water Label Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of sulfosulfuron from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur with 48 hours. The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of Federal Laws.

## Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:** long-sleeved shirt and long pants shoes plus socks, and chemical-resistant gloves made of any waterproof material, for example Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, Natural Rubber ≥ 14 mils, Poly- ethylene, Polyvinyl Chloride (PVC) ≥ 14 mils, and Viton ≥ 14 mils.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

***For WPS uses only; delete this paragraph when sub-label contains only non-WPS uses:***

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### Environmental Hazards

#### Groundwater Label Advisory

~~This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.~~

#### Surface Water Label Advisory

~~This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of sulfosulfuron from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.~~

~~The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of Federal Laws.~~

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions For Use on this label or in separately published Valent Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **NON-TARGET ORGANISM ADVISORY STATEMENT**

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

### **WINDBLOWN SOIL PARTICLES**

*Certainty* Turf Herbicide has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and directions of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying V-10142 Ag Herbicide if prevailing local conditions may be expected to result in off-site movement.

### **Agricultural Use Requirements**

***Agricultural Use Requirements box can be deleted if sub-label contains only non-WPS uses.***

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material, for example Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, Natural Rubber ≥ 14 mils, Poly-ethylene, Polyvinyl Chloride (PVC) ≥ 14 mils, and Viton ≥ 14 mils.

### **Non-Agricultural Use Requirements**

***Non-Agricultural Use Requirements box can be deleted if sub-label contains only non-WPS uses.***

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (40 CFR Part 170) for agricultural pesticides. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

***[Optional statement for Non-WPS sub-label: Keep people and pets off treated areas until spray solution has dried.]***

## STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**PESTICIDE STORAGE:** Store under cool, dry conditions (below 120°F). Do not store under moist conditions.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable federal, state or local procedures.

See individual container label for DISPOSAL information.

**CONTAINER DISPOSAL:** Emptied container retains product residue. Observe all labeled safeguards until container is destroyed. Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## PRODUCT INFORMATION

**Product Description:** This product is a postemergence, systemic herbicide with limited soil residual activity. It provides postemergence control of many annual and perennial sedges, grass and broadleaf weeds on highly managed turf, sod farms and native grass sites. It is a selective herbicide that can be used over the top of many perennial warm-season and selected cool-season turfgrasses. Refer to the appropriate sections of this label for approved turf species.

**Use Sites:** This product may be used for general weed control on highly managed turf and native grasses. This product may be applied to residential and commercial turf sites, including apartment complexes, athletic fields, cemeteries, golf course fairways, golf course roughs, golf course tees, and other golf course areas, hotel properties, office complexes, parks, public turf areas, retail sites, storage facilities, school grounds, sod and turfgrass seed farms, and other highly managed turfgrass sites. **DO NOT apply directly to or within 4 feet of golf course putting greens.**

This product may be applied at recommended rates as a post-directed spray around any established warm-season ornamental grass or established woody ornamental species in landscaped areas. Avoid contact of this product with leaves of desirable plants as foliar injury, discoloration or loss of the plant may result.

**When to Spray:** Best results are obtained when target weeds are actively growing and not disturbed by mowing for at least 2 days before and 2 days after application.

**Time to Symptoms:** This product is absorbed by both the roots and the foliage of plants and rapidly inhibits the growth of susceptible weeds. Susceptible weed growth stops within 24 hours of treatment even though visual symptoms are slow to develop. Susceptible weeds usually show yellowing or browning within 2 to 3 weeks. Warm, moist conditions following application will accelerate herbicidal activity. Cold, dry conditions will delay herbicidal activity. Weeds stressed by drought are less susceptible to this product.

**Rainfastness:** Heavy rainfall or irrigation within 2 hours after application may wash this product off the foliage and a repeat application may be required for adequate control.

**Maximum Annual Use Rate:** The combined total of all treatments must not exceed 2.66 ounces of this product per acre per year.

**IMPORTANT:** Avoid contact of this product with the roots or foliage of susceptible non-target vegetation as injury may occur. This includes areas where this product may be washed or moved into contact with roots of desirable vegetation.

Susceptible plants may be injured if seeded or transplanted into treated areas unless otherwise directed in this label.

## Weed Resistance Management

For resistance management, *Certainty* Turf Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to *Certainty* Turf Herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of *Certainty* Turf Herbicide or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes or to find out if suspected resistant weeds have been found in their region.
- For further information or to report lack of performance or suspected resistance, contact Valent U.S.A. LLC at 800-898-2536.

## MIXING

Thoroughly clean equipment prior to mixing spray solution.

Eliminate any risk of siphoning the contents of the spray or mixing tank back into the carrier source while mixing. Use approved anti-back-siphoning devices where required by State or local regulations.

### Mixing with Water

This product mixes readily with water. Mix spray solutions of this product as follows. Fill the spray tank with about three-fourths of the desired final volume. Add the recommended amount of this product. Continue the filling process while maintaining agitation. Add nonionic surfactant near the end of the filling process.

### Mixing for Hand-Held Sprayers

Hand-held sprayer applications should be made at a rate of 2 gallons of spray solution per 1000 square feet.

Using the measuring scoop provided in the product packaging, follow the instructions below to prepare the proper spray solution.

#### Measuring Scoop Instructions

Using the **LARGE SCOOP** (0.8-gram scoop) provided, refer to the following table for the appropriate **Mix Volume** (gallons of water) required to achieve the **Desired Application Rate**.

<b>Desired Application Rate<sup>1</sup></b> <b>(ounces of product / acre)</b>	<b>Number of</b> <b>Scoops</b> <b>(large scoop)</b>	<b>Mix Volume</b> <b>(gallons of water)</b>	<b>Spray Rate</b> <b>(gallons/1000 ft<sup>2</sup>)</b>
1.25	1	2.0	2
1.00	1	2.5	2
0.75	1	3.3	2
0.50	1	5.0	2
0.25	1	10.0	2

<sup>1</sup> For 2.0 ounces of product (1.5 ounces active ingredient) per acre, use 2 Large Scoops (0.8-gram scoop) in 2.5 gallons of water and apply at 2 gallons per 1000 square feet.



Using the **SMALL SCOOP** (0.16-gram scoop) provided, refer to the following table for the **Number of Scoops** of product required to achieve the **Desired Application Rate** when mixed in 2 gallons of water.

<b>Desired Application Rate (ounces of product / acre)</b>	<b>Number of Scoops (small scoop)</b>	<b>Mix Volume (gallons of water)</b>	<b>Spray Rate (gallons/1000 ft<sup>2</sup>)</b>
0.25	1	2.0	2
0.50	2	2.0	2
0.75	3	2.0	2
1.00	4	2.0	2
1.25	5	2.0	2

Ensure that product is measured as a level scoop and is not rounded.

Add 2 teaspoons (1/3 fluid ounce) of non-ionic surfactant per gallon of water.

### **Tank-Mixing Procedure**

For tank mixtures, add individual components to the spray tank in the following sequence: water, water dispersible granules (this product), water-soluble bags, dry flowables, emulsifiable concentrates, drift control additive, water-soluble liquids, nonionic surfactants.

### **Surfactants and Adjuvants**

Use a nonionic surfactant at 0.25 to 0.5 percent by volume (1 to 2 quarts per 100 gallons of spray solution). Use only nonionic surfactants that contain at least 90 percent active ingredient. **Do not use nonionic surfactants or other additives that alter the pH of the spray solution below pH 5. Use of surfactants that contain D-Limonene, methylated seed oil, or COC (crop oil concentrate) may cause temporary turf discoloration.**

### **Colorants and Dyes**

Colorants or marking dyes may be added to spray solutions of this product; however, they can reduce product performance. Use colorants and dyes according to the manufacturer's recommendations.

## **APPLICATION EQUIPMENT AND TECHNIQUES**

Apply spray solutions of this product with properly maintained and calibrated equipment capable of delivering desired volumes.

### **Ground Broadcast Equipment**

Apply this product uniformly with properly calibrated ground equipment in 10 to 50 gallons of water per acre. Select spray volumes that ensure thorough and uniform weed coverage. Use equipment that is capable of continuous agitation. Choose nozzles that provide optimum spray distribution and uniform coverage at a spray pressure appropriate for applying postemergence herbicides. Avoid streaking, skips, overlaps and spray drift during application.

|

### **Hand-Held Sprayers**

Backpack sprayers, pump-up pressure sprayers, and other, similar types of hand-held sprayers may be used to apply this product. Apply to foliage of vegetation to be controlled at a rate of 2 gallons of spray solution per 1000 square feet. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

### **Equipment Cleaning**

Thoroughly clean application equipment with a 1-percent solution of ammonia (1 quart of ammonia for every 25 gallons of rinse water) promptly after using this product. Use a sufficient volume of cleaning solution to thoroughly rinse all surfaces and to flush all hoses. Rinse with water and repeat the cleaning procedure with the ammonia solution. Complete the cleaning procedure by rinsing thoroughly with clean water.

If visible residue is present in the spray tank, use a 1-percent solution of ammonia plus 0.25 percent nonionic surfactant (8 fluid ounces per 25 gallons of water) as the cleaning solution.

Do not apply this product through any type of irrigation system.

Do not apply by air.

## MANDATORY SPRAY DRIFT MANAGEMENT

### SPRAY DRIFT

#### Ground Boom Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

#### Boom-less Ground Applications

- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

### SPRAY DRIFT ADVISORIES

#### Boom-less Ground Applications

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

#### Handheld Technology Applications

Take precautions to minimize spray drift.

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

#### Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **BOOM HEIGHT – Ground Boom**

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage.

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### **WIND**

Drift potential generally increases with wind speed.

**AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## **WARM-SEASON TURFGRASSES**

This product generally has been shown to be safe for use on the warm-season turfgrasses listed in this section.

- Bermudagrass (common or hybrid)
- Bahiagrass
- Buffalograss
- Centipedegrass
- Kikuyugrass
- St. Augustinegrass
- Seashore paspalum
- Zoysiagrass

Use of this product may result in temporary chlorosis and may affect the growth pattern or delay green-up of the desirable turf. St. Augustinegrass and seashore paspalum may be more sensitive to this product than other grasses depending on environmental conditions, cultivar differences and other influential factors. For St. Augustinegrass and seashore paspalum, test this product on a small area prior to wide-scale use to determine if this product is suitable for your management and cultural practices.

---

## **Sedge Control**

For the selective control of the weeds listed in this section, apply this product at 1.25 ounces (0.9375 ounce active ingredient) per acre after weeds have reached the 3- to 8-leaf stage of growth. A sequential application of 1.25 ounces of product (0.9375 ounce active ingredient) per acre may be made 4 or more weeks after the initial treatment, if needed. Do not apply more than 2.50 oz of product (1.9 ounces active ingredient) per acre per year. Do not apply more than 2 applications per year.

### **Kyllinga, false green**

*Kyllinga gracilima*

### **Kyllinga, fragrant**

*Kyllinga sesquiflorus*

### **Kyllinga, green**

*Kyllinga brevifolia*

### **Nutsedge, purple**

*Cyperus rotundus*

### **Nutsedge, yellow**

*Cyperus esculentus*

### **Sedge, globe**

*Cyperus croceus*

### **Sedges, annual**

*Cyperus* spp.

## **Tall Fescue Control**

Best control of tall fescue is obtained when this product is applied at 1.25 ounces of product (0.9375 ounce active ingredient) per acre followed by a second application of 1.25 ounces of product (0.9375 ounce active ingredient) per acre at 21 to 28 days after the initial application. If a single application is preferred, apply 2.0 ounces of product (1.5 ounces active ingredient) per acre. Do not apply more than 2.50 oz of product (1.9 ounces active ingredient) per acre per year. Do not make more than 2 applications per year.

---

### **Dallisgrass Suppression in Bermudagrass**

The following application includes the use of MSMA. MSMA can cause injury to common and hybrid bermudagrass turf. Test the following tank-mix recommendation on a small area prior to wide-scale use to determine if this application is suitable for your bermudagrass management objectives.

For suppression of dallisgrass in bermudagrass turf, apply this product, when dallisgrass is actively growing, at a rate of 1.25 ounces of product (0.9375 ounce active ingredient) per acre in a tank mixture with 2 pounds of MSMA per acre and 0.25 percent by volume nonionic surfactant (1 quart per 100 gallons of spray solution). Reapply this same tank mixture 2 to 4 weeks after initial application.

As an alternative program, apply MSMA at 2 pounds active ingredient per acre with 0.25 percent by volume non-ionic surfactant as an initial treatment, wait two weeks and apply 2.0 ounces of this product (1.5 ounces active ingredient) per acre. Wait an additional two weeks and apply MSMA again at 2 pounds active ingredient per acre with 0.25 percent by volume nonionic surfactant.

### **Virginia Buttonweed Suppression**

For suppression of Virginia buttonweed apply this product at 1.25 ounces (0.9375 ounce active ingredient) per acre. This application will provide suppression or partial control of Virginia buttonweed for 4 to 6 weeks.

For enhanced Virginia buttonweed control, tank-mix this product with a broadleaf herbicide labeled for Virginia buttonweed control in the desired warm-season turfgrass. Read and follow the label directions, precautionary statements and all other label information of all products used in the tank mixture. Use all products according to labeled rates and apply the tank mixture according to the most restrictive precautionary statements of the products being used.

### **Annual Bluegrass (*Poa annua*) Control in Non-Overseeded Turf**

For selective control of annual bluegrass (*Poa annua*), apply this product at 1.25 to 2.0 ounces (0.9375 to 1.5 ounces active ingredient) per acre. Use the higher rate of this product for control in areas of established, dense weed infestation. Do not apply more than 2.0 ounces of product (1.5 ounces active ingredient) per acre per year.

Best results are obtained when weeds are in the early stage of growth and prior to tillering.

IN DORMANT BERMUDAGRASS ONLY, tank mixtures of this product with Roundup PRO® or QuikPRO® herbicides may be used to increase the spectrum of vegetation controlled. Read and follow the label directions, precautionary statements, and all other label information on Roundup PRO or QuikPRO herbicides. Refer to the Roundup PRO or QuikPRO product labels for approved application rates. Always apply tank mixtures according to the most restrictive precautionary statements of the products being used.

### **Annual Bluegrass (*Poa annua*) Control Prior to Overseeding Turf with Perennial Ryegrass**

Apply this product at 2.0 ounces (1.5 ounces active ingredient) per acre to control *Poa annua* prior to overseeding warm-season turf with perennial ryegrass. Begin applications after *Poa annua* germination and 7 to 10 days prior to overseeding.



| \_\_\_\_\_

### **Removal of Overseeded Perennial Ryegrass (Transition Aid)**

Best results are obtained by applying this product at 1.25 ounces (0.9375 ounce active ingredient) per acre followed by a second application of 1.25 ounces (0.9375 ounce active ingredient) per acre at 21 to 28 days after the initial application when daily temperatures are expected to exceed 80° F during the treatment period. If a single application is preferred, apply this product at 2.0 ounces (1.5 ounces active ingredient) per acre.

---

## Additional Weeds Controlled

For selective control or suppression of annual or perennial weeds listed in this section, apply this product at 1.25 to 2.0 ounces (0.9375 to 1.5 ounces active ingredient) per acre. Use the higher rate of this product for control in areas of established, dense weed infestation. If using an initial rate of application of 1.25 ounces (0.9375 ounce active ingredient) per acre, a second application of 1.25 ounces (0.9375 ounce active ingredient) per acre may be made 4 or more weeks after the initial treatment, if needed. Do not apply more than 2 applications per year.

### **Barley, little**

*Hordeum pusillum*

### **Bedstraw, catchweed**

*Galium aparine*

### **Beggarweed, Florida<sup>1</sup>**

*Desmodium tortuosum*

### **Bentgrass, creeping<sup>1</sup>**

*Agrostis stolonifera*

### **Bluegrass, roughstalk**

*Poa trivialis*

### **Bluegrass, bulbous**

*Poa bulbosa*

### **Burweed, lawn**

*Soliva pterosperna*

### **Buttercup**

*Ranunculus arvensis*

### **Chamber bitter**

*Phyllanthus urinaria*

### **Chickweed, common**

*Stellaria media*

### **Clover, white**

*Trifolium repens*

### **Crowfootgrass**

*Dactyloctenium aegyptium*

### **Dandelion**

*Taraxacum officinale*

### **Dichondra, Carolina**

*Dichondra carolinensis*

### **Garlic, wild**

*Allium vineale*

### **Geranium, Carolina<sup>1</sup>**

*Geranium carolinianum*

### **Henbit<sup>1</sup>**

*Lamium amplexicaule*

### **Ivy, ground<sup>1</sup>**

*Glechoma hederacea*

### **Johnsongrass**

*Sorghum halepense*

### **Mustard, wild**

*Sinapis arvensis*

### **Pennycress, field**

*Thlaspi arvense*

### **Quackgrass<sup>1</sup>**

*Elytrigia repens*

### **Ryegrass, perennial**

*Lolium perenne*

### **Shepherd's-purse**

*Capsella bursa-pastoris*

### **Violet, wild<sup>1</sup>**

*Viola pratensis*

### **Woodsorrell, yellow**

*Oxalis stricta*

<sup>1</sup> Suppression or partial control only

## COOL-SEASON TURFGRASSES

**[Recommendations in this section are not intended for use in residential turf.]**

This product is recommended for use on creeping bentgrass, Kentucky bluegrass and Kentucky bluegrass mixtures with perennial ryegrass and/or fine fescues.

On cool-season turfgrass, apply this product only when turfgrass is actively growing and mowing is required. Applications made in the fall after temperatures have decreased and regular mowing is no longer required may have an increased potential to cause turf injury and may delay spring green-up.

Use of this product may result in temporary chlorosis and discoloration and may temporarily affect the growth pattern of the desirable turf. These symptoms generally appear 7 to 10 days after application and are typically gone within 21 to 28 days. Turf response is more pronounced under extreme environmental conditions. Perennial ryegrass, fine fescues and creeping bentgrass are more sensitive to this product. If undesirable turf response occurs, skip or delay additional applications to allow turf to recover to a desirable quality. Over-application may result in severe turf injury, thinning or loss of turfgrass stands.

Test this product on a small area prior to wide-scale use to determine if this product is suitable for your turf management and cultural practices.

### **Weed Control in Established Creeping Bentgrass**

For use in established creeping bentgrass on golf course fairways, roughs, approaches and tees.

### **Roughstalk Bluegrass (*Poa trivialis*) Control**

For suppression, apply 0.25 ounce of this product (0.1875 ounce active ingredient) per acre. For control, up to two additional applications of 0.25 ounce of this product (0.1875 ounce active ingredient) per acre at 14- to 21-day intervals may be made. Make all applications between June 1 and September 15. Do not make more than 3 applications per year.

For severe *Poa trivialis* infestations, apply a single application of 0.75 ounce of product (0.5625 ounce active ingredient) per acre in late summer, 21 to 28 days prior to overseeding creeping bentgrass in a renovation approach. This treatment will result in temporary chlorosis and discoloration and will temporarily affect the growth pattern of the desirable turf.

As *Poa trivialis* is controlled, bare patches may occur in the turf. Creeping bentgrass can be sodded or slit-seeded into the treated area beginning 14 days after the last application of this product in order to minimize regrowth of *Poa trivialis* and to maintain overall turf quality.

## **Weed Control in Kentucky Bluegrass**

Weed control recommendations in this section are for use ONLY on established pure stands of Kentucky bluegrass.

### **Sedge Control**

For the selective control of the weeds listed in this section, apply this product at 0.75 ounce (0.5625 ounce active ingredient) per acre after weeds have reached the 3- to 8-leaf stage of growth. A sequential application of 0.75 ounce per acre may be made 4 or more weeks after the initial treatment, if needed.

#### **Kyllinga, green**

*Kyllinga brevifolia*

#### **Nutsedge, purple**

*Cyperus rotundus*

#### **Nutsedge, yellow**

*Cyperus esculentus*

#### **Sedge, globe**

*Cyperus croceus*

#### **Sedges, annual**

*Cyperus* spp.

### **Roughstalk Bluegrass (*Poa trivialis*) Control**

Apply this product at 0.5 ounce (0.375 ounce active ingredient) per acre followed by a second application of 0.5 ounce (0.375 ounce active ingredient) per acre at 21 to 28 days after the initial application. If a single application is preferred, apply this product at 0.75 ounce (0.5625 ounce active ingredient) per acre. Do not make more than 2 applications per year.

As *Poa trivialis* is controlled, bare patches may occur in the turf. Kentucky bluegrass can be sodded or slit-seeded into the treated area beginning 14 days after the last application of this product in order to minimize regrowth of *Poa trivialis* and to maintain overall turf quality.

### **Tall Fescue Control**

Apply this product at 0.75 ounce (0.5625 ounce active ingredient) per acre followed by a second application of 0.75 ounce (0.5625 ounce active ingredient) per acre at 21 to 28 days after the initial application. Do not make more than 2 applications per year.

## Additional Weeds Controlled

For selective control or suppression of annual or perennial weeds listed in this section, apply this product at 0.75 ounce (0.5625 ounce active ingredient) per acre. A sequential application of 0.75 ounce (0.5625 ounce active ingredient) per acre may be made 4 or more weeks after the initial treatment, if needed. Do not make more than 2 applications per year.

### **Barley, little**

*Hordeum pusillum*

### **Bedstraw, catchweed**

*Galium aparine*

### **Bluegrass, annual**

*Poa annua*

### **Bluegrass, bulbous**

*Poa bulbosa*

### **Burweed, lawn**

*Soliva pterosperna*

### **Buttercup**

*Ranunculus arvensis*

### **Chamber bitter**

*Phyllanthus urinaria*

### **Chamomile, mayweed**

*Anthemus cotula*

### **Chickweed, common**

*Stellaria media*

### **Clover, white**

*Trifolium repens*

### **Dandelion**

*Taraxacum officinale*

### **Garlic, wild<sup>1</sup>**

*Allium vineale*

### **Henbit<sup>1</sup>**

*Lamium amplexicaule*

### **Ivy, ground<sup>1</sup>**

*Glechoma hederacea*

### **Johnsongrass**

*Sorghum halepense*

### **Mustard, tumble**

*Sisymbrium altissimum*

### **Mustard, wild**

*Sinapis arvensis*

### **Pennycress, field**

*Thlaspi arvense*

### **Quackgrass<sup>1</sup>**

*Elytrigia repens*

### **Shepherd's-purse**

*Capsella bursa-pastoris*

### **Violet, wild**

*Viola pratensis*

### **Woodsorrell, yellow**

*Oxalis stricta*

<sup>1</sup> Suppression or partial control only

## Weed Control in Mixed Stands of Kentucky Bluegrass

This product is recommended for use only on established stands of Kentucky bluegrass that include perennial ryegrass and/or fine fescues.

Avoid application of this product on stands with a high percentage of perennial ryegrass or fine fescue.

Perennial ryegrass and fine fescues are more sensitive to this product, therefore, as the percentage of these grasses increase in the mix, occurrence of undesirable turf response will also increase. If undesirable turf response occurs, skip or delay additional applications to allow turf to recover to a desirable quality.

Over-application may result in severe turf injury, thinning or loss of turfgrass stands.

## **Sedge Suppression**

For the selective suppression of the weeds listed below apply this product at 0.5 ounce (0.375 ounce active ingredient) per acre after weeds have reached the 3- to 8-leaf stage of growth. A sequential application of 0.5 ounce (0.375 ounce active ingredient) per acre may be made 28 or more days after the initial treatment, if needed. Do not make more than 2 applications per year.

[*Optional Statement:* Alternatively, for suppression of the weeds listed below, apply 0.25 ounce of this product per acre. For additional control, up to three additional applications of 0.25 ounce of this product per acre at 28-day intervals may be made, if needed. ]

### **Nutsedge, purple**

*Cyperus rotundus*

### **Nutsedge, yellow**

*Cyperus esculentus*

### **Sedge, globe**

*Cyperus croceus*

### **Sedges, annual**

*Cyperus* spp.

## **Roughstalk Bluegrass (*Poa trivialis*) Control**

For suppression, apply 0.25 ounce of this product (0.1875 ounce active ingredient) per acre. For control, up to two additional applications of 0.25 ounce of this product (0.1875 ounce active ingredient) per acre at 14- to 21-day intervals may be made. Make all applications between June 1 and September 15. Do not make more than 3 applications per year.

For severe *Poa trivialis* infestations, apply a single application of 0.5 ounce of product (0.375 ounce active ingredient) per acre in late summer, 21 to 28 days prior to overseeding turfgrass in a renovation approach.

As *Poa trivialis* is controlled, bare patches may occur in the turf. Kentucky bluegrass, perennial ryegrass or fine fescues can be sodded or slit-seeded into the treated area beginning 14 days after the last application of this product in order to minimize regrowth of *Poa trivialis* and to maintain overall turf quality.

## Additional Weeds Controlled

For selective control or suppression of annual or perennial weeds listed in this section, apply this product at 0.5 ounce (0.375 ounce active ingredient) per acre. A sequential application of product 0.5 ounce (0.375 ounce active ingredient) per acre may be made 4 or more weeks after the initial treatment, if needed.

[Optional Statement: Alternatively, for suppression of the weeds listed in this section, apply 0.25 ounce of this product (0.1875 ounce active ingredient) per acre. For additional control, up to three additional applications of 0.25 ounce of this product (0.1875 ounce active ingredient) per acre at 28-day intervals may be made, if needed. Do not make more than 2 applications per year.]

### **Bedstraw, catchweed<sup>1</sup>**

*Galium aparine*

### **Bluegrass, annual<sup>1</sup>**

*Poa annua*

### **Bluegrass, bulbous<sup>1</sup>**

*Poa bulbosa*

### **Chamber bitter<sup>1</sup>**

*Phyllanthus urinaria*

### **Chamomile, mayweed<sup>1</sup>**

*Anthemus cotula*

### **Chickweed, common**

*Stellaria media*

### **Clover, white<sup>1</sup>**

*Trifolium repens*

### **Dandelion<sup>1</sup>**

*Taraxacum officinale*

### **Fescue, tall<sup>1</sup>**

*Festuca arundinacea*

### **Garlic, wild<sup>1</sup>**

*Allium vineale*

### **Henbit<sup>1</sup>**

*Lamium amplexicaule*

### **Ivy, ground<sup>1</sup>**

*Glechoma hederacea*

### **Johnsongrass<sup>1</sup>**

*Sorghum halepense*

### **Mustard, tumble**

*Sisymbrium altissimum*

### **Mustard, wild**

*Sinapis arvensis*

### **Pennycress, field**

*Thlaspi arvense*

### **Quackgrass<sup>1</sup>**

*Elytrigia repens*

### **Shepherd's-purse**

*Capsella bursa-pastoris*

### **Woodsorrell, yellow<sup>1</sup>**

*Oxalis stricta*

<sup>1</sup> Suppression or partial control only



## **ORNAMENTALS**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

See "GENERAL INFORMATION" and "MIXING" sections of the label, booklet for Certainty turf herbicide for essential product performance information.

This product is for use in established woody ornamentals, perennial groundcovers and warm-season ornamental grasses growing in landscaped areas or field production nurseries. Do not apply this product to container plants or production beds of potted plants. For selective control or suppression of weeds listed in this section, apply this product at 1.25 ounces per acre. A second application of 1.25 ounces per acre may be made 4 or more weeks after the initial treatment if needed. Best results are obtained when target weeds are actively growing and not disturbed by mowing for at least 2 days before and 2 days after application.

<u>Barley, little</u>	<u><i>Hordeum pusillum</i></u>
<u>Bedstraw, catchweed</u>	<u><i>Galium aparine</i></u>
<u>Beggarweed, Florida<sup>1</sup></u>	<u><i>Desmodium tortuosum</i></u>
<u>Bluegrass, annual</u>	<u><i>Poa annua</i></u>
<u>Bluegrass, bulbous</u>	<u><i>Poa bulbosa</i></u>
<u>Bluegrass, roughstalk</u>	<u><i>Poa trivialis</i></u>
<u>Burweed, lawn</u>	<u><i>Soliva pterosperna</i></u>
<u>Buttercup</u>	<u><i>Ranunculus arvensis</i></u>
<u>Chamber bitter</u>	<u><i>Phyllanthus urinaria</i></u>
<u>Chickweed, common</u>	<u><i>Stellaria media</i></u>
<u>Clover, white</u>	<u><i>Trifolium repens</i></u>
<u>Crowfootgrass</u>	<u><i>Dactyloctenium aegyptium</i></u>
<u>Dandelion</u>	<u><i>Taraxacum officinale</i></u>
<u>Dichondra, Carolina</u>	<u><i>Dichondra carolinensis</i></u>
<u>Fescue, tall</u>	<u><i>Festuca arundinacea</i></u>
<u>Garlic, wild</u>	<u><i>Allium vineale</i></u>
<u>Geranium, Carolina<sup>1</sup></u>	<u><i>Geranium carolinianum</i></u>
<u>Henbit<sup>1</sup></u>	<u><i>Lamium amplexicaule</i></u>
<u>Ivy, ground<sup>1</sup></u>	<u><i>Glechoma hederacea</i></u>
<u>Johnsongrass</u>	<u><i>Sorghum halepense</i></u>
<u>Kyllinga, false green</u>	<u><i>Kyllinga gracilima</i></u>
<u>Kyllinga, fragrant</u>	<u><i>Kyllinga sesquiflorus</i></u>
<u>Kyllinga, green</u>	<u><i>Kyllinga brevifolia</i></u>
<u>Mustard, wild</u>	<u><i>Sinapis arvensis</i></u>
<u>Nutsedge, purple</u>	<u><i>Cyperus rotundus</i></u>

Continued

<u>Nutsedge, yellow</u>	<u>Cyperus esculentus</u>
<u>Pennycress, field</u>	<u>Thlaspi arvense</u>
<u>Pennywort, lawn<sup>1</sup> (dollarweed)</u>	<u>Hydrocotyle sibthorpioides</u>
<u>Quackgrass<sup>1</sup></u>	<u>Elytrigia repens</u>
<u>Ryegrass, perennial</u>	<u>Lolium perenne</u>
<u>Sedge, globe-</u>	<u>Cyperus croceus</u>
<u>Sedges, annual</u>	<u>Cyperus spp.</u>
<u>Shepherd's-purse</u>	<u>Capsella bursa-pastoris</u>
<u>Violet, wild<sup>1</sup></u>	<u>Viola pratincola</u>
<u>Wood sorrel, yellow<sup>1</sup></u>	<u>Oxalis stricta</u>

<sup>1</sup> Suppression or partial control only

### **Directed-Spray Applications**

This product may be applied as a post-directed spray around any established warm-season ornamental grass or established woody ornamental species in landscaped areas or field production nurseries. Avoid contact of this product with leaves of desirable plants as foliar injury, discoloration or loss of the plant may result.

### **Over-the-Top Applications**

This product may be applied in established landscaped areas as an over-the-top application on the ornamental and groundcover species listed below.

### **Pre-Plant Applications**

This product may be applied prior to planting the ornamental species listed below. Wait 14 days after the last application of this product before planting.

<b><u>Common Name</u></b>	<b><u>Scientific-Name</u></b>	<b><u>Pre-Plant</u></b>	<b><u>Over-The-Top</u></b>
<u>Arborvitae, American</u>	<u>Thuja occidentalis</u>	-	<u>X</u>
<u>Azalea, Dwarf</u>	<u>Rhododendron atlanticum</u>	-	<u>X</u>
<u>Bougainvillea*</u>	<u>Bougainvillea glabra</u>	-	<u>X</u>
<u>Boxwood, Green Mountain</u>	<u>Buxus 'Green Mountain'</u>	<u>X</u>	-
<u>Boxwood, Green Velvet</u>	<u>Buxus 'Green Velvet'</u>	<u>X</u>	<u>X</u>
<u>Burning Bush, Dwarf</u>	<u>Euonymus alatus 'Compacta'</u>	<u>X</u>	-
<u>Euonymus, Wintercreeper</u>	<u>Euonymus fortunei</u>	<u>X</u>	<u>X</u>
<u>Forsythia</u>	<u>Forsythia x intermedia</u>	<u>X</u>	-
<u>Gardenia</u>	<u>Gardenia jasminoides</u>	-	<u>X</u>
<u>Holly, Blue</u>	<u>Ilex x meserveae</u>	<u>X</u>	<u>X</u>

Continued

<u>Holly, Chinese</u>	<u>Ilex cornuta</u>	-	X
<u>Hydrangea, Panicked</u>	<u>Hydrangea paniculata</u>	X	-
<u>Ivy, English</u>	<u>Hedera helix</u>	X	-
<u>Japanese Snowball Bush</u>	<u>Viburnum plicatum var. tomentosum</u>	-	X
<u>Jasmine, Asiatic</u>	<u>Trachelospermum asiaticum</u>	-	X
<u>Jasmine, Star</u>	<u>Trachelospermum jasminoides</u>	-	X
<u>Jasmine, Winter</u>	<u>Jasminium nudiflorum</u>	X	-
<u>Juniper, Chinese</u>	<u>Juniperus chinensis</u>	X	X
<u>Juniper, Creeping</u>	<u>Juniperus horizontalis</u>	X	X
<u>Juniper, Shore</u>	<u>Juniperus conferta</u>	-	X
<u>Lilac</u>	<u>Syringa vulgaris</u>	-	X
<u>Lilac, Dwarf Korean</u>	<u>Syringa meyeri 'Palibin'</u>	X	-
<u>Mockorange, Japanese</u>	<u>Pittosporum tobira</u>	-	X
<u>Mondo Grass</u>	<u>Ophiopogon japonicus</u>	-	X
<u>Monkey Grass, Big Blue</u>	<u>Liriope muscari 'Big Blue'</u>	-	X
<u>Monkey Grass, Variegated</u>	<u>Liriope muscari 'Variegata'</u>	-	X
<u>Ninebark</u>	<u>Physocarpus opulifolius</u>	-	X
<u>Oleander</u>	<u>Nerium oleander</u>	-	X
<u>Periwinkle, Greater</u>	<u>Vinca major</u>	-	X
<u>Photinia, Fraser</u>	<u>Photinia x fraseri</u>	-	X
<u>Pine, Mugo</u>	<u>Pinus mugo</u>	X	X
<u>Privet, Golden</u>	<u>Ligustrum 'Vicaryi'</u>	X	-
<u>Redbud</u>	<u>Cercis canadensis</u>	X	-
<u>Rhododendron</u>	<u>Rhododendron spp.</u>	X	X
<u>Rosemary</u>	<u>Rosmarinus officinalis</u>	-	X
<u>Serviceberry</u>	<u>Amelanchier alnifolia</u>	X	-
<u>Spirea, Goldmound</u>	<u>Spirea japonica 'Goldmound'</u>	-	X
<u>Viburnum, American Cranberrybush</u>	<u>Viburnum trilobum</u>	X	-
<u>Viburnum, Judd</u>	<u>Viburnum x juddii</u>	-	X
<u>Viburnum, Prague</u>	<u>Viburnum 'Pragense'</u>	X	-
<u>Weigela</u>	<u>Weigela florida</u>	X	-

**\*Single application only**

## NATIVE GRASSES

This product generally has been shown to be safe for use on the warm-season native grasses listed in this section.

- Big bluestem
- Little bluestem
- Bushy bluestem
- Blue oats grama
- Buffalograss
- Indiangrass
- Lovegrass
- Switchgrass

Use of this product may result in temporary chlorosis or temporarily affect the growth pattern of these native grasses. If discoloration or excessive thinning of the native grasses occurs, skip or delay additional applications to allow the native grasses to recover to a desirable quality.

Test this product on a small area prior to wide-scale use to determine if this product is suitable for your management and cultural practices.

## Sedge Control

For the selective control of the weeds listed in this section, apply this product at 1.25 ounces (0.9375 ounce active ingredient) per acre after weeds have reached the 3- to 8-leaf stage of growth. A sequential application of 1.25 ounces (0.9375 ounce active ingredient) per acre may be made 4 or more weeks after the initial treatment, if needed. Do not apply more than 2 applications per year.

### **Kyllinga, false green**

*Kyllinga gracilima*

### **Kyllinga, fragrant**

*Kyllinga sesquiflorus*

### **Kyllinga, green**

*Kyllinga brevifolia*

### **Nutsedge, purple**

*Cyperus rotundus*

### **Nutsedge, yellow**

*Cyperus esculentus*

### **Sedge, globe**

*Cyperus croceus*

### **Sedges, annual**

*Cyperus spp.*

## Tall Fescue Control

Best control of tall fescue is obtained when this product is applied at 1.25 ounces (0.9375 ounce active ingredient) per acre followed by a second application of 1.25 ounces (0.9375 ounce active ingredient) per acre at 21 to 28 days after the initial application. If a single application is preferred, apply this product at 2.0 ounces (1.5 ounce active ingredient) per acre.

## Additional Weeds Controlled

For selective control or suppression of annual or perennial weeds listed in this section, apply this product at 1.25 ounces (0.9375 ounce active ingredient) per acre. A sequential application of 1.25 ounces (0.9375 ounce active ingredient) per acre may be made 4 or more weeks after the initial treatment, if needed. Do not apply more than 2 applications per year.

### **Barley, little**

*Hordeum pusillum*

### **Bedstraw, catchweed**

*Galium aparine*

### **Beggarweed, Florida<sup>1</sup>**

*Desmodium tortuosum*

### **Bentgrass, creeping**

*Agrostis stolonifera*

### **Bluegrass, annual<sup>1</sup>**

*Poa annua*

### **Bluegrass, roughstalk**

*Poa trivialis*

### **Bluegrass, bulbous**

*Poa bulbosa*

### **Burweed, lawn**

*Soliva pterosperma*

### **Buttercup**

*Ranunculus arvensis*

### **Buttonweed, Virginia<sup>1</sup>**

*Diodia virginiana*

### **Chamber bitter**

*Phyllanthus urinaria*

### **Chickweed, common**

*Stellaria media*

### **Clover, white**

*Trifolium repens*

### **Crowfootgrass**

*Dactyloctenium aegyptium*

### **Dallisgrass<sup>1</sup>**

*Paspalum dilatatum*

### **Dandelion**

*Taraxacum officinale*

### **Dichondra, Carolina**

*Dichondra carolinensis*

### **Garlic, wild**

*Allium vineale*

### **Geranium, Carolina<sup>1</sup>**

*Geranium carolinianum*

### **Henbit<sup>1</sup>**

*Lamium amplexicaule*

### **Ivy, ground**

*Glechoma hederacea*

### **Johnsongrass**

*Sorghum halepense*

### **Mustard, wild**

*Sinapis arvensis*

### **Pennycress, field**

*Thlaspi arvense*

### **Quackgrass<sup>1</sup>**

*Elytrigia repens*

### **Ryegrass, perennial**

*Lolium perenne*

### **Shepherd's-purse**

*Capsella bursa-pastoris*

### **Violet, wild<sup>1</sup>**

*Viola pratensis*

### **Woodsorrell, yellow**

*Oxalis stricta*

<sup>1</sup> Suppression or partial control only

[Optional Section

**WEEDS CONTROLLED**

**Chamber bitter**

*Phyllanthus urinaria*

**Chamomile, mayweed**

*Anthemus cotula*

**Chickweed, common**

*Stellaria media*

**Clover, white**

*Trifolium repens*

**Crowfootgrass**

*Dactyloctenium aegyptium*

**Dallisgrass<sup>1</sup>**

*Paspalum dilatatum*

**Dandelion**

*Taraxacum officinale*

**Dichondra, Carolina**

*Dichondra carolinensis*

**Fescue, tall**

*Festuca arundinacea*

**Garlic, wild**

*Allium vineale*

**Geranium, Carolina**

*Geranium carolinianum*

**Henbit<sup>1</sup>**

*Lamium amplexicaule*

**Ivy, ground**

*Glechoma hederacea*

**Johnsongrass**

*Sorghum halepense*

**Kyllinga, false green**

*Kyllinga gracilima*

**Kyllinga, fragrant**

*Kyllinga sesquiflorus*

**Kyllinga, green**

*Kyllinga brevifolia*

**Mustard, tumble**

*Sisymbrium altissimum*

**Mustard, wild**

*Sinapis arvensis*

**Nutsedge, purple**

*Cyperus rotundus*

**Nutsedge, yellow**

*Cyperus esculentus*

**Pennycress, field**

*Thlaspi arvense*

**Pennywort, lawn<sup>1</sup>**

*Hydrocotyle sibthorpioides*

**Quackgrass<sup>1</sup>**

*Elytrigia repens*

**Ryegrass, perennial**

*Lolium perenne*

**Sedge, globe**

*Cyperus croceus*

**Sedges, annual**

*Cyperus spp.*

**Shepherd's-purse**

*Capsella bursa-pastoris*

**Violet, wild**

*Viola pratensis*

**Woodsorrell, yellow**

*Oxalis stricta*

<sup>1</sup> Suppression or partial control only]

**RISKS OF USING THIS PRODUCT,  
LIMITED WARRANTY AND DISCLAIMER,  
AND LIMITATION OF LIABILITY**

**IMPORTANT: Read the entire Label including this section titled Risks of Using this Product, Limited Warranty and Disclaimer, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.**

**RISKS OF USING THIS PRODUCT**

The buyer and user (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

The Directions for Use of this product must be followed carefully. Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential, or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. To the extent consistent with applicable law, Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

**LIMITED WARRANTY AND DISCLAIMER**

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label **and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

**LIMITATION OF LIABILITY**

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

**PROMPT NOTICE OF CLAIM**

To the extent consistent with applicable law allowing such requirements, Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is later, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law, if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

**NO AMENDMENTS**

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Risks of Using This Product, Limited Warranty and Disclaimer, and Limitation of Liability**, which may not be modified by any oral or written agreement.

## **LIMIT OF WARRANTY AND LIABILITY**

~~Valent U.S.A. LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label pamphlet ("Directions") when used in accordance with those Directions under the conditions described therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.~~

~~To the extent consistent with applicable law, buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.~~

~~Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company to the extent consistent with applicable law, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.~~

~~TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.~~

~~This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.~~

~~Upon opening and using this product, buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement. If terms are not acceptable, return at once unopened.~~



Roundup PRO (EPA Reg. No. 524-475) and QuikPRO (EPA Reg. No. 524-435) are registered trademarks of Monsanto Technology LLC.

*Certainty* is a registered trademark of Valent U.S.A. LLC

---

In case of an emergency involving this product,  
Call day or night, (800) 892-0099.

---

©2023 Valent U.S.A. LLC

Manufactured for:

**Valent U.S.A. LLC**

P.O. Box 5075

San Ramon CA 94583



**[Nufarm]**

[Grow a better tomorrow.]

[Distributed by:

Nufarm Americas Inc.

11901 S. Austin Ave.,

Alsip, IL 60803]

EPA Reg. No. 59639-226

059639-00226.20230509.Amend.CertaintyTurfHerb.Red