



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

59639-254

Date of Issuance:

6/29/22

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

INTEGO® SUITE Canola
Seed Protectant

Name and Address of Registrant (include ZIP Code):

Valent U.S.A., LLC
4600 Norris Canyon Road
San Ramon, CA 94583

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 59639-254."

Signature of Approving Official:

Rachel C. Holloman, Chief
Fungicide and Herbicide Branch
Registration Division (7505P)

Date:

6/29/22

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 06/22/2022
- Alternate CSF 1 dated 06/23/2022

If you have any questions, please contact Lindsay DeMers via email at demers.lindsay@epa.gov.

Enclosure: stamped "accepted" label



ETHABOXAM	GROUP	22	FUNGICIDE
MANDESTROBIN	GROUP	11	FUNGICIDE
METCONAZOLE	GROUP	3	FUNGICIDE
METALAXYL	GROUP	4	FUNGICIDE

[Bracketed italicized text is information for the reader and is not part of the label.]

[Bracketed text is optional]

INTEGO® SUITE Canola Seed Protectant

A SEED TREATMENT PRODUCT PROVIDING SYSTEMIC SEED AND SEEDLING PROTECTION AGAINST LISTED DISEASES

Active Ingredients	By Wt
Ethaboxam ¹	0.830%
Mandestrobin ²	1.110%
Metconazole ³	0.166%
Metalaxyl ⁴	0.442%
Other Ingredients	97.452%
Total	100.00%

¹(*RS*)-*N*-[cyano(2-thienyl)methyl]-4-ethyl-2-(ethylamino)thiazole-5-carboxamide

²(*2RS*)-2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-*N*-methylacetamide

³(*1RS,5RS,1RS,5SR*)-5-(4-chlorobenzyl)-2,2-dimethyl-1-(1*H*-1,2,4-triazol-1-ylmethyl)cyclopentanol

⁴methyl *N*-(2,6-dimethylphenyl)-*N*-(methoxyacetyl)-DL-alaninate

One gallon of *INTEGO*® SUITE Canola Seed Protectant contains 0.085 lb ethaboxam, 0.114 lb mandestrobin, 0.045 lb metalaxyl and 0.017 lb metconazole.


KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE [NEXT] PAGE [PANEL][BOOKLET] FOR [PRECAUTIONARY STATEMENTS] [AND] [DIRECTIONS FOR USE]

NET CONTENTS

EPA Reg. No. 59639-XXX
EPA Est.



HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators and other handlers (including conducting maintenance and calibration of treater) must wear: long-sleeved shirt, long pants, socks and shoes, and chemical-resistant gloves made of any waterproof material.

Baggers, bag sewers and bag stackers must wear: long-sleeved shirt, long pants, socks and shoes.

Workers who load and plant treated seed must wear: long-sleeved shirt, long pants, socks and shoes, and chemical-resistant gloves made of any waterproof material when handling treated seed.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, oysters, and shrimp. **DO NOT** apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. **DO NOT** apply where runoff is likely to occur. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Apply this product only as specified on label.

Groundwater Advisory

This product has properties and characteristics associated with chemicals detected in ground water. It may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore, contain any product spills or equipment leaks and dispose of waste according to the disposal instructions on this label.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several days after application. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential loading of chemicals from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

For use only in commercial seed treatment facilities. Not for use in hopper box, planter box, slurry box or other on-farm seed treatment applications. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

MODES OF ACTION

Metalaxyl fungicide contained in *INTEGO*[®] SUITE Canola Seed Protectant is a member of the acylalanine chemical group (FRAC Code 4) and interrupts fungal nucleic acid synthesis. Metconazole belongs to the sterol biosynthesis inhibitor group of fungicides (FRAC Code 3). Ethaboxam is a member of the thiazole carboxamide chemical group (FRAC Code 22), and acts by inhibition of cell division. Mandestrobin belongs to the strobilurin or Quinone Outside Inhibitors (QoI) class (FRAC Code 11) of fungicides.

PRODUCT INFORMATION

INTEGO[®] SUITE Canola Seed Protectant is a fungicide product formulated for seed treatment use to provide protection against listed seed-borne and soil-borne pathogens.

The active ingredients in *INTEGO*[®] SUITE Canola Seed Protectant are metconazole, a broad-spectrum, systemic, triazole fungicide; metalaxyl, an Oomycete-active, systemic acylalanine fungicide; ethaboxam, an Oomycete-active fungicide; and mandestrobin, a strobilurin systemic fungicide. The fungicide components of *INTEGO*[®] SUITE Canola Seed Protectant provide activity against *Pythium* spp., *Rhizoctonia solani*, *Fusarium* spp. and seed decay fungi causing seed rots.

RESTRICTIONS

- *INTEGO*[®] SUITE Canola Seed Protectant must be used only in accordance with specifications on this label utilizing mechanical, slurry, or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed.
- This product is for use only in commercial seed treatment facilities.
- **DO NOT** re-use bags from treated seed to handle food or feed products. Dispose of seed packaging in accordance with local requirements.
- As stated by regulations specified in 40 CFR 153.155, all seeds treated with *INTEGO*[®] SUITE Canola Seed Protectant must be colored with an EPA-approved dye or colorant to visually distinguish seed that is treated with a pesticide. This product contains blue colorant. This regulation is necessary to prevent treated seed from accidental use as a food for man or feed for animals.
- Rape greens grown and harvested from *INTEGO*[®] SUITE Canola Seed Protectant - treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from *INTEGO*[®] SUITE Canola Seed Protectant treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.
- **DO NOT** use treated seed for feed, food, or oil purposes.
- Store treated seed away from feeds and foodstuffs. **DO NOT** store leftover treated seed where there is a chance of it becoming mixed with untreated seed.
- Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- Treated seed must be planted into the soil at a minimum depth of 0.5 inch.
- Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- The maximum annual application rate for metalaxyl/mefenoxam must not exceed 12.3 lb per acre per calendar year or the lesser value listed in the crop use section.
- The maximum total application rate (including foliar and seed treatment) on Rapeseed Subgroup 20A is 0.0008 lb ai for metalaxyl, 0.1252 lb ai for metconazole, 0.0006 lb ai for ethaboxam and 0.3758 lb ai for mandestrobin per acre per year. This seed has been treated with 0.0075 lb ai ethaboxam, 0.010 lb ai mandestrobin, 0.004 lb ai metalaxyl and 0.0015 lb ai metconazole per 100 lb seed.
- **DO NOT** plant any crop not listed on this label within 30 days of planting *INTEGO*[®] SUITE Canola Seed Protectant-treated seed.
- After treated seed has been planted, **DO NOT** enter or allow worker entry into areas planted with treated seeds during the restricted-entry interval (REI) of 24 hours.
EXCEPTION: Workers may enter these areas without restriction if there will be no worker contact with the soil/media subsurface.

APPLICATION DIRECTIONS

Always mix product thoroughly before use. Accurate application of *INTEGO*[®] SUITE Canola Seed Protectant to seed is required for best performance. Thoroughly mix *INTEGO*[®] SUITE Canola Seed Protectant in the desired seed slurry mixture, prior to application to seed. Pre-test tank mixtures with other seed treatment products to evaluate formulation compatibility and to ensure proper physical compatibility of products. Follow the most restrictive limitations or cautions on labels of all products used in a seed treatment mixture. **DO NOT** tank mix with any products which contain a prohibition on tank mixing.

It is the pesticides user's responsibility to ensure that all products are registered for intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions and precautionary statements of each product in the mixture.

RESISTANCE MANAGEMENT

For resistance management, please note that *INTEGO*[®] SUITE Canola Seed Protectant contains a Group 3 fungicide/metconazole, a Group 4 fungicide/metalaxyl, a Group 11 fungicide/mandestrobin, and a Group 22 fungicide/ethaboxam. Any fungal population may contain individuals naturally resistant to *INTEGO*[®] SUITE Canola Seed Protectant and other Group 3, 4, 11, or 22 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of *INTEGO*[®] SUITE Canola Seed Protectant or other Group 3, 4, 11 or 22, fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.

Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pathogens problems in your area.

For further information or to report suspected resistance contact Valent U.S.A. LLC at 800-6-VALENT (682-5368). You can also contact your pesticide distributor or university extension specialist to report resistance.

Seed Quality and Storage of Treated Seed

Use proper storage conditions for treated seed. Due to seed quality and seed storage conditions beyond the control of Valent U.S.A. LLC, no claims are made to guarantee the germination of carry-over treated seed.

SEED BAG TAG

The Federal Seed Act requires that the container of seed treated with *INTEGO*[®] SUITE Canola Seed Protectant containing ethaboxam, mandestrobin, metalaxyl and metconazole must be labeled with the following statements:

- This seed has been treated with *INTEGO*[®] SUITE Canola Seed Protectant which contains ethaboxam, mandestrobin, metalaxyl and metconazole.
- **DO NOT** use treated seed for feed, food or oil processing.

In addition, the U.S. Environmental Protection Agency requires the following statements on the container of seed treated with *INTEGO*[®] SUITE Canola Seed Protectant:

- This seed has been treated with 0.0075 lb ai ethaboxam, 0.010 lb ai mandestrobin, 0.004 lb ai metalaxyl and 0.0015 lb ai metconazole per 100 lb seed.
- The maximum total application rate (including foliar and seed treatment) on Rapeseed Subgroup 20A is 0.0008 lb ai for metalaxyl, 0.1252 lb ai for metconazole, 0.0006 lb ai for ethaboxam and 0.3758 lb ai for mandestrobin per acre per year.
- Wear long sleeved shirt, long pants, socks and shoes, and chemical resistant gloves made of any waterproof material when handling treated seed.
- Store away from feeds and other foodstuffs.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- **DO NOT** contaminate water bodies when disposing of planting equipment washwaters.
- **DO NOT** re-use bags from treated seed to handle food or feed products.
- Dispose of seed packaging in accordance with local requirements.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Treated seed must be planted into the soil at a minimum depth of 0.5 inch.
- **DO NOT** allow children, pets, or livestock to have access to treated seeds.
- Rape greens grown and harvested from *INTEGO*[®] SUITE Canola Seed Protectant treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from *INTEGO*[®] SUITE Canola Seed Protectant treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.
- As stated by regulations specified in 40 CFR 153.155, all seeds treated with *INTEGO*[®] SUITE Canola Seed Protectant must be colored with an EPA-approved dye or colorant to visually distinguish seed that is treated with a pesticide. This product contains blue colorant. This regulation is necessary to prevent treated seed from accidental use as a food for man or feed for animals.
- **DO NOT** plant any crop not listed on this label within 30 days of planting *INTEGO*[®] SUITE Canola Seed Protectant-treated seed.
- After treated seed has been planted, **DO NOT** enter or allow worker entry into areas planted with treated seeds during the restricted-entry interval (REI) of 24 hours. EXCEPTION: Workers may enter these areas without restriction if there will be no worker contact with the soil/media subsurface.

- **DO NOT** store leftover treated seed where there is a chance of it becoming mixed with untreated seed.
- Store treated seed in cool, dry conditions.

The following table provides use rate of *INTEGO*[®] SUITE Canola Seed Protectant with disease protection guidelines.

Rapeseed Subgroup 20A	
Borage; canola; crambe; cuphea; echium; flax seed; gold of pleasure; hare's ear mustard; lesquerella; lunaria; meadowfoam; milkweed; mustard seed; oil radish; poppy seed; rapeseed; sesame; sweet rocket; cultivars, varieties, and/or hybrids of these.	
Diseases	Application Rate
Seed rots, seedling blight and seedling root rot caused by <i>Rhizoctonia solani</i> , and <i>Fusarium</i> spp. Seed-borne blackleg (<i>Leptosphaeria maculans</i>) Seed rot/pre-emergence damping-off caused by <i>Pythium</i> spp.	11.27 fl oz/100 lb seed (Ethaboxam – 7.5 g ai/100 kg seed) (Mandestrobin – 10 g ai/100 kg seed) (Metalaxyl – 4 g ai/100 kg seed) (Metconazole – 1.5 g ai/100 kg seed)
RESTRICTIONS:	
<ul style="list-style-type: none"> • The maximum total application rate (including foliar and seed treatment) on Rapeseed Subgroup 20A is 0.0008 lb ai for metalaxyl, 0.1252 lb ai for metconazole, 0.0006 lb ai for ethaboxam and 0.3758 lb ai for mandestrobin per acre per year. 	

**RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY AND DISCLAIMER,
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this section titled Risks of Using this Product, Limited Warranty and Disclaimer, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The buyer and user (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

The Directions for Use of this product must be followed carefully. Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential, or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. To the extent consistent with applicable law, Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY AND DISCLAIMER

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label **and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

To the extent consistent with applicable law allowing such requirements, Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is later, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law, if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Risks of Using This Product, Limited Warranty and Disclaimer, and Limitation of Liability**, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law.

Read and follow the entire label of each product to be used in the tank mix with this product.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Store in a dry place away from excessive heat.

DO NOT store near food or feed.

Store in original container only.

Not for use or storage around the home.

To close package, replace and tighten cap to form an airtight seal.

PESTICIDE DISPOSAL

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER HANDLING

[Use the following statement for containers equal to or less than 5 gallons]

[Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or other procedures allowed by State and local authorities.]

[Use the following statement for containers greater than 5 gallons]

[Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or other procedures allowed by State and local authorities.]

[Use the following statement for all formulation types/all refillable container types]

[Refillable container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Offer for recycling, if available; otherwise dispose of in a sanitary landfill or other procedures allowed by State and local authorities.]

RECYCLING

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 877-952-2272 (toll free) or www.acrecycle.org.

©2022 Valent U.S.A. LLC

INTEGO is a registered trademark of Valent U.S.A. LLC

Manufactured For:

Valent U.S.A. LLC

P. O. Box 5075

San Ramon CA 94583

Made in U.S.A.

EPA Reg. No. 59639-XXX

EPA Est.

059639-00ELU.20220623.IntegoSuiteCanola.NPR.Clean