



**OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION**

WASHINGTON, D.C. 20460

November 05, 2025

Robert L. Hamilton  
robert.hamilton@valent.com  
VALENT U.S.A. LLC

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment - Adjust the restrictions and limitations in the Product Information section for pome and stone fruit; Update tank mix recommendations in the cotton use directions; Include yellow popcorn in the rotational table; Update the warranty statement.  
Product Name: VALOR EZ HERBICIDE 3  
Admin Number: 59639-273  
EPA Receipt Date: 07/30/2025  
Action Case Number: 00666008

Dear Robert L. Hamilton:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have questions, please contact Jeffrey Chang via email at [chang.jeffrey@epa.gov](mailto:chang.jeffrey@epa.gov).

Sincerely,

*Kable Bo Davis*

Kable Bo Davis, Senior Advisor  
FHB, RD  
Office of Pesticide Programs



|             |       |    |           |
|-------------|-------|----|-----------|
| FLUMIOXAZIN | GROUP | 14 | HERBICIDE |
|-------------|-------|----|-----------|

[Note: ***Bracketed Bold italicized text*** is information for the reader and is not part of the label.]

[Bracketed information is optional text].

[Roundup Ready PLUS® Crop Management Solutions]

### VALOR® EZ Herbicide 3

FOR CONTROL AND/OR SUPPRESSION OF CERTAIN WEEDS IN ALFALFA; ARTICHOKE; ASPARAGUS; BRASSICA (HEAD AND STEM); BUSHBERRIES; CACTUS (PRICKLY PEAR); CANEBERRIES; CELERY; CITRUS FRUIT; CLOVER; COTTON; CUCURBIT VEGETABLES; DRY BEANS; FIELD CORN; FIELD PEAS; FLAX; FRUITING VEGETABLES; GARLIC; GRAPE; HOPS; LENTILS; MINT; ONION (DRY BULB); OLIVE; PEANUT; POME FRUIT; POMEGRANATE; POTATO; SOYBEAN; STONE FRUIT; STRAWBERRY; SUGARCANE; SUNFLOWER AND SAFFLOWER; SWEET POTATO; TREE NUTS; WHEAT; NON-BEARING FRUIT TREES; FALLOWBED USE ON TRANSPLANTED MELON, PEPPER, AND TOMATO BEDS; FALLOW LAND AND TO MAINTAIN BARE GROUND ON NON-CROP AREAS OF FARMS; ORCHARDS AND VINEYARDS.

|                         |        |
|-------------------------|--------|
| Active Ingredient       | By Wt  |
| Flumioxazin* .....      | 41.4%  |
| Other Ingredients ..... | 58.6%  |
| Total                   | 100.0% |

\*N-[7-fluoro-3,4-dihydro-3-oxo-4-(prop-2-ynyl)-2H-1,4-benzoxazin-6-yl]cyclohex-1-ene-1,2-dicarboximide

Valor® EZ Herbicide 3 is a suspension concentrate containing 41.4% active ingredient (4.0 lb of flumioxazin per gallon).

**KEEP OUT OF REACH OF CHILDREN**

**SEE [NEXT] [PAGE] [PANEL] [BOOKLET] FOR [ADDITIONAL PRECAUTIONARY STATEMENTS] [AND] [DIRECTIONS FOR USE]**

**[Shake Well Before Using] [Shake Well Before Use] [Shake Before Using] [Shake Before Use] [Mix Thoroughly Before Use] [Always Mix Thoroughly Before Use] [Always Mix Product Thoroughly Before Use] [Agitate Well Before Using] [Shake Well, Agitate or Recirculate Before Use]**

EPA Reg. No. 59639-273

[NET CONTENTS \_\_\_\_]  
[Nonrefillable Container]  
[Refillable Container]

**ACCEPTED**

**11/05/2025**

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 59639-273

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

### PERSONAL PROTECTIVE EQUIPMENT (PPE):

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene ≥ 14 mils, polyvinyl chloride or Viton ≥ 14 mils, shoes and socks.

**For aerial application to sugarcane, mixer/loaders must also wear:** coveralls, chemical resistant apron and chemical resistant boots.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This product is toxic to non-target plants and aquatic invertebrates. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift or runoff may be hazardous to non-target plants and aquatic organisms in neighboring areas. **DO NOT** apply where runoff is likely to occur. **DO NOT** apply when weather conditions favor drift from treated areas. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is toxic to plants and should be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

Under some conditions this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, including no till, limited till and contour plowing; these methods also reduce pesticide run-off. Use of vegetation filter strips along rivers, creeks, streams, wetlands or on the downhill side of fields where run-off could occur will minimize water run-off.

**NON-TARGET ORGANISM ADVISORY:** This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including via liquid spray pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

## **PHYSICAL OR CHEMICAL HAZARDS**

**DO NOT** mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil or water is: coveralls, chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene  $\geq$  14 mils, polyvinyl chloride or Viton  $\geq$  14 mils, shoes plus socks.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries or greenhouses.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift.

**DO NOT** enter or allow others to enter treated areas until sprays have dried.

### Resistance Management

For resistance management, *Valor* EZ Herbicide 3 is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to *Valor* EZ Herbicide 3 and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of *Valor* EZ Herbicide 3 or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Fields should be scouted prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective. Fields should be scouted after application to verify that the treatment was effective. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes or to find out if suspected resistant weeds have been found in their region.
- For further information or to report lack of performance or suspected resistance, contact Valent U.S.A. LLC at 800-6-VALENT (682-5368).

## TABLE OF CONTENTS

|                                                                                                                                    |  |
|------------------------------------------------------------------------------------------------------------------------------------|--|
| PRODUCT INFORMATION.....                                                                                                           |  |
| <i>Valor</i> EZ Herbicide 3 Rate Summary Table .....                                                                               |  |
| Restrictions and Limitations.....                                                                                                  |  |
| Precautions.....                                                                                                                   |  |
| Environmental Conditions and Biological Performance .....                                                                          |  |
| Preemergence Application .....                                                                                                     |  |
| Burndown Application .....                                                                                                         |  |
| Postemergence Application .....                                                                                                    |  |
| Rainfastness .....                                                                                                                 |  |
| Soil Characteristics .....                                                                                                         |  |
| Herbicide Rate - Residual Weed Control .....                                                                                       |  |
| Carrier Volume and Spray Pressure .....                                                                                            |  |
| Preemergence Application .....                                                                                                     |  |
| Burndown Application .....                                                                                                         |  |
| Postemergence Application .....                                                                                                    |  |
| Additives Burndown Application .....                                                                                               |  |
| Jar Test to Determine Compatibility of Adjuvants and <i>Valor</i> EZ Herbicide 3.....                                              |  |
| Sprayer Preparation .....                                                                                                          |  |
| Mixing Instructions.....                                                                                                           |  |
| Sprayer Cleanup.....                                                                                                               |  |
| Application Equipment.....                                                                                                         |  |
| Broadcast Application.....                                                                                                         |  |
| Band Application.....                                                                                                              |  |
| Aerial Application.....                                                                                                            |  |
| Mandatory Spray Drift Requirements .....                                                                                           |  |
| Aerial Application .....                                                                                                           |  |
| Ground Boom Applications .....                                                                                                     |  |
| Boomless Ground Applications.....                                                                                                  |  |
| Spray Drift Advisories .....                                                                                                       |  |
| Importance of Droplet Size .....                                                                                                   |  |
| Boom Height – Ground Boom .....                                                                                                    |  |
| Release Height – Aircraft.....                                                                                                     |  |
| Shielded Sprayers .....                                                                                                            |  |
| Temperature and Humidity .....                                                                                                     |  |
| Temperature Inversions.....                                                                                                        |  |
| Wind.....                                                                                                                          |  |
| Boomless Ground Applications.....                                                                                                  |  |
| Handheld Technology Applications.....                                                                                              |  |
| Chemigation.....                                                                                                                   |  |
| Application with Dry Bulk Fertilizers .....                                                                                        |  |
| Rotational Restrictions.....                                                                                                       |  |
| <br>Table 1. Broadleaf Weeds Controlled by Residual Activity of <i>Valor</i> EZ Herbicide 3.....                                   |  |
| <br>Table 2. Weeds Suppressed by Residual Activity of <i>Valor</i> EZ Herbicide 3 .....                                            |  |
| <br>DIRECTIONS FOR USE IN FALL AND SPRING PREPLANT BURNDOWN AND FALLOW<br>SEEDBED PROGRAMS IN FIELD CORN, PEANUT AND SOYBEAN ..... |  |
| Restrictions and Limitations.....                                                                                                  |  |
| Fall Burndown and Fallow Seedbed Programs .....                                                                                    |  |

Table 3. Weeds Controlled by Fall and Spring Preplant Burndown Programs .....  
Spring Burndown Programs .....

DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN PROGRAMS IN COTTON AND  
SUGARCANE .....

Restrictions and Limitations.....  
Fall Burndown Programs .....  
Spring Burndown Programs .....

DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN PROGRAMS IN RICE,  
SORGHUM, SUNFLOWERS, TOBACCO AND WHEAT (Preplant to Crop) .....  
Restrictions and Limitations.....  
Fall Burndown Programs .....  
Spring Burndown Programs .....

DIRECTIONS FOR USE IN FALL BURNDOWN PROGRAMS IN FIELDS TO BE PLANTED TO  
BARLEY, FIELD PEAS, FLAX, LENTIL, SAFFLOWER, SUNFLOWER AND SPRING WHEAT  
(Preplant to Crop).....  
Restrictions and Limitations.....  
Fall Burndown Programs .....

DIRECTIONS FOR USE IN FALLOW LAND.....

DIRECTIONS FOR USE IN ESTABLISHED ALFALFA.....  
Restrictions and Limitations.....  
Timing to Alfalfa.....  
Timing to Weeds.....

DIRECTIONS FOR USE IN ARTICHOKE .....  
Restrictions and Limitations.....  
Timing to Artichoke.....  
Timing to Weeds.....

DIRECTIONS FOR USE IN ESTABLISHED ASPARAGUS.....  
Restrictions and Limitations.....  
Timing to Asparagus.....  
Timing to Weeds.....

DIRECTIONS FOR USE IN BRASSICA HEAD AND STEM VEGETABLES CROP GROUP 5-16 .....  
Row Middles - Restrictions and Limitations.....  
Precautions.....  
Rate .....  
Timing to Crop.....  
Weed Control and Tank Mixing .....

DIRECTIONS FOR USE IN CACTUS (PRICKLY PEAR).....  
Restrictions and Limitations.....  
Preemergence Application .....  
Postemergence Application.....  
Carrier Volume and Spray Pressure .....  
Banded Application.....

DIRECTIONS FOR USE IN CELERY .....  
Restrictions and Limitations.....  
Precautions.....  
Timing to Celery .....

|                                                                                                                                                                         |  |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
| Timing to Weeds.....                                                                                                                                                    |  |
| DIRECTIONS FOR USE IN ESTABLISHED CLOVER AND CLOVER GROWN FOR SEED .....                                                                                                |  |
| Restrictions and Limitations.....                                                                                                                                       |  |
| Precautions.....                                                                                                                                                        |  |
| Timing to Clover .....                                                                                                                                                  |  |
| Timing to Weeds.....                                                                                                                                                    |  |
| DIRECTIONS FOR USE IN COTTON .....                                                                                                                                      |  |
| Restrictions and Limitations.....                                                                                                                                       |  |
| Environmental Conditions and Biological Performance .....                                                                                                               |  |
| Herbicide Rate.....                                                                                                                                                     |  |
| Table 4. Emerged Broadleaf Weeds Controlled by Hooded, Shielded and Layby Application of <i>Valor</i> EZ Herbicide 3 Tank Mixes with Glyphosate or MSMA in Cotton ..... |  |
| Carrier Volume and Spray Pressure .....                                                                                                                                 |  |
| Additives .....                                                                                                                                                         |  |
| Application Equipment.....                                                                                                                                              |  |
| Timing to Cotton .....                                                                                                                                                  |  |
| Timing to Weeds.....                                                                                                                                                    |  |
| Tank Mixes .....                                                                                                                                                        |  |
| Table 5. Tank Mixes with <i>Valor</i> EZ Herbicide 3 for Hooded, Shielded and/or Layby Use in Cotton .....                                                              |  |
| DIRECTIONS FOR USE IN CUCURBIT VEGETABLES.....                                                                                                                          |  |
| Restrictions and Limitations.....                                                                                                                                       |  |
| Timing to Cucurbit Vegetables .....                                                                                                                                     |  |
| Timing to Weeds.....                                                                                                                                                    |  |
| DIRECTIONS FOR USE IN DRY BEANS .....                                                                                                                                   |  |
| Weed Suppression in Dry Beans and Weed Control in Chickpeas (Garbanzo Beans).....                                                                                       |  |
| Restrictions and Limitations.....                                                                                                                                       |  |
| Timing to Dry Beans and Chickpeas .....                                                                                                                                 |  |
| Timing to Weeds.....                                                                                                                                                    |  |
| Additional Residual Grass Control.....                                                                                                                                  |  |
| Harvest aid - Restrictions and Limitations .....                                                                                                                        |  |
| Timing to Dry Beans and Chickpeas .....                                                                                                                                 |  |
| DIRECTION FOR USE IN FIELD CORN .....                                                                                                                                   |  |
| Restrictions and Limitations.....                                                                                                                                       |  |
| Timing to Field Corn .....                                                                                                                                              |  |
| Burndown Use Directions – For Preplant Application in Field Corn .....                                                                                                  |  |
| Increasing Speed of Glyphosate Burndown Activity.....                                                                                                                   |  |
| Tank Mixes .....                                                                                                                                                        |  |
| Table 6. Tank Mix Partners for Burndown and/or Residual Control of Weeds in Field Corn .....                                                                            |  |
| Tank Mix Restrictions .....                                                                                                                                             |  |
| DIRECTIONS FOR USE IN FIELD PEAS .....                                                                                                                                  |  |
| Weed Control - Restrictions and Limitations .....                                                                                                                       |  |
| Timing to Field Peas.....                                                                                                                                               |  |
| Timing to Weeds.....                                                                                                                                                    |  |
| Additional Residual Grass Control.....                                                                                                                                  |  |
| Harvest Aid - Restrictions and Limitations.....                                                                                                                         |  |
| Timing to Field Peas.....                                                                                                                                               |  |
| DIRECTIONS FOR USE IN FLAX .....                                                                                                                                        |  |
| Harvest Aid - Restrictions and Limitations.....                                                                                                                         |  |
| Timing to Flax .....                                                                                                                                                    |  |

DIRECTIONS FOR USE IN FRUITING VEGETABLES .....  
Restrictions and Limitations.....  
Timing to Fruiting Vegetables .....  
Timing to Weeds.....

DIRECTIONS FOR USE IN GARLIC .....  
Restrictions and Limitations.....  
Timing to Garlic.....  
Timing to Weeds.....

DIRECTIONS FOR USE IN HOPS .....  
Restrictions and Limitations.....  
Timing to Hops for Sucker Control .....  
Timing to Hops for Preemergence Weed Control .....  
Timing to Weeds.....

DIRECTIONS FOR USE IN LENTILS.....  
Harvest Aid - Restrictions and Limitations.....  
Timing to Lentils .....

DIRECTIONS FOR USE IN MINT (Peppermint and Spearmint) .....  
Restrictions and Limitations.....  
Precautions.....  
Timing to Mint .....  
Timing to Weeds.....

Table 7. Weeds Controlled by Residual Activity of *Valor* EZ Herbicide 3.....

DIRECTIONS FOR USE IN ONION (DRY BULB) .....  
Restrictions and Limitations.....  
  
Timing to Onion (dry bulb) .....  
Timing to Weeds.....

DIRECTIONS FOR USE IN PEANUT .....  
Restrictions and Limitations.....  
Wind Management .....  
Timing to Peanuts.....  
Timing to Weeds.....  
Additional Residual Grass Control: Sequential .....  
Additional Residual Grass Control: Tank Mixed .....

DIRECTIONS FOR USE IN POTATO .....  
Restrictions and Limitations.....  
Timing to Potatoes.....  
Timing to Weeds.....

Table 8. Weeds Suppressed by Residual Activity of *Valor* EZ Herbicide 3 at 1.5 oz/A.....

DIRECTIONS FOR USE IN SOYBEAN .....  
Restrictions and Limitations.....  
Precautions.....  
Timing to Soybeans .....  
Timing to Weeds.....  
Increasing Speed of Glyphosate Burndown Activity.....  
Tank Mixes .....

Table 9. Tank Mix Partners for Control of Emerged Weeds in Reduced Tillage Soybeans .....

|                                                                                                                                                                   |  |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
| Additional Residual Broadleaf Control .....                                                                                                                       |  |
| Additional Residual Grass Control.....                                                                                                                            |  |
| Roundup Ready® Program .....                                                                                                                                      |  |
| DIRECTIONS FOR USE IN STRAWBERRY .....                                                                                                                            |  |
| Restrictions and Limitations.....                                                                                                                                 |  |
| Precautions.....                                                                                                                                                  |  |
| Table 10. Weeds Controlled by Preemergence Application of <i>Valor</i> EZ Herbicide 3.....                                                                        |  |
| DIRECTIONS FOR USE IN SUGARCANE .....                                                                                                                             |  |
| Restrictions and Limitations.....                                                                                                                                 |  |
| Timing to Sugarcane .....                                                                                                                                         |  |
| Timing to Weeds.....                                                                                                                                              |  |
| Table 11. Broadleaf Weeds Controlled by Post-Directed or Layby Application of <i>Valor</i> EZ Herbicide 3<br>in Sugarcane.....                                    |  |
| Tank Mixes .....                                                                                                                                                  |  |
| Table 12. Tank Mixes with <i>Valor</i> EZ Herbicide 3 for Post-Directed or Layby Use in Sugarcane .....                                                           |  |
| Additional Preemergence Broadleaf Control .....                                                                                                                   |  |
| Additional Preemergence Grass Control .....                                                                                                                       |  |
| DIRECTIONS FOR USE IN SUNFLOWER AND SAFFLOWER .....                                                                                                               |  |
| Harvest Aid - Restrictions and Limitations.....                                                                                                                   |  |
| Timing to Sunflower and Safflower.....                                                                                                                            |  |
| DIRECTIONS FOR USE IN SWEET POTATO.....                                                                                                                           |  |
| Restrictions and Limitations.....                                                                                                                                 |  |
| Timing to Sweet Potatoes.....                                                                                                                                     |  |
| Timing to Weeds.....                                                                                                                                              |  |
| DIRECTIONS FOR USE IN WHEAT .....                                                                                                                                 |  |
| Restrictions and Limitations.....                                                                                                                                 |  |
| Pre-plant applications, Pre-emergence Weed Control .....                                                                                                          |  |
| Restrictions and Limitations.....                                                                                                                                 |  |
| Burndown Use Directions .....                                                                                                                                     |  |
| Post-Plant, Pre-emergence Weed Control - Restrictions and Limitations .....                                                                                       |  |
| Use Directions, Post-plant, Pre-emergence Weed Control .....                                                                                                      |  |
| Harvest Aid - Restrictions and Limitations.....                                                                                                                   |  |
| Use Directions .....                                                                                                                                              |  |
| Timing to Wheat .....                                                                                                                                             |  |
| DIRECTIONS FOR USE IN BUSHBERRIES, CANEBERRIES, CITRUS FRUIT, GRAPES, OLIVE, POME<br>FRUIT, POMEGRANATE, STONE FRUIT, TREE NUTS AND NON-BEARING FRUIT TREES ..... |  |
| Restrictions and Limitations.....                                                                                                                                 |  |
| Precautions.....                                                                                                                                                  |  |
| Use Precautions for Bushberries.....                                                                                                                              |  |
| Use Precautions for Grapes .....                                                                                                                                  |  |
| Use Precautions for Citrus Fruit, Olive, Pome Fruit, Pomegranate, Stone Fruit and Tree Nuts .....                                                                 |  |
| Use Precautions on Almond and Stone Fruit in Defined Areas of Merced, San Joaquin and<br>Stanislaus Counties of California .....                                  |  |
| Use Precautions for Non-Bearing Fruit Trees .....                                                                                                                 |  |
| Use Directions .....                                                                                                                                              |  |
| Preemergence Application .....                                                                                                                                    |  |
| Postemergence Application.....                                                                                                                                    |  |
| Carrier Volume and Spray Pressure .....                                                                                                                           |  |
| Banded Application.....                                                                                                                                           |  |

|                                                                                                      |  |
|------------------------------------------------------------------------------------------------------|--|
| Table 13. Weeds Controlled by Postemergence Activity of <i>Valor</i> EZ Herbicide 3 Tank Mixes ..... |  |
| Additional Residual Weed Control .....                                                               |  |

|                                                                          |  |
|--------------------------------------------------------------------------|--|
| FALLOWBED USE ON TRANSPLANTED MELON, PEPPER AND TOMATO BEDS .....        |  |
| Restrictions and Limitations .....                                       |  |
| Use for Preemergence Fallowbed Weed Control Prior to Transplanting ..... |  |

|                                                                                                    |  |
|----------------------------------------------------------------------------------------------------|--|
| DIRECTIONS FOR USE TO MAINTAIN BARE GROUND ON NON-CROP AREAS OF FARMS, ORCHARDS OR VINEYARDS ..... |  |
| Restrictions and Limitations .....                                                                 |  |
| Preemergence Application .....                                                                     |  |
| Postemergence Application .....                                                                    |  |

|                                                                              |  |
|------------------------------------------------------------------------------|--|
| Table 14. Tank Mix Combinations to Maintain Bare Ground Non-Crop Areas ..... |  |
| STORAGE AND DISPOSAL .....                                                   |  |

## PRODUCT INFORMATION

- *Valor* EZ Herbicide 3 provides residual control of susceptible weeds.
- *Valor* EZ Herbicide 3 provides additional burndown activity when used as part of a burndown program.
- *Valor* EZ Herbicide 3 can be applied as part of a fall burndown program for control of susceptible winter annuals.
- *Valor* EZ Herbicide 3 can be applied with a hooded or shielded sprayer, as well as part of a layby application, in selected crops for postemergence weed control as well as residual control of susceptible weeds.
- *Valor* EZ Herbicide 3 can be used on farms, orchards and vineyards for non-selective vegetation control to maintain bare ground non-crop areas that must be kept weed free.
- ***Valor* EZ Herbicide 3, when applied according to label use directions, will control the weeds claimed in crop specific use directions. This label makes no claims concerning control of other weed species. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.**

| <b><i>Valor</i> EZ Herbicide 3 Rate Summary</b> |                              |
|-------------------------------------------------|------------------------------|
| <b>FL OZ of <i>Valor</i> EZ Herbicide 3</b>     | <b>Pounds of Flumioxazin</b> |
| 0.5                                             | 0.016                        |
| 1                                               | 0.032                        |
| 1.5                                             | 0.047                        |
| 2                                               | 0.063                        |
| 2.5                                             | 0.080                        |
| 3                                               | 0.094                        |
| 4                                               | 0.125                        |
| 6                                               | 0.188                        |
| 8                                               | 0.250                        |
| 12                                              | 0.375                        |
| 24                                              | 0.750                        |

## AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making decisions. Where states have more stringent regulations, they must be observed.

## RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply this product when weather conditions favor spray drift from treated areas.
- **DO NOT** apply during low-level inversion conditions, including fog.
- **DO NOT** apply to frozen or snow-covered soil.
- **DO NOT** apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation.
- **DO NOT** apply within 300 feet of non-dormant pome or stone fruit, or 300 yards of non-dormant pears.
- **DO NOT** apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be applied immediately after application.

## PRECAUTIONS

- When applying by air, observe drift management restrictions and precautions listed under "AERIAL APPLICATION".
- Mechanical incorporation into the soil will reduce residual weed control.
- Apply post directed and layby applications of *Valor* EZ Herbicide 3 only to healthy growing crops.

**Before using spray equipment to apply other products to crop foliage follow cleanout procedures identified in this label. See “SPRAYER CLEANUP” for more information.**

## **ENVIRONMENTAL CONDITIONS AND BIOLOGICAL PERFORMANCE**

### **Preemergence Application (Conventional Tillage)**

**Important:** Crop injury may occur from applications made to poorly drained soils and/or applications made under cool, wet conditions. Risk of crop injury can be minimized by using on well drained soils, planting at least 1.5 inches deep, using high quality seed and completely covering seeds with soil prior to preemergence applications. Treated soil that is splashed onto newly emerged crops may result in temporary crop injury.

Moisture is necessary to activate *Valor* EZ Herbicide 3 in soil for residual weed control. Dry weather following applications of *Valor* EZ Herbicide 3 may reduce effectiveness. However, when adequate moisture is received after dry conditions, *Valor* EZ Herbicide 3 will control susceptible germinating weeds. *Valor* EZ Herbicide 3 may not control weeds that germinate after application but before an activating rainfall/irrigation or weeds that germinate through cracks resulting from dry soil.

When adequate moisture is not received after a *Valor* EZ Herbicide 3 application, weed control may be improved by irrigation with at least 1/4 inch of water. If emerged weeds are controlled by cultivation, residual weed control will be reduced.

### **Burndown Application**

For best results, apply *Valor* EZ Herbicide 3 as part of a burndown program to actively growing weeds. Applying *Valor* EZ Herbicide 3 under conditions that do not promote active weed growth will reduce herbicide effectiveness. **DO NOT** apply *Valor* EZ Herbicide 3 when weeds are under stress due to drought, excessive water, extremes in temperature, disease or low humidity. Weeds under stress tend to become less susceptible to herbicidal action. *Valor* EZ Herbicide 3 is most effective when applied under warm sunny conditions.

Reduced residual weed control may occur when burndown applications are made to fields where heavy crop and/or weed residue exist.

### **Postemergence Application**

Only apply *Valor* EZ Herbicide 3 to healthy crops labeled for postemergence use. **DO NOT** apply *Valor* EZ Herbicide 3 to crops that have been weakened by disease, drought, flooding, excessive fertilization, soil salts, previously applied pesticides, nematodes, insects or winter injury.

### **Rainfastness**

*Valor* EZ Herbicide 3 is rainfast one hour after application. **DO NOT** make applications if rain is expected within one hour of application or postemergence efficacy may be reduced.

### **Soil Characteristics**

Application of *Valor* EZ Herbicide 3 to soils with high organic matter and/or high clay content may require higher dosages than soils with low organic matter and/or low clay content. Application to cloddy seedbeds can result in reduced weed control.

## **HERBICIDE RATE**

### **Residual Weed Control (Including Preemergence Applications or Applications as Part of a Fall or Spring Burndown and Fallow Seedbed Program)**

Based upon soil characteristics (organic matter content and texture), the most difficult to control weed species being targeted, and the crop being grown, select the proper *Valor* EZ Herbicide 3 dosage from the rate range tables contained in this label.

**CARRIER VOLUME AND SPRAY PRESSURE** (Ground Equipment only. See Information for Aerial Equipment under “AERIAL APPLICATION”.)

### **Preemergence Application (Conventional Tillage)**

To ensure uniform coverage, use 10 to 30 gal of spray solution per acre for conventional tillage applications. Nozzle selection must meet manufacturer's gallonage and pressure guidelines for preemergence herbicide application.

### **Burndown Application (Prior to Crop Emergence)**

To ensure thorough coverage in burndown applications, use 15 to 60 gallons spray solution per acre. Use 20 to 60 gallons per acre if dense vegetation or heavy crop residue is present. Nozzle selection must meet manufacturer's gallonage and pressure guidelines for postemergence herbicide application. **DO NOT** use flood jet nozzles.

### **Postemergence Application (Emerged Crop)**

Check use directions for specific crops in which *Valor* EZ Herbicide 3 can be applied postemergence. To ensure thorough coverage in burndown applications, use a minimum of 15 gallons spray solution per acre. Use a minimum of 20 gallons per acre if dense vegetation or heavy crop residue is present. Nozzle selection must meet manufacturer's gallonage and pressure guidelines for postemergence herbicide application.

## **ADDITIVES**

### **Burndown Application (Prior to Crop Emergence)**

Postemergence control of weeds from *Valor* EZ Herbicide 3 tank mixes will require the addition of an agronomically approved adjuvant to the spray mixture. When an adjuvant is to be used with *Valor* EZ Herbicide 3, Valent recommends the use of a Chemical Producers and Distributors Association certified adjuvant. Either a crop oil concentrate or methylated seed oil which contains at least 15% emulsifiers and 80% oil or a non-ionic surfactant at 0.25% v/v, may be used when applying *Valor* EZ Herbicide 3 as part of a burndown program. Some tank mix partners, for example Roundup Power Max®, are formulated with sufficient adjuvants and do not require the addition of a crop oil concentrate, methylated seed oil or non-ionic surfactant when tank mixed with *Valor* EZ Herbicide 3. The addition of a crop oil concentrate or methylated seed oil may increase the burndown activity on certain weeds including cutleaf evening primrose and Carolina geranium. Verify mixing compatibility qualities by a jar test.

A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lb/A or a 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to the spray mixture along with either a crop oil concentrate, methylated seed oil or non-ionic surfactant to enhance weed control. The addition of a nitrogen source does not replace the need for a crop oil concentrate, a methylated seed oil or a non-ionic surfactant.

### **JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND VALOR EZ HERBICIDE 3**

When using *Valor* EZ Herbicide 3 and an adjuvant, including in stale seed bed, layby, hooded/shielded or reduced tillage situations, perform a jar test before mixing commercial quantities of *Valor* EZ Herbicide 3, when using *Valor* EZ Herbicide 3 for the first time, when using new adjuvants or when a new water source is being used.

1. Add 1 pt of the water to a quart jar. Use water from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add 1 g of *Valor* EZ Herbicide 3 to the quart jar for every 3 fl oz (0.094 lb ai) of *Valor* EZ Herbicide 3 per acre being applied (4 g if 12 fl oz/A (0.375 lb ai) is the desired *Valor* EZ Herbicide 3 rate), gently mix until product goes into suspension.
3. Add 60 ml (4 Tbsp or 2 fl oz) of the crop oil or methylated seed oil to the quart jar or 1 ml of non-ionic surfactant if it is being used in place of oil, gently mix.
4. If nitrogen is being used, add 16 ml (1 Tbsp or 0.5 oz) of the 28 to 32% nitrogen source to the quart jar. If ammonium sulfate is being used, add 19 g AMS to the quart jar in place of the 28 to 32% nitrogen.
5. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
6. An ideal tank mix combination will be uniform and free of suspended particles. Question the choice of adjuvant if any of the following conditions are observed:
  - a) Layer of oil or globules on the mixture's surface.
  - b) Flocculation: fine particles in suspension or as a layer on the bottom of the jar.
  - c) Clabbering: thickening texture (coagulated) like gelatin.

## SPRAYER PREPARATION

Before applying *Valor* EZ Herbicide 3, start with clean, well maintained application equipment. The spray tank, as well as all hoses and booms, must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. Some pesticides, including but not limited to, the sulfonylurea and phenoxy herbicides, (i.e., Classic® and 2,4-D respectively) are active at very small amounts and can cause crop injury when applied to susceptible crops. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply *Valor* EZ Herbicide 3. If two or more products were tank mixed prior to *Valor* EZ Herbicide 3 application, follow the most restrictive cleanup procedure.

## MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. If a drift retardant is to be used, add 10 lb of spray grade ammonium sulfate per 100 gallon of spray solution.
3. While agitating, slowly add *Valor* EZ Herbicide 3 to the spray tank. Agitation creates a rippling or rolling action on the water surface.
4. If tank mixing *Valor* EZ Herbicide 3 with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
5. Add any required adjuvants.
6. Fill spray tank to desired level with water. **Continue agitation until all spray solution has been applied.**
7. Mix only the amount of spray solution that can be applied the day of mixing. Apply *Valor* EZ Herbicide 3 within 6 hours of mixing.

## SPRAYER CLEANUP

Spray equipment, including mixing vessels and nurse tanks, must be cleaned each day following *Valor* EZ Herbicide 3 application. After *Valor* EZ Herbicide 3 is applied, the following steps must be used to clean the spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
3. Top off tank, add 1 gallon of 3% household ammonia (or equivalent) for every 100 gallons of water, circulate through sprayer for 5 minutes, and then flush all hoses, booms, screens and nozzles for a minimum of 15 minutes. If diaphragms are being used on the spray boom, loosen diaphragms before flushing the spray system, allowing cleaning solution to spray through the open diaphragm. If spray lines have any end caps, they must be loosened before flushing the system, allowing cleaning solution to spray through the loosened caps. To enhance removal of *Valor* EZ Herbicide 3 from the spray system, add a tank cleaner for example "Valent Tank Cleaner" from Valent U.S.A. LLC, in place of ammonia and allow the cleaning solution to remain in the pressurized spray system (spray tank, hoses and boom) overnight before flushing the system for a minimum of 15 minutes.
4. Drain tank completely.
5. Add enough clean water to the spray tank to allow all hoses, booms, screens and nozzles to be flushed for 2 minutes.
6. Remove all nozzles and screens and rinse them in clean water.

Thoroughly clean spray equipment, including all tanks, hoses, booms, screens and nozzles, cleaned before it is used to apply postemergence pesticides. Equipment with *Valor* EZ Herbicide 3 residue remaining in the system may result in crop injury to the subsequently treated crop.

## APPLICATION EQUIPMENT

Ensure application equipment is clean and in good repair, nozzles are uniformly spaced on the boom and frequently checked for accuracy.

### BROADCAST APPLICATION

Apply *Valor* EZ Herbicide 3, and *Valor* EZ Herbicide 3 tank mixes, with ground equipment using standard commercial sprayers equipped with flat fan or flood nozzles (preemergence applications only) designed to deliver the desired spray pressure and spray volume.

### BAND APPLICATION

When banding, use proportionately less water and *Valor* EZ Herbicide 3 per acre. The rate of *Valor* EZ Herbicide 3 required per acre, when applied as a banded application, can be calculated with the following formula:

|                                                  |   |                                                                  |   |                         |
|--------------------------------------------------|---|------------------------------------------------------------------|---|-------------------------|
| Amount Needed per Acre for<br>Banded Application | = | $\frac{\text{Band Width in Inches}}{\text{Row Width in Inches}}$ | X | Rate per Broadcast Acre |
|--------------------------------------------------|---|------------------------------------------------------------------|---|-------------------------|

### AERIAL APPLICATION

- **Carrier Volume and Spray Pressure:** When used as part of a burndown weed control program, apply *Valor* EZ Herbicide 3 in 7 to 10 gallons of water per acre. Application at less than 7 gallons per acre may provide inadequate control. When used for preemergence weed control, apply *Valor* EZ Herbicide 3 in 5 to 10 gallons of water per acre. The higher gallonage applications afford more consistent weed control. **DO NOT** exceed the nozzle manufacturer's specified pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Adjuvants and Drift Control Additives:** Refer to tank mix partner's label for adjuvants election. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

## **MANDATORY SPRAY DRIFT REQUIREMENTS**

### **Aerial Applications**

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy unless a greater application height is necessary for pilot safety.
- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- If the windspeed is 10 miles per hour or less, applicators must use 1/2 swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 miles per hour, applicators must use 3/4 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- **DO NOT** apply during temperature inversions.
- **DO NOT** apply this product by air within 40 ft of non-target plants including non-target crops.
- **DO NOT** apply this product by air within 100 ft of emerged cotton crops.
- **DO NOT** apply this product by air within 40 ft of streams, wetlands, marshes, ponds, lakes and reservoirs.

### **Ground Boom Applications**

- User must only apply with the release height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

### **Boomless Ground Applications**

- Applicators must select nozzle and pressure that deliver Medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

## **SPRAY DRIFT ADVISORIES**

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

## **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### **Controlling Droplet Size - Ground Boom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **Controlling Droplet Size - Aircraft**

- Adjust Nozzles - Follow nozzle manufacturers' specifications for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### **BOOM HEIGHT - Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **RELEASE HEIGHT - Aircraft**

Higher release heights increase the potential for spray drift.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### **Boomless Ground Applications**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

### **Handheld Technology Applications**

Take precautions to minimize spray drift.

### **CHEMIGATION**

Follow all label directions for crops regarding rates, timing of application, special instructions and precautions.

Apply this product only through center pivot systems. End guns must be turned off due to uneven application. Restriction: **DO NOT** apply this product through any other type of irrigation system.

Crop injury, lack of efficacy or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

The system must be properly calibrated (with water only) to ensure that the amount of *Valor* EZ Herbicide 3 applied corresponds to the specified rate.

Apply *Valor* EZ Herbicide 3 in 1/2 to 3/4 inches of water during the first sprinkler set. Allow time for all lines to flush the herbicide through all nozzles before turning off irrigation water. To ensure the lines are flushed and free of remaining herbicide, a dye indicator may be injected into the lines to mark the end of the application period. Once chemigation has begun, the run must be completed to ensure no product is left in the system.

If you have any questions about calibration, contact your State Extension Service Specialist, equipment manufacturers or other experts.

### Special Precautions for Chemigation

1. **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
2. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.
3. The system must be free of leaks and clogged nozzles.
4. The pesticide must be supplied continuously for the duration of the aqueous application. An uneven application may cause injury to the crop or poor weed control.
5. Agitation must be maintained in the nurse tank.
6. The sprinkler chemigation system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
7. The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in the case where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, for example a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with the pesticides and capable of being fitted with a system interlock.
12. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

### Chemigation Systems Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption, if such a system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to the public water system must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, discharge the water from the public water system into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. All chemigation systems connected to the public water system must also follow restrictions listed in the preceding section titled **"Special Precautions for Chemigation"**.

### APPLICATION WITH DRY BULK FERTILIZERS

Dry bulk fertilizer may be impregnated or coated with *Valor* EZ Herbicide 3. Application of dry bulk fertilizer with *Valor* EZ Herbicide 3 provides weed control equal to, or slightly below, the same rate of *Valor* EZ Herbicide 3 applied in liquid carriers, due to better coverage with application via spray equipment. Follow label directions for *Valor* EZ Herbicide 3 regarding rates, special instructions, cautions and special precautions. Apply 400 to 700 lb of the fertilizer/herbicide mixture per acre to obtain adequate soil coverage. Apply the mixture to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury and to obtain uniform weed control.

**DO NOT** use ammonium nitrate and/or limestone as the sole source of fertilizer, as the *Valor* EZ Herbicide 3 may not adhere to these materials.

Compliance with all Federal and State regulations relating to blending pesticide mixtures with dry bulk fertilizer, registrations, labeling and application are the responsibility of the individual and/or company offering the fertilizer and *Valor* EZ Herbicide 3 mixture for sale.

*Valor* EZ Herbicide 3 must be premixed with water to form a slurry prior to impregnation on dry bulk fertilizer. For best results, use a minimum of 1 pt of water for each 2 fl oz of *Valor* EZ Herbicide 3. Use a minimum of 6 pt of the *Valor* EZ Herbicide 3 slurry to impregnate 2000 lb of the fertilizer for uniform coverage of the fertilizer. Closed drum, belt, ribbon or other commonly used dry bulk blenders may be used.

The amount of *Valor* EZ Herbicide 3 required can be calculated with the following formula:

|                                                                         |   |                                                            |   |      |   |                                  |
|-------------------------------------------------------------------------|---|------------------------------------------------------------|---|------|---|----------------------------------|
| fluid ounces of<br><i>Valor</i> EZ Herbicide 3<br>per ton of fertilizer | = | fluid ounces of<br><i>Valor</i> EZ Herbicide 3 per<br>acre | X | 2000 | ÷ | pounds of fertilizer<br>per acre |
|-------------------------------------------------------------------------|---|------------------------------------------------------------|---|------|---|----------------------------------|

Thoroughly clean dry fertilizer blending equipment after *Valor* EZ Herbicide 3 has been placed in the system to avoid injury to sensitive crops that may be treated with fertilizers blended after the equipment has been used for *Valor* EZ Herbicide 3. Rinse the sides of the blender and the herbicide tank with water. Then impregnate the rinsate onto a load of dry fertilizer intended for an approved crop. Use a maximum rate of 1 gallon of rinsate per ton of fertilizer. Follow with 1 to 2 loads of unimpregnated fertilizer in the blender before switching herbicides.

## ROTATIONAL RESTRICTIONS

The following rotational crops may be planted after applying *Valor* EZ Herbicide 3 at the listed rate. Planting earlier than the specified rotational interval may result in crop injury.

- **DO NOT plant any crop, except corn (field), cotton, peanut, soybean, sugarcane and sweet potato earlier than 30 days after applying *Valor* EZ Herbicide 3.**

| Application Rates                       | Crops                                                                                                                                                                              | Rotation Intervals                                                                    |
|-----------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| 1 fl oz/A (0.032 lb ai)                 | Cotton (no-till or strip-till only)                                                                                                                                                | 14 days <sup>(1)</sup>                                                                |
| 1.5 to 2 fl oz/A (0.047 to 0.063 lb ai) | Cotton (no-till or strip-till only)                                                                                                                                                | 21 days <sup>(1)</sup>                                                                |
| 2 fl oz/A (0.063 lb ai) or less         | Peanut, Soybean, Sugarcane and Sweet Potato                                                                                                                                        | immediately                                                                           |
|                                         | Field Corn (minimum and no-till)                                                                                                                                                   | 7 days                                                                                |
|                                         | Cotton and Field Corn (conventional tillage), Rice, Sorghum, Sunflower, Tobacco and Wheat                                                                                          | 30 days <sup>(1)</sup>                                                                |
|                                         | Barley, Dry and Snap Beans, Flax, Peas, Rye, Safflower, Sweet Corn, and Yellow Popcorn                                                                                             | 3 months                                                                              |
|                                         | Alfalfa, Canola, Clover, Oats, Potato, Sugar Beet and all other crops not listed <sup>(2)</sup>                                                                                    | 4 months if soil is tilled prior to planting<br>8 months if no tillage is performed   |
|                                         | Lentil                                                                                                                                                                             | 6 months                                                                              |
| Up to 3 fl oz/A (0.094 lb ai)           | Peanut, Soybean, Sugarcane and Sweet Potato                                                                                                                                        | immediately                                                                           |
|                                         | Field Corn (minimum and no-till)                                                                                                                                                   | 14 days                                                                               |
|                                         | Field Corn (conventional tillage) and Sorghum                                                                                                                                      | 30 days <sup>(1)</sup>                                                                |
|                                         | Cotton, Rice, Sunflower, Tobacco and Wheat                                                                                                                                         | 2 months <sup>(1)</sup>                                                               |
|                                         | Barley, Dry and Snap Beans, Flax, Pea, Rye, Safflower, Sweet Corn, and Yellow Popcorn                                                                                              | 4 months                                                                              |
|                                         | Alfalfa, Clover, Oats, Potato, Sugar Beet                                                                                                                                          | 5 months if soil is tilled prior to planting<br>10 months if no tillage is performed  |
|                                         | Canola and all other crops not listed <sup>(2)</sup>                                                                                                                               | 6 months if soil is tilled prior to planting<br>12 months if no tillage is performed  |
|                                         | Lentil                                                                                                                                                                             | 7 months                                                                              |
| Up to 4 fl oz/A (0.125 lb ai)           | Sugarcane                                                                                                                                                                          | Immediately                                                                           |
|                                         | Alfalfa, Canola, Potato, Sugar Beet and all other crops not listed <sup>(2)</sup>                                                                                                  | 6 months if soil is tilled prior to planting<br>12 months if no tillage is performed  |
|                                         | Cotton, Field Corn, Peanut, Rice, Sorghum, Soybean, Sunflower, Tobacco and Wheat                                                                                                   | 4 months                                                                              |
|                                         | Transplanted on raised beds only: melon, pepper and tomato                                                                                                                         | 2 months (if the top 4 inches of the beds have been removed)                          |
| 6 to 12 fl oz/A (0.188 to 0.375 lb ai)  | Cotton, Field Corn, Peanut, Rice, Sorghum, Soybean, Sunflower, Tobacco and Wheat                                                                                                   | 9 months                                                                              |
|                                         | Alfalfa, Canola, Sugar Beet and all other crops not listed <sup>(2)</sup><br>Trees can be transplanted 2 months after an application of <i>Valor</i> EZ Herbicide 3 <sup>(3)</sup> | 12 months if soil is tilled prior to planting<br>18 months if no tillage is performed |

<sup>(1)</sup> At least one inch of rainfall/irrigation must occur between application and planting or crop injury may occur.

<sup>(2)</sup> Successful soil bioassay must be performed prior to planting these crops.

<sup>(3)</sup> Transplanted apple, apricot, avocado, bushberries (including blueberry), cherry, fig, grape, grapefruit, lemon, nectarine, olive, orange, peach, pear, plum (including dried plum), tangerine and tree nuts (including pistachio) can be planted 2 months after a *Valor* EZ Herbicide 3 application of 2 to 12/A.

**Table 1. Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3**

| <b>BROADLEAF WEED SPECIES</b> |                                                 |                       |                  |                            |
|-------------------------------|-------------------------------------------------|-----------------------|------------------|----------------------------|
| <b>Section A</b>              |                                                 |                       |                  |                            |
| <b>Common Name</b>            | <b>Scientific Name</b>                          | <b>Organic Matter</b> | <b>Soil Type</b> | <b>Application Rate</b>    |
| Carpetweed                    | <i>Mollugo verticillata</i>                     | Up to 5%              | All Soil Types   | 2 fl oz/A<br>(0.063 lb ai) |
| Chickweeds                    |                                                 |                       |                  |                            |
| Common                        | <i>Stellaria media</i>                          |                       |                  |                            |
| Mouseear                      | <i>Cerastium vulgatum</i>                       |                       |                  |                            |
| Dandelion                     | <i>Taraxacum officinale</i>                     |                       |                  |                            |
| Eclipta                       | <i>Eclipta prostrate</i>                        |                       |                  |                            |
| Evening primrose, Cutleaf     | <i>Oenothera laciniata</i>                      |                       |                  |                            |
| Field Pennycress[*]           | <i>Thlaspi arvense</i>                          |                       |                  |                            |
| Florida Pusley                | <i>Richardia scabra</i>                         |                       |                  |                            |
| Henbit                        | <i>Lamium amplexicaule</i>                      |                       |                  |                            |
| Lambsquarters, Common         | <i>Chenopodium album</i>                        |                       |                  |                            |
| Little Mallow                 | <i>Malva parviflora</i>                         |                       |                  |                            |
| Marestail/Horseweed           | <i>Conyza canadensis</i>                        |                       |                  |                            |
| Mayweed/False Chamomile       | <i>Matricaria maritime</i>                      |                       |                  |                            |
| Nightshades                   |                                                 |                       |                  |                            |
| Black                         | <i>Solanum nigrum</i>                           |                       |                  |                            |
| Eastern Black                 | <i>Solanum ptycanthum</i>                       |                       |                  |                            |
| Hairy                         | <i>Solanum sarrachoides</i>                     |                       |                  |                            |
| Pigweeds                      |                                                 |                       |                  |                            |
| Redroot                       | <i>Amaranthus retroflexus</i>                   |                       |                  |                            |
| Smooth                        | <i>Amaranthus hybridus</i>                      |                       |                  |                            |
| Spiny Amaranth                | <i>Amaranthus spinosus</i>                      |                       |                  |                            |
| Tumble                        | <i>Amaranthus albus</i>                         |                       |                  |                            |
| Prickly Lettuce               | <i>Lactuca serriola</i>                         |                       |                  |                            |
| Prickly Sida (Teaweed)        | <i>Sida spinosa</i>                             |                       |                  |                            |
| Puncturevine                  | <i>Tribulus terrestris</i>                      |                       |                  |                            |
| Purslane, Common              | <i>Portulaca oleracea</i>                       |                       |                  |                            |
| Radish, Wild                  | <i>Raphanus raphanistrum</i>                    |                       |                  |                            |
| Redmaids                      | <i>Calandrinia ciliata</i> var <i>menziesii</i> |                       |                  |                            |
| Shepherd's-purse              | <i>Capsella bursa-pastoris</i>                  |                       |                  |                            |
| Smallflower Morningglory      | <i>Jacquemontia tamnifolia</i>                  |                       |                  |                            |
| Sowthistle, Prickly[*]        | <i>Sonchus asper</i>                            |                       |                  |                            |
| Spotted Spurge                | <i>Euphorbia maculate</i>                       |                       |                  |                            |
| Venice Mallow                 | <i>Hibiscus trionum</i>                         |                       |                  |                            |

[\*Not for use in California]

continued

**Table 1. Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3 (continued)**

| <b>Section B</b>                           |                                                   |                       |                                                                                                           |                                                                                                                                                                        |
|--------------------------------------------|---------------------------------------------------|-----------------------|-----------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>All weeds listed in Section A plus:</b> |                                                   |                       |                                                                                                           |                                                                                                                                                                        |
| <b>Common Name</b>                         | <b>Scientific Name</b>                            | <b>Organic Matter</b> | <b>Soil Type</b>                                                                                          | <b>Application Rates<sup>[(2)]</sup></b>                                                                                                                               |
| Coffee Senna                               | <i>Cassia occidentalis</i>                        | Up to 3%              | All Soil Types                                                                                            | 2 fl oz/A (0.063 lb ai)<br>Cotton and Dry Bean<br>2.5 fl oz/A (0.080 lb ai)<br>Field Corn and Soybean<br>3 fl oz/A (0.094 lb ai)<br>Peanut and all other               |
| Common Ragweed <sup>(1)</sup>              | <i>Ambrosia artemisiifolia</i>                    |                       |                                                                                                           |                                                                                                                                                                        |
| False Chamomile[*]                         | <i>Tripleurospermum maritima</i>                  |                       |                                                                                                           |                                                                                                                                                                        |
| Florida Beggarweed                         | <i>Desmodium tortuosum</i>                        |                       |                                                                                                           |                                                                                                                                                                        |
| Golden Crownbeard                          | <i>Verbesina encelioides</i>                      |                       |                                                                                                           |                                                                                                                                                                        |
| Hairy Indigo                               | <i>Indigofera hirsute</i>                         |                       |                                                                                                           |                                                                                                                                                                        |
| Hemp Sesbania                              | <i>Sesbania exaltata</i>                          | 3 to 5%               | Coarse and Medium Soils:<br>(sandy loam, loamy sand, loamy, silt-loam, silt, sandy clay, sandy clay loam) | 2 fl oz/A (0.063 lb ai)<br>Cotton and Dry Bean<br>2.5 fl oz/A (0.080 lb ai)<br>Field Corn and Soybean<br>3 fl oz/A (0.094 lb ai)<br>Peanut and all other labeled crops |
| Jimsonweed                                 | <i>Datura stramonium</i>                          |                       |                                                                                                           |                                                                                                                                                                        |
| Kochia                                     | <i>Kochia scoparia</i>                            |                       |                                                                                                           |                                                                                                                                                                        |
| London Rocket[*]                           | <i>Sisymbrium irio</i>                            |                       |                                                                                                           |                                                                                                                                                                        |
| Morningglories <sup>(3)</sup>              |                                                   |                       |                                                                                                           |                                                                                                                                                                        |
| Entireleaf                                 | <i>Ipomoea hederacea</i> var. <i>integruscula</i> |                       |                                                                                                           |                                                                                                                                                                        |
| Ivyleaf                                    | <i>Ipomoea hederacea</i>                          |                       |                                                                                                           |                                                                                                                                                                        |
| Red/Scarlet                                | <i>Ipomoea coccinea</i>                           |                       |                                                                                                           |                                                                                                                                                                        |
| Tall                                       | <i>Ipomoea purpurea</i>                           |                       |                                                                                                           |                                                                                                                                                                        |
| Mustard, Wild                              | <i>Brassica kaber</i>                             |                       |                                                                                                           |                                                                                                                                                                        |
| Palmer Amaranth                            | <i>Amaranthus palmeri</i>                         |                       |                                                                                                           |                                                                                                                                                                        |
| Spurred Anoda                              | <i>Anoda cristata</i>                             |                       |                                                                                                           |                                                                                                                                                                        |
| Tropic Croton                              | <i>Croton glandulosus</i>                         |                       |                                                                                                           |                                                                                                                                                                        |
| Waterhemp <sup>(1)</sup>                   |                                                   |                       | Fine Soils:<br>(silty clay, silty clay loam, clay, clay loam)                                             | 2 fl oz/A (0.063 lb ai)<br>Cotton and Dry Bean<br>3 fl oz/A (0.094 lb ai) Field Corn, Peanut, Soybean and all other labeled crops                                      |
| Common                                     | <i>Amaranthus rudis</i>                           |                       |                                                                                                           |                                                                                                                                                                        |
| Tall                                       | <i>Amaranthus tuberculatus</i>                    |                       |                                                                                                           |                                                                                                                                                                        |
| Wild Poinsettia                            | <i>Euphorbia heterophylla</i>                     |                       |                                                                                                           |                                                                                                                                                                        |
| Yellow Rocket[*]                           | <i>Barbarea vulgaris</i>                          |                       |                                                                                                           |                                                                                                                                                                        |

[\*Not for use in California.]

<sup>(1)</sup>A postemergence herbicide, including Cobra® Herbicide, Phoenix™ Herbicide glyphosate (Roundup Ready® soybeans only) may be needed following a preemergence application of Valor EZ Herbicide 3 to adequately control common ragweed or waterhemp in soybean fields with heavy pressure.

<sup>(2)</sup>Valor EZ Herbicide 3 will provide residual control of these weeds at 2 fl oz/A (0.063 lb ai) when applied under a cotton canopy.]

<sup>(3)</sup>Morningglory species are not adequately controlled on fine soils or soils with greater than 3% organic matter.

**Table 2. Weeds Suppressed by Residual Activity of Valor EZ Herbicide 3 Application Rates**

| Broadleaf Weed Species  |                                |                |                                            |
|-------------------------|--------------------------------|----------------|--------------------------------------------|
| Common Name             | Scientific Name                | Organic Matter | Application Rates                          |
| Bristly Starbur         | <i>Acanthospermum hispidum</i> | Up to 5%       | 2 to 3 fl oz/A<br>(0.063 to 0.094 lb ai)   |
| Copperleaf, Hophornbeam | <i>Acalypha ostryifolia</i>    |                |                                            |
| Ragweed, Giant          | <i>Ambrosia trifida</i>        |                |                                            |
| Russian Thistle         | <i>Salsola iberica</i>         |                |                                            |
| Smartweeds              |                                |                |                                            |
| Ladysthumb              | <i>Polygonum persicaria</i>    |                |                                            |
| Pennsylvania            | <i>Polygonum pensylvanicum</i> |                |                                            |
| Smellmelon[*]           | <i>Cucumis melo</i>            |                |                                            |
| Velvetleaf              | <i>Abutilon theophrasti</i>    |                |                                            |
| Wild Buckwheat          | <i>Polygonum convolvulus</i>   |                |                                            |
| Wormwood, Biennial      | <i>Artemisia biennis</i>       |                |                                            |
| Grass Weed Species      |                                |                |                                            |
| Barnyardgrass           | <i>Echinochloa crus-galli</i>  |                |                                            |
| Bluegrass, Annual       | <i>Poa annua</i>               |                |                                            |
| Crabgrass, Large        | <i>Digitaria sanguinalis</i>   |                |                                            |
| Foxtail, Giant          | <i>Setaria faberi</i>          |                |                                            |
| Goosegrass              | <i>Eleusine indica</i>         |                |                                            |
| Lovegrass, California   | <i>Eragrostis diffusa</i>      |                |                                            |
| Panicums                |                                |                |                                            |
| Fall                    | <i>Panicum dichotomiflorum</i> |                |                                            |
| Texas                   | <i>Panicum texanum</i>         |                |                                            |
| Ryegrass, Italian[*]    | <i>Lolium multiflorum</i>      |                |                                            |
| Signalgrass, Broadleaf  | <i>Brachiaria platyphylla</i>  |                |                                            |
| Cheat                   | <i>Bromus secalinus</i>        | Up to 5%       | 1.5 to 3 fl oz/A<br>(0.047 to 0.094 lb ai) |
| Downy Brome[*]          | <i>Bromus tectorum</i>         |                |                                            |

[\*Not for use in California]

**DIRECTIONS FOR USE IN FALL AND SPRING PREPLANT BURNDOWN AND FALLOW  
SEEDBED PROGRAMS IN FIELD CORN, PEANUT AND SOYBEAN  
(Preemergence to Crop)**

[For Use in Arizona, California and Hawaii Only]

**RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply to frozen or snow covered soil.
- **DO NOT** perform any tillage operation after application or residual weed control will be reduced.
- Observe all rotational intervals prior to planting as listed in the "ROTATIONAL RESTRICTIONS" table.

**FALL BURNDOWN AND FALLOW SEEDBED PROGRAMS**

*Valor* EZ Herbicide 3 [at 2 to 4 fl oz/A (0.063 to 0.125 lb ai)] can be used in the fall to provide residual weed control in fields that will be planted the following spring with field corn, peanut or soybean (refer to Rotational Restrictions table for rates and rotational intervals prior to planting). Weeds controlled by residual activity are listed in Table 1 *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3* (sections A and B), Broadleaf Weeds Controlled by Residual Activity of *Valor* EZ Herbicide 3; Table 3, *Weeds Controlled by Fall and Spring Preplant Burndown Programs*; and Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*. If weeds have emerged at the time of application, use *Valor* EZ Herbicide 3 in combination with a labeled burndown herbicide.

*Valor* EZ Herbicide 3 can be used in a fall burndown or fallow seedbed program, however the length of residual control may be variable.

Abnormally warm or wet winters will reduce the length of weed control observed in the spring.

Weeds controlled by postemergence or residual activity are listed in Table 3, *Weeds Controlled by Fall and Spring Preplant Burndown Programs*. Preplant burndown treatment tank mixes and rates are:

| Herbicide                                                          | Product Rate                                                        |
|--------------------------------------------------------------------|---------------------------------------------------------------------|
| <b>Program 1<sup>(1)</sup></b>                                     |                                                                     |
| Valor EZ Herbicide 3<br><b>Plus</b>                                | 2 to 3 fl oz/A (0.063 to 0.094 lb ai)                               |
| glyphosate<br><b>Plus</b>                                          | 0.5 to 1.0 lb ai/A (equivalent to 1 to 2 pt/A of Roundup Original®) |
| 2,4-D LVE (2,4-D for use on preplant soybeans only)<br><b>Plus</b> | 0.5 to 1.0 lb ai/A (equivalent to 1 to 2 pt/A of 2,4-D 4 LVE)       |
| NIS + AMS                                                          | 0.5% v/v + 17 lb/100 gallons of water                               |

or

|                                       |                                                                    |
|---------------------------------------|--------------------------------------------------------------------|
| <b>Program 2<sup>(1)</sup></b>        |                                                                    |
| Valor EZ Herbicide 3<br><b>Plus</b>   | 2 to 3 fl oz/A (0.063 to 0.094 lb ai)                              |
| glyphosate<br><b>Plus</b>             | 0.5 to 1.0 lb ai/A (equivalent to 1 to 2 pt/A of Roundup Original) |
| COC <sup>(2)</sup><br>or<br>NIS + AMS | 1pt/A<br>or<br>0.5% v/v + 17 lb/100 gallons of water               |

or

|                                                                    |                                                               |
|--------------------------------------------------------------------|---------------------------------------------------------------|
| <b>Program 3<sup>(1)</sup></b>                                     |                                                               |
| Valor EZ Herbicide 3<br><b>Plus</b>                                | 2 to 3 fl oz/A (0.063 to 0.094 lb ai)                         |
| 2,4-D LVE (2,4-D for use on preplant soybeans only)<br><b>Plus</b> | 0.5 to 1.0 lb ai/A (equivalent to 1 to 2 pt/A of 2,4-D 4 LVE) |
| COC                                                                | 1 pt/A                                                        |

<sup>(1)</sup>Dicamba (Banvel®), at 0.188 lb ai/A (6 fl oz/A of Banvel 4) can be added to Programs 1, 2 & 3 to assist in the control emerged broadleaves. Refer to dicamba label for rotational restrictions.

<sup>(2)</sup>Crop oil concentrate has been found to increase glyphosate burndown of emerged cutleaf evening primrose and Carolina geranium.

**Table 3. Weeds Controlled by Fall and Spring Preplant Burndown Programs**

| Weeds Controlled <sup>(1)</sup>          |                                | Postemergence           |                    |                    | Residual |
|------------------------------------------|--------------------------------|-------------------------|--------------------|--------------------|----------|
| Common Name                              | Scientific Name                | Program 1               | Program 2          | Program 3          |          |
|                                          |                                | Weeds 3 inches or less  |                    |                    |          |
| Chamomile, False                         | <i>Matricaria maritime</i>     | Yes                     | Yes                | No                 | Yes      |
| Cheatgrass                               | <i>Bromus tectorum</i>         | Yes                     | Yes                | No                 | Yes      |
| Chickweed, Common                        | <i>Stellaria media</i>         | Yes                     | Yes                | No                 | Yes      |
| Chickweed, Mouseear                      | <i>Cerastium vulgatum</i>      | Yes                     | Yes                | No                 | Yes      |
| Cockle, White                            | <i>Silene latifolia</i>        | No                      | Yes                | Yes                | Yes      |
| Dandelion                                | <i>Taraxacum officinale</i>    | Yes                     | No                 | Yes <sup>(2)</sup> | Yes      |
| Deadnettle, Purple                       | <i>Lamium purpureum</i>        | Yes                     | Yes                | Yes                | Yes      |
| Groundsel, Cressleaf                     | <i>Senecio glabellus</i>       | Yes                     | Yes                | -                  | Yes      |
| Henbit                                   | <i>Lamium amplexicaule</i>     | Yes                     | Yes                | Yes                | Yes      |
| Kochia                                   | <i>Kochia scoparia</i>         | Yes                     | Yes                | Yes                | Yes      |
| Marestail/Horseweed                      | <i>Conyza canadensis</i>       | Yes                     | Yes <sup>(3)</sup> | Yes                | Yes      |
| Mallow, Common                           | <i>Malva neglecta</i>          | Yes                     | Yes                | No                 | Yes      |
| Prickly Lettuce                          | <i>Lactuca serriola</i>        | Yes                     | Yes                | Yes                | Yes      |
| Wormwood, Biennial                       | <i>Artemisia biennis</i>       | Yes                     | Yes                | Yes                | Yes      |
|                                          |                                | Weeds 12 inches or less |                    |                    |          |
| Canola, Volunteer                        | <i>Brassica napus</i>          | Yes                     | Yes                | Yes                | Yes      |
| Carolina Geranium                        | <i>Geranium carolinianum</i>   | Yes                     | Yes                | Yes                | -        |
| Evening primrose, Cutleaf <sup>(4)</sup> | <i>Oenothera laciniata</i>     | Yes                     | Yes                | Yes                | Yes      |
| Flixweed                                 | <i>Descurainia sophia</i>      | Yes                     | Yes                | Yes                | Yes      |
| Mustard, Tansy                           | <i>Descurainia pinnata</i>     | Yes                     | Yes                | Yes                | Yes      |
| Mustard, Wild                            | <i>Brassica kaber</i>          | Yes                     | Yes                | Yes                | Yes      |
| Shepherd's-purse                         | <i>Capsella bursa-pastoris</i> | Yes                     | Yes                | Yes                | Yes      |

<sup>(1)</sup>Refer to glyphosate and/or 2,4-D labels for additional weeds controlled and rotational restrictions.

<sup>(2)</sup>Use 1 lb ai/A of 2,4-D LVE (equivalent to 2 pt/A of 2,4-D 4 LVE) for control of emerged dandelion.

<sup>(3)</sup>Program 2 will not control emerged glyphosate resistant marestail/horseweed.

<sup>(4)</sup>Use Program 1 to control cutleaf evening primrose that are nearing 12 inches in height or are past the rosette stage.

Use Programs 2 or 3 to control cutleaf evening primrose that are 12 inches or less and in the rosette stage.

### SPRING BURNDOWN PROGRAMS

*Valor* EZ Herbicide 3 can be used in combination with labeled preplant burndown herbicides to assist in the postemergence burndown of emerged weeds and provide residual weed control prior to crop emergence. Weeds controlled by residual activity are listed in Table 1 *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row. Apply *Valor* EZ Herbicide 3 after planting peanuts and soybeans when these types of planters are used (within 3 days after planting soybeans, within 2 days after planting peanuts and before the crop emerges). *Valor* cannot be applied after planting field corn.

*Valor* EZ Herbicide 3 can be used [at 1 to 3 fl oz/A (0.032 to 0.094 lb ai)] with labeled preplant burndown herbicides to enhance the speed of burndown and increase weed spectrum.

*Valor* EZ Herbicide 3 can be used [at 1 to 3 fl oz/A (0.032 to 0.094 lb ai)] [1 to 2 fl oz/A (0.032 to 0.063 lb ai)] in field corn, peanut and soybean burndown programs. See "DIRECTIONS FOR USE IN FIELD CORN", "DIRECTIONS FOR USE IN PEANUT", "DIRECTIONS FOR USE IN SOYBEAN" for more information.

**DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN  
PROGRAMS IN COTTON AND SUGARCANE**  
[For Use in Arizona, California and Hawaii Only]

**RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply to frozen or snow covered soil.
- **DO NOT** perform any tillage operation after application or residual weed control will be reduced.
- *Valor* EZ Herbicide 3 can be used [at 1 to 2 fl oz/A (0.032 lb to 0.063 lb flumioxazin)] with labeled burndown herbicides to enhance the speed of burndown and increase weed spectrum.
- A minimum of 30 days must pass, and 1 inch of rainfall/irrigation must occur, between *Valor* EZ Herbicide 3 application and planting of conventionally tilled cotton.
- A minimum of 14 days must pass, and 1 inch of rainfall/irrigation must occur, between *Valor* EZ Herbicide 3 application and planting of no-till or strip-till cotton when a *Valor* EZ Herbicide 3 rate of 1 fl oz/A (0.315 lb flumioxazin) is used and 21 days when a *Valor* EZ Herbicide 3 rate of 1.5 to 2 fl oz/A (0.0315 lb to 0.063 lb flumioxazin) is used. The field must contain the stubble from the previous crop.
- *Valor* EZ Herbicide 3 can be applied as part of a burndown application to sugarcane until cane emergence.
- Observe all rotational intervals prior to planting as listed in the “ROTATIONAL RESTRICTIONS” table.
- Refer to most restrictive label for minimum interval between application and planting.

**FALL BURNDOWN PROGRAMS**

*Valor* EZ Herbicide 3 [at 2 to 4 fl oz/A (0.063 to 0.125 lb ai)] can be used in the fall to provide residual weed control in fields that will be planted the following spring with cotton or sugarcane (refer to Rotational Restrictions table for rates and rotational intervals prior to planting). Weeds controlled by residual activity are listed in Table 1 *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3* and Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*. If weeds have emerged at the time of application, use *Valor* EZ Herbicide 3 in combination with a labeled burndown herbicide.

Abnormally warm or wet winters will reduce the length of weed control observed in the spring.

**SPRING BURNDOWN PROGRAMS**

*Valor* EZ Herbicide 3 [at 1 to 2 fl oz/A (0.032 to 0.063 lb ai)] can be used in combination with labeled preplant burndown herbicides to assist in the postemergence burndown of emerged weeds and provide residual weed control prior to crop emergence in fields that will be planted with cotton or sugarcane. Weeds controlled by residual activity are listed in Table 1 *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row.

**DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN PROGRAMS IN RICE,  
SORGHUM, SUNFLOWER, TOBACCO AND WHEAT  
(Preplant to Crop)**

[For Use in Arizona, California and Hawaii Only]

**RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply to frozen or snow covered soil.
- **DO NOT** perform any tillage operation after application or residual weed control will be reduced.
- *Valor* EZ Herbicide 3 can be used [at 1 to 2 fl oz/A (0.032 lb to 0.063 lb flumioxazin)] with labeled burndown herbicides to enhance the speed of burndown and increase weed spectrum. A minimum of 30 days must pass, and 1 inch of rainfall/irrigation must occur, between *Valor* EZ Herbicide 3 application and planting of rice, sorghum, sugarcane, sunflowers, tobacco or wheat. Refer to most restrictive label for minimum interval between application and planting.
- Observe all rotational intervals prior to planting as listed in the "ROTATIONAL RESTRICTIONS" table.

**FALL BURNDOWN PROGRAMS**

*Valor* EZ Herbicide 3 can be used in combination with labeled burndown programs to control emerged weeds and provide residual weed control in fields that will be planted the following spring (refer to Rotational Restrictions table for rates and rotational intervals prior to planting).

Abnormally warm winters may reduce the length of weed control observed in the spring.

**SPRING BURNDOWN PROGRAMS**

*Valor* EZ Herbicide 3 [at 1 to 2 fl oz/A (0.032 to 0.063 lb ai),] can be used in combination with labeled burndown programs to control emerged weeds and provide residual weed control prior to crop emergence. Weeds controlled by residual activity are listed in Table 1 *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3* Section A. Crops that will be planted following application must be in compliance with the rotational interval listed in the "Rotational Restriction" table above.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row.

**DIRECTIONS FOR USE IN FALL BURNDOWN PROGRAMS IN FIELDS TO BE PLANTED TO  
BARLEY, FIELD PEAS, FLAX, LENTIL, SAFFLOWER, SUNFLOWER AND SPRING WHEAT  
(Preplant to Crop)**

[For Use in Arizona, California and Hawaii Only]

**RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply to frozen or snow-covered soil.
- **DO NOT** perform any tillage operation after application or residual weed control will be reduced.
- *Valor* EZ Herbicide 3 can be mixed with 2,4-D and/or glyphosate formulations labeled for burndown programs (preplant to crop) in accordance with the most restrictive label limitations and precautions.
- **DO NOT** mix *Valor* EZ Herbicide 3 with any product containing a label prohibition against such mixing.
- Observe all rotational intervals prior to planting as listed in the "ROTATIONAL RESTRICTIONS" table.

**FALL BURNDOWN PROGRAMS**

*Valor* EZ Herbicide 3 can be used [at 2 to 4 fl oz/A (0.063 to 0.125 lb ai)] with labeled burndown herbicides to enhance the speed of burndown, increase weed spectrum and provide residual weed control of the weeds listed in Table 3, *Weeds Controlled by Fall and Spring Preplant Burndown Programs* until the following spring. Rotational intervals must be followed for crop to be planted in the spring following the fall *Valor* EZ Herbicide 3 application. Refer to most restrictive label for minimum interval between application and planting.

## **DIRECTIONS FOR USE IN FALLOW LAND**

[For Use in Arizona, California and Hawaii Only]

*Valor* EZ Herbicide 3 may be used as a preemergence fallow treatment. Weeds controlled by residual activity are listed in Table 1 *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*.

*Valor* EZ Herbicide 3 [at 2 to 4 fl oz/A (0.063 to 0.125 lb ai)] can be used in the fall to provide residual weed control in fallow fields (refer to Rotational Restrictions table for rates and rotational intervals prior to planting). If weeds have emerged at the time of application, use *Valor* EZ Herbicide 3 in combination with a labeled fallow herbicide. Abnormally warm or wet winters will reduce the length of weed control observed in the spring.

*Valor* EZ Herbicide 3 [at 1 to 4 fl oz/A (0.032 to 0.125 lb ai),] can be used in spring in combination with labeled burndown herbicides to control emerged weeds and provide residual weed control.

## **DIRECTIONS FOR USE IN ESTABLISHED ALFALFA**

### **RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply more than 4 fl oz (0.125 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** apply more than 2 applications of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 8 fl oz (0.250 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** make a sequential *Valor* EZ Herbicide 3 application within 60 days of the first *Valor* EZ Herbicide 3 application.
- **DO NOT** apply to alfalfa with greater than 6 inches of growth. Application will result in burning of treated leaves and stems. **Understand and accept this risk before using *Valor* EZ Herbicide 3 on alfalfa.**
- **DO NOT** apply within 25 days of harvest or grazing.
- **DO NOT** use on alfalfa grown for seed unless approved by a State authority to support a Special Local Need (SLN) under FIFRA section 24(c).
- Only apply with an adjuvant or tank mix with products formulated as an emulsifiable concentrate "EC" when targeting control of emerged weeds (expect and accept crop burn and/or stunting if *Valor* EZ Herbicide 3 is used with an adjuvant, a tank mix partner formulated as an emulsifiable concentrate (EC) or a tank mix partner formulated with an adjuvant.)
- Application with paraquat can be used to burndown winter annuals prior to winter dormant period.
- **DO NOT** use on intended mixed alfalfa-grass stands.

### **TIMING TO ALFALFA**

*Valor* EZ Herbicide 3 may be applied to established alfalfa with a maximum amount of growth of 6 inches or less for the preemergence control of the weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*. Established alfalfa is defined as alfalfa planted in the fall or spring which has gone through a first cutting/mowing. Application to alfalfa with greater than 6 inches of growth may result in unacceptable crop injury.

For control of winter annual weeds: the best timing for preemergence control is in the fall immediately after the last cutting or sheepling-off has occurred.

For control of summer annual weeds: the best timing for preemergence control is in the spring prior to alfalfa growth and before 6 inches of growth.

## **TIMING TO WEEDS**

### **Preemergence – Preemergence To Weeds**

Apply *Valor* EZ Herbicide 3 before alfalfa growth exceeds 6 inches in height for the preemergence control of weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*. Make applications as soon as possible after cutting and removing alfalfa to minimize injury to alfalfa growth.

### **Postemergence Dodder Suppression**

Apply *Valor* EZ Herbicide 3 at 4 fl oz (0.125 lb ai) per acre with an adjuvant for postemergence suppression of dodder. Tank mixes with Pursuit® Herbicide or Raptor® Herbicide will increase control.

## **DIRECTIONS FOR USE IN ARTICHOKE**

### **RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply more than 4 fl oz (0.125 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application on annual or perennial artichoke varieties after new planting.
- **DO NOT** apply more than 6 fl oz (0.188 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application on perennial artichoke varieties after cutback.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 6 fl oz (0.188 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- Application to artichoke foliage may result in unacceptable crop injury.

### **TIMING TO ARTICHOKE**

**Annual Varieties:** *Valor* EZ Herbicide 3 may be applied to artichoke beds prior to transplanting.

Application of *Valor* EZ Herbicide 3 must be made to the beds no later than 2 days prior to transplanting. Irrigation or rainfall after transplanting is necessary to activate the *Valor* EZ Herbicide 3.

**DO NOT** irrigate the *Valor* EZ Herbicide 3 before transplanting. Heavy irrigation or rainfall may result in crop injury. The injury is usually transitory and the plants will quickly grow out of the crop damage. Take care to minimize soil disturbance during transplanting, as preemergence weed control will decrease as soil disturbance increases.

**Perennial Varieties:** *Valor* EZ Herbicide 3 may be applied to artichokes after planting of crown pieces or “cut back” of mature plants. Applications of *Valor* EZ Herbicide 3 must be made within 2 days after planting or cut back and prior to artichoke emergence. Application after the artichokes have begun to crack, or are emerged, will result in crop injury. Apply before artichokes have begun to emerge (cracking).

### **TIMING TO WEEDS**

#### **Pre-plant (annual)/Preemergence (perennial) to Artichokes - Preemergence to Weeds**

Apply *Valor* EZ Herbicide 3 pre-plant to annual artichokes for preemergence control of the weeds. For perennial artichokes apply before cracking for preemergence control the weeds. Apply prior to weed emergence. A post-emergence herbicide may be necessary to control emerged weeds. *Valor* EZ Herbicide 3 may be applied to annual or perennial artichokes as specified above for preemergence control of weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*.

## DIRECTIONS FOR USE IN ESTABLISHED ASPARAGUS

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 6 fl oz (0.188 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 6 fl oz (0.188 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- Apply only to dormant asparagus no less than 14 days before spears emerge. Application to non-dormant asparagus may result in unacceptable crop injury.
- **[DO NOT** work soil within 60 days prior to application in the spring. Soil can be worked after spear harvest in preparation for *Valor* Herbicide application prior to fern emergence. Treated soil that is splashed onto the ferns may result in spotting.]

### TIMING TO ASPARAGUS - Dormant

*Valor* EZ Herbicide 3 may be applied to dormant asparagus for preemergence control of the weeds listed in Table 10, *Weeds Controlled by Preemergence Application of Valor EZ Herbicide 3*. Application to non-dormant asparagus will result in unacceptable crop injury. Apply a minimum of two weeks before spear emergence. Scoring may result if a minimum of 0.5 inch of either rainfall or irrigation has not occurred two weeks prior to emergence.

### TIMING TO ASPARAGUS – Post Harvest

Apply *Valor* EZ Herbicide 3 after the final harvest of the year, but prior to fern emergence, for preemergence control of the weeds listed in Table 10, *Weeds Controlled by Preemergence Application of Valor EZ Herbicide 3*. Application after fern emergence will result in unacceptable crop injury. Apply no less than two weeks prior to fern emergence and must be sprinkler or rainfall incorporated with 1/2 to 3/4 inches of water. Add a burndown tank mix partner for the control of emerged weeds labeled for asparagus in accordance with the most restrictive labeled limitations and precautions.

### TIMING TO WEEDS

#### Burndown – Dormant Asparagus, Postemergence to Weeds

*Valor* EZ Herbicide 3 may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds where asparagus is dormant. For control of emerged weeds, tank mix *Valor* EZ Herbicide 3 with paraquat. Refer to paraquat label for rates and application parameters. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. *Valor* EZ Herbicide 3 tank mixes applied to assist in the control of emerged weeds must be applied with a non-ionic surfactant at 0.25% v/v. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lb/A or 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to increase herbicidal activity.

#### Burndown – After Last Harvest of Season, Postemergence to Weeds

Use *Valor* EZ Herbicide 3 for residual weed control and to assist in postemergence burndown for many annual and perennial weeds where asparagus harvest has been completed for the year. For control of emerged weeds, use a labeled tank mix partner with activity on the emerged weeds.

#### Preemergence – Dormant Asparagus or After Last Harvest of the Year, Preemergence to Weeds

Apply *Valor* EZ Herbicide 3 for the preemergence control of weeds listed in Table 10, *Weeds Controlled by Preemergence Application of Valor EZ Herbicide 3*.

## DIRECTIONS FOR USE ON BRASSICA HEAD AND STEM VEGETABLES CROP GROUP 5-16

**Includes:** Broccoli; Brussels Sprouts; Cabbage; Cabbage, Chinese, napa; Cauliflower; cultivars, varieties, and/or hybrids of these.

**[FOR DISTRIBUTION AND USE ONLY WHERE THIRD PARTY INDEMNIFICATION IS IN EFFECT]**

### ROW MIDDLES

#### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply after crops are transplanted.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application. For Cabbage **DO NOT** apply more than 4 oz (0.125 lb flumioxazin) of *Valor* Herbicide per acre per application.
- **DO NOT** make more than 2 applications of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 6 fl oz (0.188 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year. For Cabbage **DO NOT** apply more than 8 fl oz (0.250 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.

#### PRECAUTIONS

- *Valor* EZ Herbicide 3 can only be applied in row middles between raised plastic mulched beds that are at least **4 inches higher** than the treated row middle and the mulched bed must have **a minimum of a 24-inch bed width**.
- Spray must remain between raised beds and contact no more than the bottom 1 inch of the side of the raised bed.
- All applications must be made with shielded or hooded equipment.
- Injury can occur if soil particles treated with *Valor* EZ Herbicide 3 contact the crop.
- A rainfall after application but prior to transplanting is required.

#### RATE

Up to 4 fl oz (0.125 lb ai) per acre per application.

#### TIMING TO CROP

*Valor* EZ Herbicide 3 may be applied at 3 fl oz (0.094 lb ai) per acre (except cabbage may be applied at 4 oz/A (0.125 lb ai)) as a shielded or hooded application to row middles after plastic is laid up to transplanting. Transplanting or seeding can take place any time after spray has dried. Spray must be directed to the row middle and contact no more than approximately the bottom 1 inch of the side of the raised bed. If the top of the mulch beds (where plants are to be transplanted) is contacted, severe injury can occur due to foliage contact with treated plastic.

#### WEED CONTROL AND TANK MIXING

*Valor* EZ Herbicide 3 provides preemergence residual control of the weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*, as well as to assist in the postemergence control of emerged weeds. A registered preemergence grass herbicide may be added for control of additional grassy weeds. For control of emerged weeds, tank mix *Valor* EZ Herbicide 3 with paraquat, Aim™, glyphosate, or other registered burndown herbicide. Refer to tank mix partner label for rates and application parameters.

## DIRECTIONS FOR USE ON CACTUS (PRICKLY PEAR)

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 12 fl oz (0.375 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 2 applications of *Valor* EZ Herbicide 3 per acre per year at the 6 fl oz (0.188 lb flumioxazin) rate.
- **DO NOT** apply more than 12 fl oz (0.375 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- Use a maximum *Valor* EZ Herbicide 3 rate of 6 fl oz/A (0.188 lb flumioxazin) per application on any soil that has a sand plus gravel content over 80% if plants are less than 3 years of age. (Two applications of 6 fl oz/A (0.188 lb flumioxazin) in a 12-month period can still be made as long as there have been 60 days between applications).
- **DO NOT** apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation.
- Raise mower height during all mowing to reduce dust. Dust created by mowing can drift onto desirable vegetation resulting in injury.
- **DO NOT** mow treated areas. Dust created by mowing may drift onto desirable vegetation resulting in injury.
- Follow the most restrictive label limitations and precautions of the tank mix product(s) being used.
- Avoid direct or indirect spray contact to foliage.
- **DO NOT** apply within 60 days prior to harvest.
- **DO NOT** apply to plants established less than one year.

Apply *Valor* EZ Herbicide 3 as a uniform broadcast application to the plantation floor or as a uniform band directed at the base of the cactus. The preferred application timing for *Valor* EZ Herbicide 3 is in the fall to maximize the potential for rainfall to activate and set the herbicide. **DO NOT** apply over the top of crop or allow spray to come in contact with crop as a result of application or drift.

### Preemergence Application

Apply 6 to 12 fl oz (0.188 to 0.375 lb ai) of *Valor* EZ Herbicide 3 per broadcast acre as a preemergence application. *Valor* EZ Herbicide 3 applications must be made prior to weed emergence for control of weeds listed in Table 10, *Weeds Controlled by Preemergence Application of Valor EZ Herbicide 3*. Make preemergence (to weed emergence) applications of *Valor* EZ Herbicide 3 to a weed-free soil surface. Preemergence applications of *Valor* EZ Herbicide 3 must be completed prior to weed emergence. Moisture is necessary to activate *Valor* EZ Herbicide 3 on soil for residual weed control. Dry weather following application of *Valor* EZ Herbicide 3 may reduce effectiveness. However, when adequate moisture is received after dry conditions, *Valor* EZ Herbicide 3 will control susceptible germinating weeds.

### Postemergence Application

Apply 6 to 12 fl oz (0.188 to 0.375 lb ai) of *Valor* EZ Herbicide 3 per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 qt/A crop oil concentrate). The addition of an adjuvant enhances *Valor* EZ Herbicide 3 activity on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of *Valor* EZ Herbicide 3.

Refer to Table 13, *Weeds Controlled by Postemergence Activity of Valor EZ Herbicide 3 Tank Mixes* for weeds controlled by the residual activity of *Valor* EZ Herbicide 3. Tank mix *Valor* EZ Herbicide 3 with a labeled burndown herbicide for control of the emerged weeds.

Residual weed control will be reduced if vegetation prevents the *Valor* EZ Herbicide 3 from reaching the soil surface. If vegetation is heavy, use a burndown herbicide with *Valor* EZ Herbicide 3 and make a sequential *Valor* EZ Herbicide 3 application prior to the emergence of new weeds.]

### Carrier Volume and Spray Pressure

To ensure thorough coverage in burndown applications, use a minimum of 15 gallons of spray solution per acre. Use higher gallonage if dense vegetation or heavy crop residue is present.

Nozzle selection must meet manufacturer's gallonage and pressure guidelines.

**Banded Application**

Rates listed in Table 13, *Weeds Controlled by Postemergence Activity of Valor EZ Herbicide 3 Tank Mixes*, refer to a broadcast application covering the entire acre. Refer to the Band Application table in Use Information Section to calculate amount needed per acre when making a banded application.

**DIRECTIONS FOR USE IN CELERY**

[For Use in] [California,] [Michigan and Wisconsin Only]

**RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre during a pre-transplant application.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre during a post-transplant application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** use with an adjuvant.
- Post-transplant applications must be made between 3 to 7 days following transplanting.
- **DO NOT** apply as part of a tank mix.

**[PRECAUTIONS]**

- [In the state of California, use as pre-transplant application only.]

**TIMING TO CELERY**

Apply *Valor* EZ Herbicide 3 at 3 fl oz/A (0.094 lb ai) prior to transplanting, or between 3 and 7 days following transplanting, for preemergence control of the weeds listed in Table 1, Broadleaf Weeds Controlled by Residual Activity of *Valor* EZ Herbicide 3.

**TIMING TO WEEDS**

Use *Valor* EZ Herbicide 3 prior to weed emergence for residual control.

**Refer to Product Information section for tank mix guidance. *Valor* EZ Herbicide 3, when applied according to label use directions, will control the weeds listed in Table 1, *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*.**

## DIRECTIONS FOR USE IN ESTABLISHED CLOVER AND CLOVER GROWN FOR SEED

For Use in Idaho, Oregon and Washington Only

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 4 fl oz (0.125 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 4 fl oz (0.125 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply within 25 days of harvest or grazing.
- Application to clover with greater than 6 inches of growth may result in unacceptable crop injury.

### PRECAUTIONS

- Applications to clover with 6 inches of growth will result in burning of treated leaves and stems. **Understand and accept this risk before using *Valor* EZ Herbicide 3 on clover.**
- Only apply with an adjuvant or tank mix with products formulated as an emulsifiable concentrate "EC" when targeting control of emerged weeds (expect and accept crop may be burned and/or stunting when applying tank mixes of *Valor* EZ Herbicide 3 with an adjuvant).
- Application with paraquat can be used to burndown winter annuals prior to winter dormant period.
- Application to mixed clover grass stands may result in unacceptable injury to the grass.

### TIMING TO CLOVER

*Valor* EZ Herbicide 3 may be applied to established clover with a maximum amount of growth of 6 inches or less for the preemergence control of the weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*. Established Clover is defined as clover planted in the fall or spring which has gone through a first cutting/mowing.

For control of winter annual weeds: the best timing for preemergence control is in the fall immediately after the last cutting or sheepling-off has occurred.

For control of summer annual weeds: the best timing for preemergence control is in the spring prior to clover growth and before 6 inches of growth.

### TIMING TO WEEDS

#### Preemergence – Preemergence to Weeds

Apply *Valor* EZ Herbicide 3 before clover growth exceeds 6 inches in height for the preemergence control of weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*. Make applications as soon as possible after cutting and removing clover to minimize injury to clover growth.

#### Postemergence Dodder Suppression

Apply *Valor* EZ Herbicide 3 at 4 fl oz per acre (0.125 lb ai) with an adjuvant for postemergence suppression of dodder. Tank mixes with Pursuit Herbicide or Raptor Herbicide will increase control.

**DIRECTIONS FOR USE IN COTTON**  
[For Use in Arizona, California and Hawaii Only]

**RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply more than 2 fl oz (0.063 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 2 applications of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 4 fl oz (0.125 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** make a sequential *Valor* EZ Herbicide 3 application within 30 days of the first *Valor* EZ Herbicide 3 application.
- **DO NOT** apply within 60 days of harvest.

**ENVIRONMENTAL CONDITIONS AND BIOLOGICAL PERFORMANCE**

**Hooded, Shielded and Layby Application**

For best results, apply *Valor* EZ Herbicide 3 to actively growing weeds within the growth stages indicated in this label. Applying *Valor* EZ Herbicide 3 under conditions that do not promote active weed growth will reduce herbicide effectiveness. **DO NOT** apply *Valor* EZ Herbicide 3 when the crop or weeds are under stress due to drought, excessive water, extremes in temperature, disease or low humidity. Weeds under stress tend to become less susceptible to herbicidal action. *Valor* EZ Herbicide 3 is most effective when applied under sunny conditions at temperatures above 65°F.

*Valor* EZ Herbicide 3 is rainfast one hour after application. **DO NOT** make applications if rain is expected within one hour of application or postemergence efficacy may be reduced.

**HERBICIDE RATE**

**Hooded, Shielded and Layby Application**

For postemergence weed control, apply *Valor* EZ Herbicide 3 through a hooded or shielded sprayer or at layby, at 2 fl oz/A (0.063 lb ai), in combinations with MSMA or at 1 to 2 fl oz/A (0.032 to 0.063 lb ai) in combination with glyphosate, to assist in the control of weeds listed in Table 4. *Emerged Broadleaf Weeds Controlled by Hooded, Shielded and Layby Application of Valor EZ Herbicide 3 Tank Mixes with Glyphosate or MSMA in Cotton*. Residual weed control can also be obtained through hooded, shielded and layby application of *Valor* EZ Herbicide 3. Weeds that are controlled through residual activity of *Valor* EZ Herbicide 3 are listed in Table 1 *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*. Weeds that are suppressed by residual activity of *Valor* EZ Herbicide 3 are listed in Table 2. *Weeds Suppressed by Residual Activity of Valor EZ Herbicide 3 Application Rates*.

**Table 4. Emerged Broadleaf Weeds Controlled by Hooded, Shielded and Layby Application of Valor EZ Herbicide 3 Tank Mixes with Glyphosate or MSMA in Cotton**

| Broadleaf Weed Species         |                                                    | Weed Height (inches)<br>2 fl oz/A<br>(0.063 lb ai) |
|--------------------------------|----------------------------------------------------|----------------------------------------------------|
| Common Name                    | Scientific Name                                    |                                                    |
| Bindweed, Field <sup>(1)</sup> | <i>Convolvulus arvensis</i>                        | 4                                                  |
| Carpetweed                     | <i>Mollugo verticillata</i>                        | 4                                                  |
| Chickweed, Common              | <i>Stellaria media</i>                             | 4                                                  |
| Cocklebur, Common              | <i>Xanthium strumarium</i>                         | 4                                                  |
| Florida Beggarweed             | <i>Desmodium tortuosum</i>                         | 2                                                  |
| Hemp Sesbania                  | <i>Sesbania exaltata</i>                           | 6                                                  |
| Jimsonweed                     | <i>Datura stramonium</i>                           | 4                                                  |
| Lambsquarters, Common          | <i>Chenopodium album</i>                           | 4                                                  |
| Morningglories                 |                                                    |                                                    |
| Entireleaf                     | <i>Ipomoea hederacea</i> var. <i>integriuscula</i> | 4                                                  |
| Ivyleaf                        | <i>Ipomoea hederacea</i>                           | 4                                                  |
| Pitted                         | <i>Ipomoea lacunose</i>                            | 4                                                  |
| Red                            | <i>Ipomoea coccinea</i>                            | 4                                                  |
| Tall                           | <i>Ipomoea purpurea</i>                            | 2                                                  |
| Mustard, Wild                  | <i>Brassica kaber</i>                              | 6                                                  |
| Nightshades                    |                                                    |                                                    |
| Black                          | <i>Solanum nigrum</i>                              | 4                                                  |
| Eastern Black                  | <i>Solanum ptycanthum</i>                          | 4                                                  |
| Hairy                          | <i>Solanum sarrachoides</i>                        | 4                                                  |
| Pigweeds                       |                                                    |                                                    |
| Palmer Amaranth                | <i>Amaranthus palmeri</i>                          | 4                                                  |
| Redroot                        | <i>Amaranthus retroflexus</i>                      | 4                                                  |
| Smooth                         | <i>Amaranthus hybridus</i>                         | 4                                                  |
| Plantain, Broadleaf            | <i>Plantago major</i>                              | 6                                                  |
| Prickly Sida (Teaweed)         | <i>Sida spinosa</i>                                | 4                                                  |
| Purslane, Common               | <i>Portulaca oleracea</i>                          | 2                                                  |
| Ragweeds                       |                                                    |                                                    |
| Common                         | <i>Ambrosia artemisiifolia</i>                     | 2                                                  |
| Giant                          | <i>Ambrosia trifida</i>                            | 4                                                  |
| Rice Flatsedge                 | <i>Cyperus iria</i>                                | 2                                                  |
| Sicklepod                      | <i>Senna obtusifolia</i>                           | 4                                                  |
| Smartweeds                     |                                                    |                                                    |
| Ladysthumb                     | <i>Polygonum persicaria</i>                        | 4                                                  |
| Pale                           | <i>Polygonum lapathifolium</i>                     | 4                                                  |
| Pennsylvania                   | <i>Polygonum pennsylvanicum</i>                    | 4                                                  |
| Spotted Spurge                 | <i>Euphorbia maculata</i>                          | 4                                                  |
| Velvetleaf                     | <i>Abutilon theophrasti</i>                        | 4                                                  |
| Venice Mallow                  | <i>Hibiscus trionum</i>                            | 2                                                  |
| Waterhemp                      |                                                    |                                                    |
| Common                         | <i>Amaranthus rudis</i>                            | 2                                                  |
| Tall                           | <i>Amaranthus tuberculatus</i>                     | 2                                                  |

<sup>(1)</sup> Valor EZ Herbicide 3 tank mixes will control the above ground portion of field bindweed. Repeated applications will be needed to control regrowth.

## CARRIER VOLUME AND SPRAY PRESSURE

### Hooded, Shielded and Layby Application

To ensure thorough coverage in hooded, shielded and layby applications, use 15 to 30 gallons spray solution per treated acre. Use 20 to 30 gallons per treated acre under heavy weed pressure. Nozzle selection must meet manufacturer's gallonage and pressure guidance for application method being used. **DO NOT** use "Flood Jet" nozzles, as they tend to increase the chance of crop injury.

## ADDITIVES

### Hooded, Shielded and Layby Application

Weed control from hooded, shielded or layby application of *Valor* EZ Herbicide 3 in cotton requires the addition of an agronomically approved non-ionic surfactant to the spray mixture. Non-ionic surfactant must contain at least 80% active ingredient. Verify mixing compatibility qualities by a jar test. **The use of crop oil concentrates, methylated seed oils, organo-silicant surfactants or products containing these ingredients, may result in severe crop injury.**

## APPLICATION EQUIPMENT

Apply *Valor* EZ Herbicide 3 tank mixes, with ground equipment using standard commercial sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume. Application equipment must be clean and in good repair. Nozzles must meet manufacturer's guidelines for spray pattern and placement on spray boom and must be checked frequently for accuracy.

## TIMING TO COTTON

### Hooded and Shielded Application

*Valor* EZ Herbicide 3 tank mixes may be applied with a hooded or shielded sprayer after cotton has reached a minimum of 6 inches in height. All nozzles must be under the hood or behind the shield to ensure no spray solution comes in contact with the cotton. **Care must be taken to ensure the spray solution or drift does not come in contact with the cotton or severe crop injury can occur.**

### Layby Application

Layby application of *Valor* EZ Herbicide 3 tank mixes may be made once cotton has reached a minimum of 16 inches in height. Cotton that is smaller than 16 inches in height may be injured by *Valor* EZ Herbicide 3 applications. *Valor* EZ Herbicide 3 application must be directed to the lower 2 inches of the cotton stem to avoid crop injury.

## TIMING TO WEEDS

*Valor* EZ Herbicide 3 tank mix applications must be made to weeds within the height range given in Table 4.

## TANK MIXES

*Valor* EZ Herbicide 3 must be tank mixed with one of the herbicides listed in Table 5. *Tank Mixes with Valor EZ Herbicide 3 for Hooded, Shielded and/or Layby Use in Cotton* for postemergence control of the weeds listed in Table 4. *Emerged Broadleaf Weeds Controlled by Hooded, Shielded and Layby Application of Valor EZ Herbicide 3 Tank Mixes with Glyphosate or MSMA in Cotton.* Refer to tank mix partner label for additional weeds controlled.

**Table 5. Tank Mixes with *Valor* EZ Herbicide 3 for Hooded, Shielded and/or Layby Use in Cotton**

| Tank Mix Partner | Hooded and Shielded | Layby            |
|------------------|---------------------|------------------|
| glyphosate       | X                   | X <sup>(1)</sup> |
| MSMA             | X                   | X                |
| Diuron           | X                   | X                |
| MSMA + Diuron    | X                   | X                |
| Glufosinate      | X                   | X <sup>(2)</sup> |
| Paraquat         | X                   | X                |

<sup>(1)</sup>For use only in cotton with the Roundup Ready gene.

<sup>(2)</sup>For use only in cotton with the LibertyLink® gene or glufosinate-resistant cotton.

## DIRECTIONS FOR USE IN CUCURBIT VEGETABLES

**Cucurbit Vegetables (Crop Group 9) including:** chayote (fruit); Chinese Waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp. (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes cantaloupe); pumpkin; squash, summer; squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon

### [FOR DISTRIBUTION AND USE ONLY WHERE THIRD PARTY INDEMNIFICATION IS IN EFFECT] ROW MIDDLES

Many weather-related factors, including high wind or heavy rains or cool conditions at or near crop transplanting, may result in crop injury in fields treated with *Valor* EZ Herbicide 3. On occasion this has resulted in a delay in maturity. Understand and accept these risks before using *Valor* EZ Herbicide 3.

Refer to Product Information section for tank mix guidance. *Valor* EZ Herbicide 3, when applied according to label use directions, will control the weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*.

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 4 fl oz (0.125 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 2 applications of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 8 fl oz (0.250 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** use with an adjuvant.
- **PRECAUTIONS**
- Grow plants on raised plastic mulched beds that are higher than the treated row middle.
- Spray must be directed to the row middle, away from the crop bed and with minimal contact with plastic, including the sides of the bed. If top of mulch beds (where plants are to be transplanted) is contacted, severe injury can occur due to foliage contact with treated plastic. In this scenario, a rainfall event of 1/2 inch (natural or irrigation) must occur prior to transplanting to reduce *Valor* EZ Herbicide 3 residues.
- Drift of treated soil particles onto plants may cause contact injury.
- Irrigate treated field after application and prior to transplanting with minimum of 1/4 inch of water if rainfall does not occur between application and transplanting.
- All applications must be made with hooded or shielded equipment.

### TIMING TO CUCURBIT VEGETABLES

Apply *Valor* EZ Herbicide 3 at 4 fl oz (0.125 lb ai) per acre as a hooded or shielded application to row middles up to 14 days prior to transplanting or seeding for preemergence control of the weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*, as well as to assist in the postemergence control of emerged weeds. A second application of *Valor* EZ Herbicide 3 at 4 fl oz (0.125 lb ai) per acre may be applied up to 21 days after transplanting or emergence if needed. **DO NOT** apply during or after bloom.

### TIMING TO WEEDS

*Valor* EZ Herbicide 3 may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds in row middles. A registered preemergence grass herbicide may be added for control of additional grassy weeds. For assisting in the control of emerged weeds, tank mix *Valor* EZ Herbicide 3 with paraquat, Aim™ or other registered burndown herbicide. **DO NOT** tank mix with glyphosate after transplanting. Refer to tank mix partner's label for rates and use directions.

## DIRECTIONS FOR USE IN DRY BEANS

Dried cultivars of bean (*Lupinus*); bean (*Phaseolus*) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean); bean (*Vigna*) (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean); broad bean (dry); chickpea (garbanzo bean); guar; lablab bean and lentil

### **WEED SUPPRESSION IN DRY BEANS AND WEED CONTROL IN CHICKPEAS (GARBANZO BEANS)**

**[Arizona, California, Colorado, Hawaii, Idaho, Montana, Nebraska, Oregon and Washington only]**

#### **RESTRICTIONS AND LIMITATIONS**

- For Chickpeas, **DO NOT** apply more than 2 fl oz (0.063 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application. For all other Dry Beans, **DO NOT** apply more than 1.5 fl oz (0.047 lb ai) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- For Chickpeas, **DO NOT** apply more than 2 fl oz (0.063 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year. For all other Dry Beans, **DO NOT** apply more than 1.5 fl oz (0.047 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.

**Many weather-related factors, including high wind, splashing or heavy rains or cool conditions at or near crop emergence, may result in dry bean injury in fields treated with *Valor* EZ Herbicide 3. On occasion this has resulted in a delay in maturity. Understand and accept these risks before using *Valor* EZ Herbicide 3.**

#### **TIMING TO DRY BEANS AND CHICKPEAS**

*Valor* EZ Herbicide 3 may be applied to dry beans within 2 days after planting for the preemergence suppression of the weeds listed in Table 1, *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3* or Table 8. *Weeds Suppressed by Residual Activity of Valor EZ Herbicide 3 at 1.5 oz/A*. Tank mix *Valor* EZ Herbicide 3 with other labeled herbicides for broad spectrum weed control.

#### **TIMING TO WEEDS**

*Valor* EZ Herbicide 3 may be applied to dry beans prior to planting or preemergence (after planting). Preemergence application of *Valor* EZ Herbicide 3 must be made within 2 days after planting and prior to dry bean emergence. To avoid severe crop injury, **DO NOT** apply to dry beans after beans begin to crack or have emerged.

Preplant incorporation (PPI) applications may result in reduced weed control.

#### **ADDITIONAL RESIDUAL GRASS CONTROL**

*Valor* EZ Herbicide 3 can be tank mixed with pendimethalin for additional grass control.

#### **HARVEST AID**

**[All states]**

#### **RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** harvest within 5 days of application.

Desiccation from *Valor* EZ Herbicide 3 requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 2% v/v. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lb/A or a 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to the spray mixture along with either a crop oil concentrate or methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for a crop oil concentrate or a methylated seed oil. Tank mixing *Valor* EZ Herbicide 3 with glyphosate or paraquat will increase control of emerged weeds and aid in harvest. Add a burndown tank mix partner for the control of emerged weeds labeled for dry bean in accordance with the most restrictive labeled limitations and precautions.

## TIMING TO DRY BEANS AND CHICKPEAS

Apply when crop is mature and at least 80% of the pods are yellowing and mostly ripe with no more than 40% (bush type beans) or 30% (vine type beans) of the leaves still green in color. Dry beans can be harvested 5 days after application. To ensure thorough coverage use 15 to 30 gallons spray solution per acre. Nozzle selection must meet manufacturer's gallonage and pressure guidelines for postemergence application.

## DIRECTIONS FOR USE IN FIELD CORN

[For Use in Arizona, California and Hawaii Only]

### RESTRICTIONS AND LIMITATIONS

- Use only on no-till or minimum tillage fields where last years crop residue has not been incorporated into the soil.
- Corn must be planted between 14 and 30 days after application unless the application is made as part of a Fall burndown program.
- Corn can be planted 7 days after an application of 2 fl oz/A (0.063 lb ai) if a minimum of 25% of the soil surface is covered with the residue of the preceding crop and a minimum of 1/4 inch of rainfall has occurred between application and planting.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** irrigate between emergence and 2-leaf corn.
- **DO NOT** use on popcorn, sweet corn or corn grown for seed.

### TIMING TO FIELD CORN

- Apply *Valor* EZ Herbicide 3, at 2 to 3 fl oz/A (0.063 to 0.094 lb ai), between 7 and 30 days prior to planting field corn for the preemergence control of the weeds listed in Table 1, *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*.
- Apply *Valor* EZ Herbicide 3 at 2 fl oz/A (0.063 lb ai) between 7 and 30 days prior to planting field corn if a minimum of 25% of the soil surface is covered with the residue of the preceding crop and a minimum of 1/4 inch of rainfall has occurred between application and planting.
- Apply *Valor* EZ Herbicide 3 at 3 fl oz/A (0.094 lb ai) between 14 and 30 days prior to planting field corn.

### Burndown Use Directions – For Preplant Applications in Field Corn

*Valor* EZ Herbicide 3, applied as part of a burndown program, may be used for residual weed control, as well as to assist in postemergence burndown of many weeds where field corn will be planted directly into the residue of the previous year. See Directions for Use in Fall and Spring Preplant Burndown and Fallow Seedbed Programs in Field Corn, Peanut and Soybean for rates and timing of applications. For control of emerged weeds, *Valor* EZ Herbicide 3 must be applied with an appropriate burndown tank mix partner listed in Table 6. *Tank Mix Partners for Burndown and/or Residual Control of Weeds in Field Corn*. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Refer to tank mix partner's label for application pressures and adjuvant systems.

#### INCREASING SPEED OF GLYPHOSATE BURNDOWN ACTIVITY

*Valor* EZ Herbicide 3, at 1 fl oz/A (0.032 lb ai), may be tank mixed with glyphosate (Roundup®) to increase the speed of burndown activity compared to glyphosate applied alone. Residual weed control will not be provided at rates lower than 2 fl oz/A (0.063 lb ai); however, suppression of the weeds in Table 2. *Weeds Suppressed by Residual Activity of Valor EZ Herbicide 3 Application Rates* may occur at *Valor* EZ Herbicide 3 rates as low as 1 fl oz/A (0.032 lb ai). Applications of *Valor* EZ Herbicide 3 at 1 fl oz/A (0.032 lb ai) must be made a minimum of 14 days prior to planting field corn.

### TANK MIXES

*Valor* EZ Herbicide 3 may be tank mixed with the herbicides listed in Table 6. *Tank Mix Partners for Burndown and/or Residual Control of Weeds in Field Corn* for pre-plant burndown applications.

Refer to tank mix partner's label for adjuvants.

**Table 6. Tank Mix Partners for Burndown and/or Residual Control of Weeds in Field Corn**

| <b>TANK MIX PARTNERS<sup>1</sup></b> |             |
|--------------------------------------|-------------|
| 2,4-D LVE                            | metribuzin  |
| atrazine                             | paraquat    |
| Basis®                               | Python®     |
| dicamba                              | Resolve®    |
| Express®                             | simazine    |
| glyphosate                           | Weedmaster® |
| Hornet®                              |             |

<sup>1</sup>Refer to tank mix product labels for specific application directions.

#### **TANK MIX RESTRICTIONS**

Tank mixes with flufenacet (Axiom or Domain), metolachlor or s-metolachlor (Dual Magnum or Dual II Magnum), dimethenamid or dimethenamid-p (Frontier or Outlook), or acetochlor (Surpass or Harness) may result in injury to field corn when application is followed by prolonged periods of cool wet weather.

## DIRECTIONS FOR USE IN FIELD PEAS

### WEED CONTROL

[For Use in Idaho, Montana, Oregon and Washington only.]

#### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 2 fl oz (0.063 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 2 fl oz of (0.063 lb flumioxazin) *Valor* EZ Herbicide 3 per acre per year.

**Many weather-related factors, including high wind, splashing or heavy rains or cool conditions at or near crop emergence, may result in peas injury in fields treated with *Valor* EZ Herbicide 3. On occasion this has resulted in a delay in maturity. Understand and accept these risks before using *Valor* EZ Herbicide 3.**

#### TIMING TO FIELD PEAS

*Valor* EZ Herbicide 3 may be applied to field peas within 2 days after planting for the preemergence control of the weeds listed in Table 1, *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3* or Table 8. *Weeds Suppressed by Residual Activity of Valor EZ Herbicide 3 at 1.5 oz/A*. Tank mix *Valor* EZ Herbicide 3 with other labeled herbicides for broad spectrum weed control.

#### TIMING TO WEEDS

*Valor* EZ Herbicide 3 may be applied to field peas prior to planting or preemergence (after planting). Preemergence application of *Valor* EZ Herbicide 3 must be made within 2 days after planting and prior to field pea emergence. To avoid severe crop injury, **DO NOT** apply to field peas after peas begin to crack or have emerged.

Preplant incorporation (PPI) applications may result in reduced weed control.

#### ADDITIONAL RESIDUAL GRASS CONTROL

*Valor* EZ Herbicide 3 can be tank mixed with pendimethalin for additional grass control.

### HARVEST AID

[All states]

#### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** harvest within 5 days of application.

Desiccation from *Valor* EZ Herbicide 3 requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lb/A or a 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* EZ Herbicide 3 with glyphosate will increase control of emerged weeds and aid in harvest.

#### TIMING TO FIELD PEAS

Apply *Valor* EZ Herbicide 3, at 1.5 to 2 fl oz/A (0.047 to 0.063 lb ai), when crop is physiologically mature and a minimum of 80% of the pods are yellow to tan in color and 20% are yellow in color. If field peas are treated too early, a reduction in seed quality may occur. **DO NOT** spray *Valor* Herbicide on any area of the field with a significant amount of plants with green color. Peas can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure guidelines for postemergence application.

## DIRECTIONS FOR USE IN FLAX

### **HARVEST AID**

#### **RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 2 applications of *Valor* EZ Herbicide 3 per acre per year at the 1.5 fl oz (0.047 lb ai) rate.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** harvest within 5 days of application.

Desiccation from *Valor* EZ Herbicide 3 requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lb/A or a 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil.

#### **TIMING TO FLAX**

Apply *Valor* EZ Herbicide 3, at 1.5 to 2 fl oz/A (0.047 to 0.094 lb ai), when crop is physiologically mature and at least 75% of the bolls are brown in color. Flax can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure guidelines for postemergence application.

## DIRECTIONS FOR USE IN FRUITING VEGETABLES

African eggplant; Bush Tomato; Bell Pepper; Cocona; Currant Tomato; Eggplant, Garden Huckleberry; Goji Berry; Groundcherry, Martynia; Naranjilla; Okra, Pea Eggplant; Pepino; Nonbell Pepper; Roselle; Scarlet Eggplant; Sunberry; Tomatillo; Tomato; Tree Tomato; cultivars, varieties and /or hybrids of these.

### [FOR DISTRIBUTION AND USE ONLY WHERE THIRD PARTY INDEMNIFICATION IS IN EFFECT] ROW MIDDLES

Many weather-related factors, including high wind or heavy rains or cool conditions at or near crop transplanting, may result in crop injury in fields treated with *Valor* EZ Herbicide 3. On occasion this has resulted in a delay in maturity. Understand and accept these risks before using *Valor* Herbicide.

## RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 4 fl oz (0.125 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 2 applications of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 8 fl oz (0.250 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.

## PRECAUTIONS

- Grow plants on raised or plastic mulched beds that are higher than the treated row middle.
- Spray must be directed to the row middle, away from the crop bed and with minimal contact with plastic, including the sides of the bed. If top of mulch beds (where plants are to be transplanted) is contacted, severe injury can occur due to foliage contact with treated plastic. In this scenario, a rainfall event of 1/2 inch (natural or irrigation) must occur prior to transplanting to reduce *Valor* EZ Herbicide 3 residues.
- Drift of treated soil particles onto plants may cause contact injury.
- Irrigate treated field after application and prior to transplanting with minimum of 1/4 inch of water if rainfall does not occur between application and transplanting.
- All applications must be made with hooded or shielded equipment.

## TIMING TO FRUITING VEGETABLES

Apply *Valor* EZ Herbicide 3 at 4 fl oz (0.125 lb ai) per acre as a hooded or shielded application to row middles up to 14 days prior to transplanting or seeding for preemergence control of the weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*, as well as to assist in the postemergence control of emerged weeds. A second application of *Valor* EZ Herbicide 3 at 4 fl oz (0.125 lb ai) per acre may be applied up to 21 days after transplanting or emergence if needed. **DO NOT** apply during or after bloom.

## TIMING TO WEEDS

*Valor* EZ Herbicide 3 may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds in row middles. A registered preemergence grass herbicide may be added for control of additional grassy weeds. For assisting in the control of emerged weeds, tank mix *Valor* EZ Herbicide 3 with paraquat, Aim™ or other registered burndown herbicide. **DO NOT** tank mix with glyphosate after transplanting or crop emergence. Refer to tank mix partner's label for rates and application parameters.

## DIRECTIONS FOR USE IN GARLIC

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 6 fl oz (0.188 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 6 fl oz (0.188 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.

### TIMING TO GARLIC

*Valor* EZ Herbicide 3 may be applied, at 6 fl oz/A (0.188 lb ai), to garlic prior to garlic emergence. Make application within 3 days after planting garlic.

### TIMING TO WEEDS

#### Preemergence – Preemergence To Weeds

Apply *Valor* EZ Herbicide 3 to weed free garlic for preemergence control of the weeds listed in Table 10, *Weeds Controlled by Preemergence Application of Valor EZ Herbicide 3*.

## DIRECTIONS FOR USE IN HOPS

[Not For Use in California and New York]

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 6 fl oz (0.188 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 6 fl oz (0.188 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** allow spray to contact green stem (unless used for sucker control), foliage, flowers or cones or unacceptable injury may occur.
- **DO NOT** apply within 30 days of harvest.
- **DO NOT** use with an adjuvant.

*Valor* EZ Herbicide 3 can be used in hops for preemergence weed control as well as sucker control.

### TIMING TO HOPS FOR SUCKER CONTROL

Apply *Valor* EZ Herbicide 3 at 6 fl oz/A (0.188 lb ai) as a directed application after hops have reached a minimum of 6 feet in height for sucker control. Direct application to the lower 2 feet of the hops.

### TIMING TO HOPS FOR PREEMERGENCE WEED CONTROL

Apply *Valor* EZ Herbicide 3 at 6 fl oz/A (0.188 lb ai) as a 1 to 1.5 foot band to each side of the hop row, to dormant hops November thru February to ensure time for rain incorporation and activation. If weeds are emerged at the time of application, tank mix *Valor* EZ Herbicide 3 with a labeled burndown herbicide including paraquat or glyphosate to assist with control of emerged weeds. **DO NOT** mow or rake over treated areas, as dust created by mowing may drift onto sensitive crops or vegetation resulting in injury.

### TIMING TO WEEDS

*Valor* EZ Herbicide 3 applications must be made prior to weed emergence for control of weeds listed in Table 10, *Weeds Controlled by Preemergence Application of Valor EZ Herbicide 3*.

**Refer to Product Information section for tank mix guidance. *Valor* EZ Herbicide 3, when applied according to label use directions, will control the weeds listed in Table 10, *Weeds Controlled by Preemergence Application of Valor EZ Herbicide 3*.**

## DIRECTIONS FOR USE IN LENTILS

### HARVEST AID

#### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** harvest within 5 days of application.

Desiccation from *Valor* EZ Herbicide 3 requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lb/A or a 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* EZ Herbicide 3 with glyphosate or paraquat will increase control of emerged weeds and aid in harvest.

#### TIMING TO LENTILS

Apply *Valor* EZ Herbicide 3, at 1.5 to 2 fl oz/A (0.047 to 0.063 lb ai), when crop is physiologically mature and a minimum of 80% of the pods are yellow to tan in color and 20% are yellow in color. If lentils are treated too early, a reduction in seed quality may occur. **DO NOT** spray *Valor* EZ Herbicide 3 on any area of the field with a significant amount of plants with green color. Lentils can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure guidelines for postemergence application.

## DIRECTIONS FOR USE IN MINT (Peppermint and Spearmint)

#### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 4 fl oz (0.125 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 2 applications of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 8 fl oz (0.250 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** make a sequential *Valor* EZ Herbicide 3 application within 60 days of the first *Valor* EZ Herbicide 3 application.
- Apply only to dormant mint. Application to non-dormant mint may result in unacceptable crop injury.
- **DO NOT** apply within 80 days of harvest.

#### PRECAUTIONS

##### To avoid crop injury:

- Application to stands established longer than 3 years may result in crop injury.
- Applications to stands with weak, thin, or damaged roots or rhizomes may result in crop injury.
- Application to mint in Southern Union County (south of Ladd Canyon) or Baker County in Oregon may result in unacceptable crop injury.
- Use only on established meadow mint.
- Applications to mint that has been weakened by diseases, insects (example mint root borer), nematodes, drought, soil salts, high soil pH, previous pesticides, winter injury or double cutting, may result in severe injury. Apply only to healthy vigorous mint with undamaged rhizomes.

**Many weather-related factors, including high wind, splashing or heavy rains or cool conditions at or near mint emergence, may result in mint injury in fields treated with *Valor* EZ Herbicide 3. Understand and accept these risks before using *Valor* EZ Herbicide 3.**

Tank mixes with labeled rates of paraquat are recommended to control emerged weeds and increase crop safety.

#### **TIMING TO MINT**

As a spray, *Valor* EZ Herbicide 3 may be applied only to established, dormant mint for preemergence control of the weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3* as well as to assist in the postemergence control of emerged weeds. Application to non-dormant mint or to baby (row) mint (time from planting of mint roots through the first cutting), may result in unacceptable crop injury. As a bulk fertilizer application, *Valor* EZ Herbicide 3 may be applied at least 80 days prior to harvest. Leaves must be dry at the time of applications or severe injury may occur.

#### **TIMING TO WEEDS**

##### **Burndown – Dormant Mint, Postemergence to Weeds**

*Valor* EZ Herbicide 3 may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds where established mint is dormant. For control of emerged weeds, tank mix *Valor* EZ Herbicide 3 with paraquat. Refer to the paraquat label for rates and use directions. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. *Valor* EZ Herbicide 3 tank mixes applied to assist in the control of emerged weeds must be applied with a non-ionic surfactant at 0.25% v/v. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lb/A or 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to increase herbicidal activity.

##### **Preemergence – Dormant Mint, Preemergence to Weeds**

Apply *Valor* EZ Herbicide 3 to dormant mint for the preemergence control of weeds listed in Table 7, *Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*. Fall applications of *Valor* EZ Herbicide 3, followed by a sequential application in the Spring, have resulted in better Summer annual weed control than a single Fall or single Spring application.

Fall application is most effective for Fall germinating weeds for example groundsel. Fields plowed or harrowed after a *Valor* EZ Herbicide 3 application will result in less effective preemergence activity. In furrow irrigated fields, corrugating that is done after a *Valor* EZ Herbicide 3 application will expose untreated soil and break the herbicide barrier resulting in poor weed control.

**Table 7. Weeds Controlled by Residual Activity of Valor EZ Herbicide 3**

| <b>Broadleaf Weed Species</b>               |                                  |                       |                  |                            |
|---------------------------------------------|----------------------------------|-----------------------|------------------|----------------------------|
| <b>Common Name</b>                          | <b>Scientific Name</b>           | <b>Organic Matter</b> | <b>Soil Type</b> | <b>Application Rate</b>    |
| Bristly Starbur                             | <i>Acanthospermum hispidum</i>   | Up to 5%              | All Soil Types   | 4 fl oz/A<br>(0.125 lb ai) |
| Carpetweed                                  | <i>Mollugo verticillata</i>      |                       |                  |                            |
| Chickweeds                                  |                                  |                       |                  |                            |
| Common                                      | <i>Stellaria media</i>           |                       |                  |                            |
| Mouseear                                    | <i>Cerastium vulgatum</i>        |                       |                  |                            |
| Coffee Senna                                | <i>Cassia occidentalis</i>       |                       |                  |                            |
| Copperleaf, Hophornbeam                     | <i>Acalypha ostryifolia</i>      |                       |                  |                            |
| Dandelion                                   | <i>Taraxacum officinale</i>      |                       |                  |                            |
| Dodder (suppression only)[*] <sup>(1)</sup> | <i>Cuscuta</i> spp.              |                       |                  |                            |
| Eclipta                                     | <i>Eclipta prostrate</i>         |                       |                  |                            |
| Evening primrose, Cutleaf                   | <i>Oenothera laciniata</i>       |                       |                  |                            |
| False Chamomile[*]                          | <i>Tripleurospermum maritima</i> |                       |                  |                            |
| Fiddleneck, Coast[*]                        | <i>Amsinckia menziesii</i>       |                       |                  |                            |
| Field Pennycress[*]                         | <i>Thlaspi arvense</i>           |                       |                  |                            |
| Fleabane, Hairy[*]                          | <i>Conyza bonariensis</i>        |                       |                  |                            |
| Flixweed[*]                                 | <i>Descurainia sophia</i>        |                       |                  |                            |
| Florida Beggarweed                          | <i>Desmodium tortuosum</i>       |                       |                  |                            |
| Florida Pusley                              | <i>Richardia scabra</i>          |                       |                  |                            |
| Golden Crownbeard                           | <i>Verbesina encelioides</i>     |                       |                  |                            |
| Groundsel, Common                           | <i>Senecio vulgaris</i>          |                       |                  |                            |
| Hairy Indigo                                | <i>Indigofera hirsuta</i>        |                       |                  |                            |
| Hemp Sesbania                               | <i>Sesbania exaltata</i>         |                       |                  |                            |
| Henbit                                      | <i>Lamium amplexicaule</i>       |                       |                  |                            |
| Jimsonweed                                  | <i>Datura stramonium</i>         |                       |                  |                            |
| Kochia                                      | <i>Kochia scoparia</i>           |                       |                  |                            |
| Lambsquarters, Common                       | <i>Chenopodium album</i>         |                       |                  |                            |
| Little Mallow                               | <i>Malva parviflora</i>          |                       |                  |                            |
| London Rocket[*]                            | <i>Sisymbrium irio</i>           |                       |                  |                            |
| Marestail/Horseweed                         | <i>Conyza canadensis</i>         |                       |                  |                            |
| Mayweed/False Chamomile[*]                  | <i>Matricaria maritima</i>       |                       |                  |                            |

[\*Not for use in California.]

continued

<sup>(1)</sup>Valor EZ Herbicide 3 at 4 fl oz/A (0.125 lb ai) will provide postemergence dodder suppression when applied in combination with Pursuit Herbicide or Raptor Herbicide at labeled rates. Pursuit Herbicide and Raptor Herbicide require the use of NIS, which will result in burn and stunting of alfalfa. Understand and accept these risks before tank mixing with Valor EZ Herbicide 3.

**Table 7. Weeds Controlled by Residual Activity of Valor EZ Herbicide 3 (continued)**

| <b>Broadleaf Weed Species</b>      |                                                    |                       |                  |                            |
|------------------------------------|----------------------------------------------------|-----------------------|------------------|----------------------------|
| <b>Common Name</b>                 | <b>Scientific Name</b>                             | <b>Organic Matter</b> | <b>Soil Type</b> | <b>Application Rate</b>    |
| Morningglories                     |                                                    | Up to 5%              | All Soil Types   | 4 fl oz/A<br>(0.125 lb ai) |
| Entireleaf                         | <i>Ipomoea hederacea</i> var. <i>integriuscula</i> |                       |                  |                            |
| Ivyleaf                            | <i>Ipomoea hederacea</i>                           |                       |                  |                            |
| Red/Scarlet                        | <i>Ipomoea coccinea</i>                            |                       |                  |                            |
| Smallflower                        | <i>Jacquemontia tamnifolia</i>                     |                       |                  |                            |
| Tall                               | <i>Ipomoea purpurea</i>                            |                       |                  |                            |
| Mustard                            |                                                    |                       |                  |                            |
| Tansy[*]                           | <i>Descurainia pinnata</i>                         |                       |                  |                            |
| Tumble[*]                          | <i>Sisymbrium altissimum</i>                       |                       |                  |                            |
| Wild                               | <i>Brassica kaber</i>                              |                       |                  |                            |
| Nettle, Burning[*]                 | <i>Urtica urens</i>                                |                       |                  |                            |
| Nightshades                        |                                                    |                       |                  |                            |
| Black                              | <i>Solanum nigrum</i>                              |                       |                  |                            |
| Eastern Black                      | <i>Solanum ptycanthum</i>                          |                       |                  |                            |
| Hairy                              | <i>Solanum sarrachoides</i>                        |                       |                  |                            |
| Pigweeds                           |                                                    |                       |                  |                            |
| Palmer Amaranth                    | <i>Amaranthus palmeri</i>                          |                       |                  |                            |
| Redroot                            | <i>Amaranthus retroflexus</i>                      |                       |                  |                            |
| Smooth                             | <i>Amaranthus hybridus</i>                         |                       |                  |                            |
| Spiny Amaranth                     | <i>Amaranthus spinosus</i>                         |                       |                  |                            |
| Tumble                             | <i>Amaranthus albus</i>                            |                       |                  |                            |
| Prickly Lettuce<br>(China Lettuce) | <i>Lactuca serriola</i>                            |                       |                  |                            |
| Prickly Sida (Teaweed)             | <i>Sida spinosa</i>                                |                       |                  |                            |
| Puncturevine                       | <i>Tribulus terrestris</i>                         |                       |                  |                            |
| Purslane                           |                                                    |                       |                  |                            |
| Common                             | <i>Portulaca oleracea</i>                          |                       |                  |                            |
| Horse[*]                           | <i>Trianthema portulacastrum</i>                   |                       |                  |                            |
| Radish, Wild                       | <i>Raphanus raphanistrum</i>                       |                       |                  |                            |
| Ragweed, Common                    | <i>Ambrosia artemisiifolia</i>                     |                       |                  |                            |
| Redmaids                           | <i>Calandrinia ciliata</i> var. <i>menziesii</i>   |                       |                  |                            |
| Russian Thistle                    | <i>Salsola iberica</i>                             |                       |                  |                            |
| Shepherd's-purse                   | <i>Capsella bursa-pastoris</i>                     |                       |                  |                            |
| Smartweeds                         |                                                    |                       |                  |                            |
| Ladysthumb                         | <i>Polygonum persicaria</i>                        |                       |                  |                            |
| Pennsylvania                       | <i>Polygonum pennsylvanicum</i>                    |                       |                  |                            |
| Smellmelon[*]                      | <i>Cucumis melo</i>                                |                       |                  |                            |
| Sowthistle, Prickly[*]             | <i>Sonchus asper</i>                               |                       |                  |                            |
| Spotted Spurge                     | <i>Euphorbia maculate</i>                          |                       |                  |                            |
| Spurred Anoda                      | <i>Anoda cristata</i>                              |                       |                  |                            |
| Tropic Croton                      | <i>Croton glandulosus</i>                          |                       |                  |                            |
| Velvetleaf                         | <i>Abutilon theophrasti</i>                        |                       |                  |                            |
| Venice Mallow                      | <i>Hibiscus trionum</i>                            |                       |                  |                            |
| Waterhemp                          |                                                    |                       |                  |                            |
| Common                             | <i>Amaranthus rudis</i>                            |                       |                  |                            |
| Tall                               | <i>Amaranthus tuberculatus</i>                     |                       |                  |                            |
| White Cockle[*]                    | <i>Silene latifolia</i>                            |                       |                  |                            |
| Wild Poinsettia                    | <i>Euphorbia heterophylla</i>                      |                       |                  |                            |
| Wormwood, Biennial                 | <i>Artemisia biennis</i>                           |                       |                  |                            |
| Yellow Rocket[*]                   | <i>Barbarea vulgaris</i>                           |                       |                  |                            |

[\*Not for use in California.]

continued

**Table 7. Weeds Controlled by Residual Activity of Valor EZ Herbicide 3 (continued)**

| Grass Weed Species     |                                | Organic Matter | Soil Type      | Application Rate           |
|------------------------|--------------------------------|----------------|----------------|----------------------------|
| Common Name            | Scientific Name                |                |                |                            |
| Barnyardgrass          | <i>Echinochloa crus-galli</i>  | Up to 5%       | All Soil Types | 4 fl oz/A<br>(0.125 lb ai) |
| Bluegrass, Annual      | <i>Poa annua</i>               |                |                |                            |
| Crabgrass, Large       | <i>Digitaria sanguinalis</i>   |                |                |                            |
| Foxtail, Giant         | <i>Setaria faberi</i>          |                |                |                            |
| Goosegrass             | <i>Eleusine indica</i>         |                |                |                            |
| Lovegrass, California  | <i>Eragrostis diffusa</i>      |                |                |                            |
| Panicums               |                                |                |                |                            |
| Fall                   | <i>Panicum dichotomiflorum</i> |                |                |                            |
| Texas                  | <i>Panicum texanum</i>         |                |                |                            |
| Ryegrass, Italian[*]   | <i>Lolium multiflorum</i>      |                |                |                            |
| Signalgrass, Broadleaf | <i>Brachiaria platyphylla</i>  |                |                |                            |

[\*Not for use in California.]

### **DIRECTIONS FOR USE IN ONION (DRY BULB)**

[For Use in Michigan, New York, North Dakota and Wisconsin Only]

#### **RESTRICTIONS AND LIMITATIONS**

- **DO NOT** apply more than 2 fl oz (0.063 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 6 applications of *Valor* EZ Herbicide 3 per acre per year at the 0.5 fl oz (0.016 lb ai) rate.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** make sequential application within 14 days of the first application.
- **DO NOT** apply more than 1 fl (0.032 lb ai) of *Valor* EZ Herbicide 3 per year on soils that contain greater than 90% sand plus gravel.
- **DO NOT** apply as part of a tank mix, other than Prowl® H<sub>2</sub>O Herbicide, or unacceptable injury may result. **DO NOT** tank mix other formulations of pendimethalin with *Valor* EZ Herbicide 3 for use in onions.
- **DO NOT** apply with any type of adjuvant.
- **DO NOT** apply within 45 days of harvest.

**Use of *Valor* EZ Herbicide 3 may result in necrotic spotting of onion leaves that come in contact with the spray. Understand and accept this risk before using *Valor* EZ Herbicide 3.**

#### **[Microrate Application]**

[Sequential applications of *Valor* EZ Herbicide 3 may be applied to onions (dry bulb), between the 2-leaf and 6-leaf stage, at rates of 0.5 to 1 fl oz/A (0.016 to 0.032 lb ai), on a 7 day interval.]

#### **TIMING TO ONIONS (dry bulb)**

Apply *Valor* EZ Herbicide 3 to transplanted onions (dry bulb) between the 2-leaf and 6-leaf stage and on direct seed onions (dry bulb) between the 3-leaf and 6-leaf stage.

#### **TIMING TO WEEDS**

##### **Preemergence – Emerged Onions (dry bulb), Preemergence To Weeds**

Apply *Valor* EZ Herbicide 3 to weed free onions (dry bulb) for preemergence control of the weeds listed in Table 1, Section A, *Broadleaf Weeds Controlled by Residual Activity to Valor EZ Herbicide 3*.

## DIRECTIONS FOR USE IN PEANUT

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** apply more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** irrigate when peanuts are cracking.
- **DO NOT** graze treated fields or feed treated hay to livestock.

Many weather-related factors, including high wind, splashing or heavy rains or cool conditions at or near peanut emergence, may result in peanut injury in fields treated with *Valor* EZ Herbicide 3. On occasion this has resulted in a delay in maturity or even a slight decrease in yield.

### WIND MANAGEMENT

In areas where shallow cultivation is used between rows to reduce wind-borne sand damage to peanuts, weed control from *Valor* EZ Herbicide 3 may be reduced.

### TIMING TO PEANUTS

*Valor* EZ Herbicide 3 may be applied to peanuts prior to planting or preemergence (after planting). Preemergence applications of *Valor* EZ Herbicide 3 must be made within 2 days after planting and prior to peanut emergence. Application after the peanuts have begun to crack, or are emerged, will result in severe crop injury. Apply before peanuts have begun to crack. Select *Valor* EZ Herbicide 3 rate from Table 1 *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3* according to anticipated weed spectrum.

### TIMING TO WEEDS

#### Burndown – Preemergence to Peanuts, Postemergence to Weeds

*Valor* EZ Herbicide 3, applied as part of a burndown program, may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds where peanuts will be planted directly into a stale seedbed, cover crop or in previous crop residues. Apply *Valor* EZ Herbicide 3 before planting, during planting or after planting, but before the crop emerges. For control of emerged weeds, tank mix *Valor* EZ Herbicide 3 with glyphosate. Refer to glyphosate label for rates and application pressure. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. *Valor* EZ Herbicide 3 tank mixes applied to assist in the control of emerged weeds must be applied with an adjuvant, including a non-ionic surfactant at 0.25% v/v or a crop oil concentrate or a methylated seed oil at 1 to 2 pt/A. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lb/A or 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to increase herbicidal activity.

Preemergence (conventional tillage) applications of *Valor* EZ Herbicide 3 must be applied prior to weed emergence.

### ADDITIONAL RESIDUAL GRASS CONTROL: SEQUENTIAL

*Valor* EZ Herbicide 3 may be applied sequentially following a preplant incorporated application of trifluralin (states of New Mexico, Oklahoma and Texas only), Sonalan®, Dual® (metolachlor), pendimethalin or Frontier®.

### ADDITIONAL RESIDUAL GRASS CONTROL: TANK MIXED

*Valor* EZ Herbicide 3 can be tank mixed with alachlor, metolachlor or Frontier for additional grass and broadleaf weed control. *Valor* EZ Herbicide 3 can also be tank mixed with pendimethalin or Sonalan® in states where they are labeled, provided overhead irrigation guidelines on the pendimethalin and/or Sonalan® labels are followed.

## DIRECTIONS FOR USE IN POTATO

[For Use in Arizona, California, Colorado, Delaware, Florida, Hawaii, Idaho, Maryland, Minnesota, Montana, Nebraska, Nevada, New Jersey, New Mexico, North Carolina, North Dakota, Oregon, South Carolina, South Dakota, Texas, Utah, Virginia, Washington, Washington DC and Wyoming only.]

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 1.5 fl oz (0.047 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** apply more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 1.5 fl oz (0.047 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply to Rill (furrow) irrigated potatoes.

**Many weather-related factors, including high wind, splashing or heavy rains or cool conditions at or near potato emergence, may result in potato injury in fields treated with *Valor* EZ Herbicide 3. On occasion this has resulted in a delay in maturity. Understand and accept these risks before using *Valor* EZ Herbicide 3.**

### TIMING TO POTATOES

*Valor* EZ Herbicide 3 may be applied to potatoes after hilling for the preemergence suppression of the weeds listed in Table 8, *Weeds Suppressed by Residual Activity of Valor EZ Herbicide 3 at 1.5 fl oz/A*. Tank mix *Valor* EZ Herbicide 3 with other labeled herbicides for broad spectrum weed control. A minimum of 2 inches of settled soil must cover the vegetative portion of the potato plant at the time of *Valor* EZ Herbicide 3 application. Application to potatoes with less than 2 inches of soil covering the vegetative portion of the potato may result in crop injury. In areas with historically higher amounts of rainfall during the time of preemergence herbicide applications, including the Red River Valley, Minnesota and North Dakota, the requirement for 2 inches of settled soil is critical to avoid crop injury. Mechanical incorporation of *Valor* EZ Herbicide 3 will result in decreased weed control. In areas with sprinkler irrigation, incorporate *Valor* EZ Herbicide 3 with 1/4 to 3/4 inches of irrigation, after application and before any sprouts are within 2 inches of the settled soil surface if a rainfall event has not yet occurred.

### TIMING TO WEEDS

#### Preemergence – Soil Covered Potatoes, Preemergence to Weeds

Apply *Valor* EZ Herbicide 3 to soil covered potatoes for the preemergence suppression of the weeds listed in Table 8. Harrowing, cultivation or corrugating after *Valor* EZ Herbicide 3 application will reduce weed control.

**Table 8. Weeds Suppressed by Residual Activity of *Valor* EZ Herbicide 3 at 1.5 fl oz/A**

| Common Name                        | Scientific Name               | Organic Matter | Application Rate             |
|------------------------------------|-------------------------------|----------------|------------------------------|
| Lambsquarters, Common              | <i>Chenopodium album</i>      | Up to 5%       | 1.5 fl oz/A<br>(0.047 lb ai) |
| Mustard, Wild                      | <i>Brassica kaber</i>         |                |                              |
| Nightshades                        |                               |                |                              |
| Black                              | <i>Solanum nigrum</i>         |                |                              |
| Eastern Black                      | <i>Solanum ptycanthum</i>     |                |                              |
| Hairy                              | <i>Solanum sarrachoides</i>   |                |                              |
| Pigweeds                           |                               |                |                              |
| Palmer Amaranth                    | <i>Amaranthus palmeri</i>     |                |                              |
| Redroot                            | <i>Amaranthus retroflexus</i> |                |                              |
| Smooth                             | <i>Amaranthus hybridus</i>    |                |                              |
| Spiny Amaranth                     | <i>Amaranthus spinosus</i>    |                |                              |
| Tumble                             | <i>Amaranthus albus</i>       |                |                              |
| Prickly Lettuce<br>(China Lettuce) | <i>Lactuca serriola</i>       |                |                              |
| Radish, Wild                       | <i>Raphanus raphanistrum</i>  |                |                              |

## DIRECTIONS FOR USE IN SOYBEAN

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** apply more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- Graze treated fields or feed treated hay to livestock no sooner than 21 days after application.

### PRECAUTIONS

- If *Valor* EZ Herbicide 3 is tank mixed with flufenacet (Axiom®, Domain®), metolachlor (Dual Magnum, Dual II Magnum, Boundary®) or dimethenamid (Frontier or Outlook®) and applied within 14 days of planting soybeans, plant under no-till or minimum tillage conditions on wheat stubble or field corn stubble.
- Irrigation when soybeans are cracking may result in severe injury.

### TIMING TO SOYBEANS

*Valor* EZ Herbicide 3 may be applied to soybeans prior to planting or preemergence (after planting). Preemergence application of *Valor* EZ Herbicide 3 must be made within 3 days after planting and prior to soybean emergence. Application after the soybeans have begun to crack, or are emerged, will result in severe crop injury. Apply before soybeans have begun to crack. Select *Valor* EZ Herbicide 3 rate from Table 1 *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3* according to anticipated weed spectrum.

### TIMING TO WEEDS

#### Burndown – Preemergence to Soybeans, Postemergence to Weeds

*Valor* EZ Herbicide 3, applied as part of a burndown program, may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds where soybeans will be planted directly into a stale seedbed, cover crop or in previous crop residues. For control of emerged weeds, choose the most appropriate tank mix partner from Table 9. *Tank Mix Partners for Control of Emerged Weeds in Reduced Tillage Soybeans*. Apply *Valor* EZ Herbicide 3 with ground equipment before planting, during planting or within 3 days after planting, **but before the crop emerges**. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Refer to tank mix partner's label for rates and application pressures. All *Valor* EZ Herbicide 3 tank mixes applied to assist in the control of emerged weeds must be applied with crop oil concentrate or methylated seed oil at 1 to 2 pt/A or a non-ionic surfactant at 0.25% v/v.

#### INCREASING SPEED OF GLYPHOSATE BURNDOWN ACTIVITY

*Valor* EZ Herbicide 3, at rates as low as 1 fl oz/A (0.032 lb ai), may be tank mixed with glyphosate (Roundup) to increase the speed of burndown activity compared to glyphosate applied alone. Residual weed control will not be provided at rates lower than 2 fl oz/A (0.063 lb ai); however, suppression of the weeds in Table 2. *Weeds Suppressed by Residual Activity of Valor EZ Herbicide 3 Application Rates* may occur at *Valor* EZ Herbicide 3 rates as low as 1 fl oz/A (0.032 lb ai).

### TANK MIXES

*Valor* EZ Herbicide 3 may be tank mixed with the herbicides listed in Table 9. *Tank Mix Partners for Control of Emerged Weeds in Reduced Tillage Soybeans* for increased burndown activity, additional residual broadleaf and/or additional grass control. Refer to tank mix partner's label for adjuvant selection.

**Table 9. Tank Mix Partners for Control of Emerged Weeds in Reduced Tillage Soybeans**

| <b>Tank Mix Partner</b>    | <b>Target Weeds <sup>(1)</sup></b>      |
|----------------------------|-----------------------------------------|
| 2,4-D LVE                  | Marestail<br>Giant Ragweed<br>Dandelion |
| paraquat                   | Annual Grasses<br>Henbit                |
| glyphosate                 | General Burndown                        |
| Select Max <sup>®</sup>    | Annual Grasses                          |
| Scepter <sup>®</sup> 70 DG | Cocklebur<br>Common Sunflower           |
| Weedmaster <sup>®</sup>    | Marestail<br>Giant Ragweed<br>Dandelion |

<sup>(1)</sup>Refer to tank mix product labels for specific use directions for control of emerged weeds present.

#### **ADDITIONAL RESIDUAL BROADLEAF CONTROL**

*Valor* EZ Herbicide 3 can be tank mixed with metribuzin, Firstrate<sup>®</sup>, Lorox<sup>®</sup>, Pursuit Plus<sup>®</sup>, PYTHON<sup>®</sup>, Squadron<sup>®</sup>, Scepter or Steel<sup>®</sup> for additional broadleaf control.

#### **ADDITIONAL RESIDUAL GRASS CONTROL**

*Valor* EZ Herbicide 3 can be tank mixed with pendimethalin or Command<sup>®</sup> for additional grass control. [In the states of **[Note to EPA Reviewer: specific states will be listed here]** *Valor* EZ Herbicide 3 can be tank mixed with microencapsulated acetochlor (Warrant<sup>®</sup>) at 2 fl oz per acre.] [Tank mixes with flufenacet (Axiom or Domain), metolachlor (Dual products or Boundary) or dimethenamid (Frontier or Outlook) may result in severe injury to soybeans when application is followed by prolonged periods of cool wet weather.]

#### **ROUNDUP READY PROGRAM**

*Valor* EZ Herbicide 3 may be applied as part of a burndown program or preemergence in conventional tillage programs, at 2 to 3 fl oz/A (0.063 to 0.094 lb ai) to reduce early season weed competition from waterhemp, velvetleaf, nightshade and morningglories as well as other weeds listed in Tables 2 and 3 in Roundup Ready programs. A sequential post emergence application of glyphosate will be required to control weeds not controlled by *Valor* EZ Herbicide 3.

### **DIRECTIONS FOR USE IN STRAWBERRY**

#### **RESTRICTIONS AND LIMITATIONS:**

- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** apply more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.

#### **PRECAUTIONS**

- *Valor* EZ Herbicide 3, at 3 fl oz (0.094 lb ai) per acre, can be applied to the soil a minimum of 30 days prior to transplanting strawberries provided the strawberries will be transplanted through a plastic mulch.
- *Valor* EZ Herbicide 3 at 3 fl oz (0.094 lb ai) per acre can be applied to dormant (established or newly planted) strawberries for the preemergence control of the weeds listed in Table 1, *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*.
- *Valor* EZ Herbicide 3, at 3 fl oz (0.094 lb ai) per acre, can be applied in strawberry row middles with a shielded or hooded sprayer for the preemergence control of the weeds listed in Table 1, *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*.

| Application Method                                    | Minimum Time From Application to Harvest (PHI) | Use Rate Per Acre Per Application (oz) | Use Rate Per Acre Per Year (oz) | Special Use Instructions                                                                                                                                                                                                                                                                                                    |
|-------------------------------------------------------|------------------------------------------------|----------------------------------------|---------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Pre-transplant                                        | Not applicable                                 | 3<br>(0.094 lb ai)                     | 3<br>(0.094 lb ai)              | Apply a minimum of 30 days prior to transplanting and prior to plastic mulch being laid.<br><br>Apply as part of a tank mix to control emerged weeds.                                                                                                                                                                       |
| Preemergence to dormant strawberries                  | Not applicable                                 | 3<br>(0.094 lb ai)                     | 3<br>(0.094 lb ai)              | Crop oil concentrate, at 1% v/v, or non-ionic surfactant, at 0.25% v/v, may be added to help control emerged broadleaf weeds.                                                                                                                                                                                               |
| Hooded or shielded sprayer application to row middles | <b>DO NOT</b> apply after fruit set            | 3<br>(0.094 lb ai)                     | 3<br>(0.094 lb ai)              | <b>Apply only to row middles - DO NOT apply over strawberries.</b><br><br>Apply prior to weed emergence.<br><br>Crop spotting may occur if an adjuvant is added.<br><br><b>DO NOT apply after fruit set or spotting of fruit may occur.</b><br><br><b>DO NOT</b> allow spray drift to come in contact with fruit or foliage |

**Table 10. Weeds Controlled by Preemergence Application of Valor EZ Herbicide 3**

| Broadleaf Weed Species     |                                                    |                          |                               |                                                       |
|----------------------------|----------------------------------------------------|--------------------------|-------------------------------|-------------------------------------------------------|
| Common Name                | Scientific Name                                    | Organic Matter           | Soil Type                     | Application Rates                                     |
| Bristly Starbur            | <i>Acanthospermum hispidum</i>                     | Up to 10% <sup>(1)</sup> | All Soil Types <sup>(2)</sup> | 6 to 12 fl oz/A (0.188 to 0.375 lb ai) <sup>(3)</sup> |
| Carpetweed                 | <i>Mollugo verticillata</i>                        |                          |                               |                                                       |
| Chickweeds                 |                                                    |                          |                               |                                                       |
| Common                     | <i>Stellaria media</i>                             |                          |                               |                                                       |
| Mouseear                   | <i>Cerastium vulgatum</i>                          |                          |                               |                                                       |
| Coffee Senna               | <i>Cassia occidentalis</i>                         |                          |                               |                                                       |
| Dandelion                  | <i>Taraxacum officinale</i>                        |                          |                               |                                                       |
| Eclipta                    | <i>Eclipta prostrata</i>                           |                          |                               |                                                       |
| Evening primrose, Cutleaf  | <i>Oenothera laciniata</i>                         |                          |                               |                                                       |
| False Chamomile[*]         | <i>Tripleurospermum maritima</i>                   |                          |                               |                                                       |
| Filaree                    |                                                    |                          |                               |                                                       |
| Redstem                    | <i>Erodium cicutarium</i>                          |                          |                               |                                                       |
| Whitestem                  | <i>Erodium moschatum</i>                           |                          |                               |                                                       |
| Fiddleneck, Coast[*]       | <i>Amsinckia menziesii</i>                         |                          |                               |                                                       |
| Fleabane, Hairy            | <i>Conyza bonariensis</i>                          |                          |                               |                                                       |
| Field Pennycress[*]        | <i>Thlaspi arvense</i>                             |                          |                               |                                                       |
| Florida Beggarweed         | <i>Desmodium tortuosum</i>                         |                          |                               |                                                       |
| Florida Pusley             | <i>Richardia scabra</i>                            |                          |                               |                                                       |
| Golden Crownbeard          | <i>Verbesina encelioides</i>                       |                          |                               |                                                       |
| Groundsel, Common          | <i>Senecio vulgaris</i>                            |                          |                               |                                                       |
| Hairy Indigo               | <i>Indigofera hirsuta</i>                          |                          |                               |                                                       |
| Hemp Sesbania              | <i>Sesbania exaltata</i>                           |                          |                               |                                                       |
| Henbit                     | <i>Lamium amplexicaule</i>                         |                          |                               |                                                       |
| Jimsonweed                 | <i>Datura stramonium</i>                           |                          |                               |                                                       |
| Kochia                     | <i>Kochia scoparia</i>                             |                          |                               |                                                       |
| Lambsquarters, Common      | <i>Chenopodium album</i>                           |                          |                               |                                                       |
| Mallow                     |                                                    |                          |                               |                                                       |
| Common (Cheeseweed)        | <i>Malva neglecta</i>                              |                          |                               |                                                       |
| Little                     |                                                    |                          |                               |                                                       |
| Horseweed/Marestail        | <i>Conyza canadensis</i>                           |                          |                               |                                                       |
| Mayweed/False Chamomile[*] | <i>Matricaria maritima</i>                         |                          |                               |                                                       |
| Morningglories             |                                                    |                          |                               |                                                       |
| Entireleaf                 | <i>Ipomoea hederacea</i> var. <i>integriuscula</i> |                          |                               |                                                       |
| Ivyleaf                    |                                                    |                          |                               |                                                       |
| Red/Scarlet                | <i>Ipomoea coccinea</i>                            |                          |                               |                                                       |
| Smallflower                | <i>Jacquemontia tamnifolia</i>                     |                          |                               |                                                       |
| Tall                       | <i>Ipomoea purpurea</i>                            |                          |                               |                                                       |

[\*Not for use in California.]

continued

<sup>(1)</sup>Valor EZ Herbicide 3 can be used on soils with greater than 10% organic matter; however, length of residual control may be shorter than on soils with lower organic matter content.

<sup>(2)</sup>Use a maximum Valor EZ Herbicide 3 rate of 6 fl oz/A (0.188 lb ai) per application on any soil that has a sand plus gravel content over 80% if bushes, trees or vines are under 3 years of age.

<sup>(3)</sup>See specific crop instructions for application rate

**Table 10. Weeds Controlled by Preemergence Application of Valor EZ Herbicide 3 (continued)**

| Broadleaf Weed Species             |                                                   |                          |                               |                                                          |
|------------------------------------|---------------------------------------------------|--------------------------|-------------------------------|----------------------------------------------------------|
| Common Name                        | Scientific Name                                   | Organic Matter           | Soil Type                     | Application Rates                                        |
| Mustards                           |                                                   | Up to 10% <sup>(1)</sup> | All Soil Types <sup>(2)</sup> | 6 to 12 fl oz/A <sup>(3)</sup><br>(0.188 to 0.375 lb ai) |
| London Rocket[*]                   | <i>Sisymbrium irio</i>                            |                          |                               |                                                          |
| Tansey[*]                          | <i>Descurainia pinnata</i>                        |                          |                               |                                                          |
| Tumble                             | <i>Sisymbrium altissimum</i>                      |                          |                               |                                                          |
| Wild                               | <i>Brassica kaber</i>                             |                          |                               |                                                          |
| Nettle, Burning[*]                 | <i>Urtica urens</i>                               |                          |                               |                                                          |
| Nightshades                        |                                                   |                          |                               |                                                          |
| Black                              | <i>Solanum nigrum</i>                             |                          |                               |                                                          |
| Eastern Black                      | <i>Solanum ptycanthum</i>                         |                          |                               |                                                          |
| Hairy                              | <i>Solanum sarrachoides</i>                       |                          |                               |                                                          |
| Pigweeds                           |                                                   |                          |                               |                                                          |
| Palmer Amaranth                    | <i>Amaranthus palmeri</i>                         |                          |                               |                                                          |
| Redroot                            | <i>Amaranthus retroflexus</i>                     |                          |                               |                                                          |
| Smooth                             | <i>Amaranthus hybridus</i>                        |                          |                               |                                                          |
| Spiny Amaranth                     | <i>Amaranthus spinosus</i>                        |                          |                               |                                                          |
| Tumble                             | <i>Amaranthus albus</i>                           |                          |                               |                                                          |
| Prickly Lettuce<br>(China Lettuce) | <i>Lactuca serriola</i>                           |                          |                               |                                                          |
| Prickly Sida (Teaweed)             | <i>Sida spinosa</i>                               |                          |                               |                                                          |
| Puncturevine                       | <i>Tribulus terrestris</i>                        |                          |                               |                                                          |
| Purslane                           |                                                   |                          |                               |                                                          |
| Common                             | <i>Portulaca oleracea</i>                         |                          |                               |                                                          |
| Horse[*]                           | <i>Trianthema portulacastrum</i>                  |                          |                               |                                                          |
| Radish, Wild                       | <i>Raphanus raphanistrum</i>                      |                          |                               |                                                          |
| Ragweed, Common                    | <i>Ambrosia artemisiifolia</i>                    |                          |                               |                                                          |
| Redmaids                           | <i>Calandrinia ciliata</i> var <i>menziesii</i> . |                          |                               |                                                          |
| Redweed                            | <i>Melochia corchorifolia</i>                     |                          |                               |                                                          |
| Shepherd's-purse                   | <i>Capsella bursa-pastoris</i>                    |                          |                               |                                                          |
| Smellmelon[*]                      | <i>Cucumis melo</i>                               |                          |                               |                                                          |
| Sowthistle, Annual                 | <i>Sonchus oleraceus</i>                          |                          |                               |                                                          |
| Spotted Spurge                     | <i>Euphorbia maculata</i>                         |                          |                               |                                                          |
| Spurred Anoda                      | <i>Anoda cristata</i>                             |                          |                               |                                                          |
| Thistle, Russian                   | <i>Salsola iberica</i>                            |                          |                               |                                                          |
| Tropic Croton                      | <i>Croton glandulosus</i>                         |                          |                               |                                                          |
| Venice Mallow                      | <i>Hibiscus trionum</i>                           |                          |                               |                                                          |
| Waterhemp                          |                                                   |                          |                               |                                                          |
| Common                             | <i>Amaranthus rudis</i>                           |                          |                               |                                                          |
| Tall                               | <i>Amaranthus tuberculatus</i>                    |                          |                               |                                                          |
| Wild Poinsettia                    | <i>Euphorbia heterophylla</i>                     |                          |                               |                                                          |
| White Cockle[*]                    | <i>Silene latifolia</i>                           |                          |                               |                                                          |
| Wormwood, Biennial                 | <i>Artemisia biennis</i>                          |                          |                               |                                                          |
| Yellow Rocket[*]                   | <i>Barbarea vulgaris</i>                          |                          |                               |                                                          |

[\*Not for use in California.]

continued

<sup>(1)</sup>Valor EZ Herbicide 3 can be used on soils with greater than 10% organic matter; however, length of residual

control may be shorter than on soils with lower organic matter content.

<sup>(2)</sup>Use a maximum Valor EZ Herbicide 3 rate of 6 fl oz/A (0.188 lb ai) per application on any soil that has a sand plus gravel content over 80% if bushes, trees or vines are under 3 years of age.

<sup>(3)</sup>See specific crop instructions for application rate

**Table 10. Weeds Controlled by Preemergence Application of Valor EZ Herbicide 3 (continued)**

| <b>Broadleaf Weed Species</b> |                                |                          |                               |                                                          |
|-------------------------------|--------------------------------|--------------------------|-------------------------------|----------------------------------------------------------|
| <b>Common Name</b>            | <b>Scientific Name</b>         | <b>Organic Matter</b>    | <b>Soil Type</b>              | <b>Application Rates</b>                                 |
| <b>GRASS WEED SPECIES</b>     |                                | Up to 10% <sup>(1)</sup> | All Soil Types <sup>(2)</sup> | 6 to 12 fl oz/A <sup>(3)</sup><br>(0.188 to 0.375 lb ai) |
| Barnyardgrass                 | <i>Echinochloa crus-galli</i>  |                          |                               |                                                          |
| Bluegrass, Annual             | <i>Poa annua</i>               |                          |                               |                                                          |
| Crabgrass                     |                                |                          |                               |                                                          |
| Large                         | <i>Digitaria sanguinalis</i>   |                          |                               |                                                          |
| Smooth                        | <i>Digitaria ischaemum</i>     |                          |                               |                                                          |
| Foxtails                      |                                |                          |                               |                                                          |
| Bristly                       | <i>Setaria verticillata</i>    |                          |                               |                                                          |
| Giant                         | <i>Setaria faberi</i>          |                          |                               |                                                          |
| Green                         | <i>Setaria viridis</i>         |                          |                               |                                                          |
| Yellow                        | <i>Setaria glauca</i>          |                          |                               |                                                          |
| Goosegrass                    | <i>Eleusine indica</i>         |                          |                               |                                                          |
| Guineagrass                   | <i>Panicum maximum</i>         |                          |                               |                                                          |
| Johnsongrass, Seedling        | <i>Sorghum halepense</i>       |                          |                               |                                                          |
| Lovegrass, California         | <i>Eragrostis diffusa</i>      |                          |                               |                                                          |
| Panicum                       |                                |                          |                               |                                                          |
| Fall                          | <i>Panicum dichotomiflorum</i> |                          |                               |                                                          |
| Texas                         | <i>Panicum texanum</i>         |                          |                               |                                                          |
| Ryegrass, Italian[*]          | <i>Lolium multiflorum</i>      |                          |                               |                                                          |
| Signalgrass, Broadleaf        | <i>Brachiaria platyphylla</i>  |                          |                               |                                                          |

[\*Not for use in California.]

<sup>(1)</sup>Valor EZ Herbicide 3 can be used on soils with greater than 10%; however, length of residual control may be shorter than on soils with lower organic matter content.

<sup>(2)</sup>Use a maximum Valor EZ Herbicide 3 rate of 6 fl oz/A (0.188 lb ai) per application on any soil that has a sand plus gravel content over 80% if bushes, trees or vines are under 3 years of age.

<sup>(3)</sup>See specific crop instructions for application rate.

## DIRECTIONS FOR USE IN SUGARCANE

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 8 fl oz (0.250 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make a sequential application within 14 days of the first application.
- **DO NOT** apply more than 4 applications of *Valor* EZ Herbicide 3 per acre per year at the 3 fl oz (0.094 lb flumioxazin) rate.
- **DO NOT** apply more than 12 fl oz (0.375 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply within 90 days of harvest.

### TIMING TO SUGARCANE

*Valor* EZ Herbicide 3 may be applied from 2 weeks prior to planting to before the sugarcane emerges, post directed or at layby. Select the proper *Valor* EZ Herbicide 3 rate from Table 10 according to anticipated weed spectrum and soil organic matter content for preemergence applications. Select *Valor* EZ Herbicide 3 rate from Table 11. *Broadleaf Weeds Controlled by Post-Directed or Layby Application of Valor EZ Herbicide 3 in Sugarcane* according to emerged weed spectrum and weed heights for post-directed and layby applications.

### TIMING TO WEEDS

#### Burndown – Preemergence to Sugarcane, Postemergence to Weeds

*Valor* EZ Herbicide 3 may be used for preemergence control, and to assist in postemergence burndown, of many annual broadleaf weeds in sugarcane. For control of emerged weeds, choose the most appropriate tank mix partner from Table 12. *Tank Mixes with Valor EZ Herbicide 3 for Post-Directed or Layby Use in Sugarcane*. Apply *Valor* EZ Herbicide 3 **before the crop emerges**. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. All *Valor* EZ Herbicide 3 tank mixes applied to assist in the control of emerged weeds must be applied with crop oil concentrate or methylated seed oil at 1 qt/A or a non-ionic surfactant at 0.25% v/v. Some tank mix products, for example Roundup Original Max (glyphosate), may be formulated with a suitable adjuvant and do not require additional adjuvant.

#### Preemergence – Preemergence to Sugarcane, Preemergence to Weeds

*Valor* EZ Herbicide 3 may be used for preemergence control of many annual broadleaf and grassy weeds in sugarcane. Select rate based on anticipated weed spectrum and soil organic matter content from Table 10. Apply *Valor* EZ Herbicide 3 **before the crop emerges**.

#### Post-Directed – Postemergence to Sugarcane, Postemergence to Weeds

Make post-directed applications to upright sugarcane varieties after the sugarcane has exceeded 24 inches in height and has begun to joint. Post-directed applications to “PINEAPPLE” varieties or to upright varieties that are less than 24 inches in height and have not begun to joint, may result in unacceptable crop injury. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Post-directed applications of *Valor* EZ Herbicide 3 must include a crop oil concentrate or methylated seed oil at 1 qt/A or a non-ionic surfactant at 0.25% v/v. Select the proper *Valor* EZ Herbicide 3 rate based on weed spectrum and weed height from Table 11. *Broadleaf Weeds Controlled by Post-Directed or Layby Application of Valor EZ Herbicide 3 in Sugarcane*.

#### Layby – Postemergence to Sugarcane, Postemergence to Weeds

Layby applications can be made to upright and “PINEAPPLE” varieties after the sugarcane has exceeded 30 inches in height and the spray solution will not contact foliage above 6 inches from the base of the sugarcane. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Layby applications of *Valor* EZ Herbicide 3 must be applied with crop oil concentrate or methylated seed oil at 1 qt/A or a non-ionic surfactant at 0.25% v/v. Select the proper *Valor* EZ Herbicide 3 rate based on weed spectrum and weed height from Table 11. *Broadleaf Weeds Controlled by Post-Directed or Layby Application of Valor EZ Herbicide 3 in Sugarcane*.

**Table 11. Broadleaf Weeds Controlled by Post-Directed or Layby Application of Valor EZ Herbicide 3 in Sugarcane**

| Broadleaf Weed Species         |                                                    | Weed Height (inches)       |                            |
|--------------------------------|----------------------------------------------------|----------------------------|----------------------------|
| Common Name                    | Scientific Name                                    | 3 fl oz/A<br>(0.094 lb ai) | 4 fl oz/A<br>(0.125 lb ai) |
| Bindweed, Field <sup>(1)</sup> | <i>Convolvulus arvensis</i>                        | 4                          | 8                          |
| Carpetweed                     | <i>Mollugo verticillata</i>                        | 4                          | 4                          |
| Cocklebur, Common              | <i>Xanthium strumarium</i>                         | 4                          | 4                          |
| Florida Beggarweed             | <i>Desmodium tortuosum</i>                         | 2                          | 2                          |
| Hemp Sesbania                  | <i>Sesbania exaltata</i>                           | 6                          | 8                          |
| Jimsonweed                     | <i>Datura stramonium</i>                           | 4                          | 4                          |
| Lambsquarters, Common          | <i>Chenopodium album</i>                           | 4                          | 4                          |
| Morningglories                 |                                                    |                            |                            |
| Entireleaf                     | <i>Ipomoea hederacea</i> var. <i>integriuscula</i> | -                          | 4                          |
| Ivyleaf                        | <i>Ipomoea hederacea</i>                           | 4                          | 4                          |
| Pitted                         | <i>Ipomoea lacunosa</i>                            | 4                          | 6                          |
| Red                            | <i>Ipomoea coccinea</i>                            | -                          | 4                          |
| Tall                           | <i>Ipomoea purpurea</i>                            | 2                          | 4                          |
| Mustard, Wild                  | <i>Brassica kaber</i>                              | 6                          | 6                          |
| Pigweeds                       |                                                    |                            |                            |
| Palmer Amaranth                | <i>Amaranthus palmeri</i>                          | 4                          | 6                          |
| Redroot                        | <i>Amaranthus retroflexus</i>                      | 4                          | 6                          |
| Smooth                         | <i>Amaranthus hybridus</i>                         | 4                          | 6                          |
| Plantain, Broadleaf            | <i>Plantago major</i>                              | 6                          | 6                          |
| Prickly Sida                   | <i>Sida spinosa</i>                                | 4                          | 6                          |
| Purslanes                      |                                                    |                            |                            |
| Common                         | <i>Portulaca oleracea</i>                          | 2                          | 4                          |
| Rock                           | <i>Calandrinia</i> spp.                            | -                          | 2                          |
| Ragweeds                       |                                                    |                            |                            |
| Common                         | <i>Ambrosia artemisiifolia</i>                     | 2                          | 2                          |
| Giant                          | <i>Ambrosia trifida</i>                            | 4                          | 4                          |
| Rice Flatsedge                 | <i>Cyperus iria</i>                                | 2                          | 4                          |
| Sicklepod                      | <i>Senna obtusifolia</i>                           | 4                          | 4                          |
| Smartweeds                     |                                                    |                            |                            |
| Ladysthumb                     | <i>Polygonum persicaria</i>                        | 4                          | 4                          |
| Pale                           | <i>Polygonum lapathifolium</i>                     | 4                          | 4                          |
| Pennsylvania                   | <i>Polygonum pennsylvanicum</i>                    | 4                          | 4                          |
| Spotted Spurge                 | <i>Euphorbia maculata</i>                          | 4                          | 4                          |
| Velvetleaf                     | <i>Abutilon theophrasti</i>                        | 4                          | 6                          |
| Venice Mallow                  | <i>Hibiscus trionum</i>                            | 2                          | 2                          |
| Waterhemp                      |                                                    |                            |                            |
| Common                         | <i>Amaranthus rudis</i>                            | 2                          | 2                          |
| Tall                           | <i>Amaranthus tuberculatus</i>                     | 2                          | 2                          |

<sup>(1)</sup>Valor EZ Herbicide 3 tank mixes will only control the above ground portion of field bindweed. Repeated applications will be needed to control regrowth.

## TANK MIXES

Valor EZ Herbicide 3 may be tank mixed with the herbicides listed in Table 12 for additional weed control in burndown, preemergence, post-directed and layby applications. Refer to tank mix partner's label for adjuvants.

**Table 12. Tank Mixes with Valor EZ Herbicide 3 for Post-Directed or Layby Use in Sugarcane**

| Tank Mix Partner <sup>(1)</sup> | Target Weeds                         | Burndown | Post-Directed <sup>(2)</sup> | Layby |
|---------------------------------|--------------------------------------|----------|------------------------------|-------|
| 2,4-D amine                     | Annual and Perennial Broadleaf Weeds | X        |                              |       |
| atrazine                        | Pigweeds<br>Cocklebur                | X        | X                            | X     |
| Asulox <sup>®(3)</sup>          | Annual Grasses                       |          | X                            | X     |
| Evik <sup>®(4)</sup>            | Annual Grasses                       |          | X                            | X     |
| glyphosate <sup>(5)</sup>       | Annual and Perennial Weeds           | X        |                              | X     |
| metribuzin <sup>(6)</sup>       | Broadleaf Panicum<br>Goosegrass      |          | X                            | X     |
| Sempre <sup>®</sup>             | Purple Nutsedge<br>Yellow Nutsedge   | X        | X                            | X     |
| Weedmaster                      | Annual and Perennial Broadleaf Weeds | X        |                              |       |

<sup>(1)</sup>Refer to tank mix product labels for specific use directions for control of emerged weeds present not listed in Table 11. *Broadleaf Weeds Controlled by Post-Directed or Layby Application of Valor EZ Herbicide 3 in Sugarcane.*

<sup>(2)</sup>Make post-directed applications to upright sugarcane varieties after the sugarcane has exceeded 24 inches in height. Post-directed applications to "PINEAPPLE" varieties or to upright varieties that are less than 24 inches in height may result in unacceptable crop injury.

<sup>(3)</sup>Apply to sugarcane at least 24 inches tall.

<sup>(4)</sup>Apply before weeds are greater than 6 inches tall.

<sup>(5)</sup>Glyphosate applications must be made with a hooded sprayer. Sugarcane must be at least 3 feet tall. Contact with the sugarcane foliage by either the spray mixture or the treated weed foliage will result in sugarcane injury.

<sup>(6)</sup>Refer to metribuzin label for restrictions based on soil type.

## ADDITIONAL PREEMERGENCE BROADLEAF CONTROL

Valor EZ Herbicide 3 can be tank mixed with atrazine or diuron for additional preemergence broadleaf control.

## ADDITIONAL PREEMERGENCE GRASS CONTROL

Valor EZ Herbicide 3 can be tank mixed with Prowl (or other pendimethalin products) for additional preemergence grass control provided sugarcane has not emerged.

## DIRECTIONS FOR USE IN SUNFLOWER AND SAFFLOWER

### HARVEST AID

#### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** apply more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** harvest within 5 days of application.

Desiccation from *Valor* EZ Herbicide 3 requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lb/A or a 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* EZ Herbicide 3 with glyphosate or paraquat will increase control of emerged weeds and aid in harvest for sunflowers. Tank mixing *Valor* EZ Herbicide 3 with glyphosate will increase control of emerged weeds and aid in harvest for safflower.

#### TIMING TO SUNFLOWER AND SAFFLOWER

Apply *Valor* EZ Herbicide 3, at 1.5 to 2 fl oz/A (0.047 to 0.063 lb ai), when crop is mature (when seed is 35% moisture or less). For many varieties, this is when the backs of the heads are turning yellow and the bracts are turning brown. Sunflower and safflower can be harvested 5 days after application.

To ensure thorough coverage, use 15 to 30 gallons of spray solution per acre and select nozzle type using manufacturer's gallonage and pressure guidelines for postemergence application.

## DIRECTIONS FOR USE IN SWEET POTATO [For Use in Arizona, California and Hawaii Only]

#### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** apply more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 3 fl oz (0.094 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply postemergence to sweet potatoes.
- **DO NOT** use greenhouse grown transplants.
- **DO NOT** use transplants harvested more than 2 days prior to transplanting.
- **DO NOT** use on any sweet potato variety other than "BEAUREGARD", unless user has tested *Valor* EZ Herbicide 3 on other variety and has found crop tolerance to be acceptable.
- **DO NOT** apply as a part of any tank mix, except with labeled rates of Command, if tank mix is applied prior to transplanting.

#### TIMING TO SWEET POTATOES

*Valor* EZ Herbicide 3 must be applied prior to transplanting sweet potatoes.

#### TIMING TO WEEDS

##### Preemergence To Weeds

Apply *Valor* EZ Herbicide 3 to soil prior to transplanting sweet potato slips for the preemergence control of the weeds listed in Table 1 *Broadleaf Weeds Controlled by Residual Activity of Valor EZ Herbicide 3*.

## DIRECTIONS FOR USE IN WHEAT

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 2 fl oz (0.063 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** apply more than 1 application of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 2 fl oz (0.063 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.

### PRE-PLANT APPLICATIONS, PRE-EMERGENCE WEED CONTROL

[For Use in Delaware, Idaho, Kentucky, Maryland, Minnesota, Montana, North Carolina, North Dakota, New Jersey, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Virginia, Washington and Wisconsin Only]

### RESTRICTIONS AND LIMITATIONS

- For pre-plant weed control, use only on no-till or minimum tillage fields where the previous year's crop residue has not been incorporated into the soil.
- [Plant wheat no sooner than 7 days after *Valor* EZ Herbicide 3 application in the states of DE, ID, KY, MD, MN, MT, NC, ND, NJ, OR, PA, SC, SD, TN, VA, WA or WI]
- [Plant wheat no sooner than 14 days after *Valor* EZ Herbicide 3 application in the states of DE, ID, KY, MD, MN, MT, NC, ND, NJ, OR, PA, SC, SD, TN, VA, WA or WI]
- [DO NOT use on Durum wheat.]
- **DO NOT** irrigate between emergence and spike.
- Wheat must be planted a minimum of 1" deep.
- **DO NOT** graze until wheat has reached 5 inches in height.

### Burndown Use Directions

#### [All states]

*Valor* EZ Herbicide 3, applied as part of a burndown program, at 2 fl oz/A (0.063 lb ai), may be used for residual weed control, as well as to assist in postemergence burndown of many weeds where wheat will be planted directly into the residue of the previous crop. See Directions for Use in Fall Burndown Programs in Fields to be Planted to Barley, Field Pea, Flax, Lentil, Safflower, Sunflower and Spring Wheat for rates and timing of applications. For control of emerged weeds, *Valor* EZ Herbicide 3 must be applied with an appropriate burndown tank mix partner. To ensure thorough coverage, use a minimum of 15 gallons of spray solution per acre. Refer to tank mix partner's label for application pressure and adjuvant systems.

### POST-PLANT, PRE-EMERGENCE WEED CONTROL

[For Use in Delaware, Idaho, Kentucky, Maryland, Minnesota, Montana, North Carolina, North Dakota, New Jersey, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Virginia, Washington and Wisconsin Only]

### RESTRICTIONS AND LIMITATIONS

- For post-plant, pre-emergence weed control, use only on no-till or minimum tillage fields where the previous crop residue has not been incorporated into the soil.
- Apply *Valor* EZ Herbicide 3 up to 2 days after planting.
- [DO NOT use on Durum wheat.]
- **DO NOT** irrigate between emergence and spike.
- Wheat must be planted a minimum of 1" deep.
- **DO NOT** graze until wheat has reached 5 inches in height.

### Use Directions

*Valor* EZ Herbicide 3, applied at 2 fl oz/A (0.063 lb ai), may be used for residual weed control, where wheat has been planted directly into the residue of the previous year. Application must be made no later than 2 days after planting.

## **HARVEST AID**

**[All states]**

### **RESTRICTIONS AND LIMITATIONS**

- **DO NOT** harvest within 10 days of application.

### **Use Directions**

*Valor* EZ Herbicide 3, applied at 2 fl oz/A (0.063 lb ai) for desiccation requires the addition of an agronomically approved adjuvant to the spray mixture. Use a methylated seed oil which contains at least 15% emulsifiers and 80% oil at 1 qt/A. A spray grade nitrogen source (either ammonium sulfate at 2 to 2.5 lb/A or a 28 to 32% nitrogen solution at 1 to 2 qt/A) may be added to the spray mixture along with methylated seed oil to enhance desiccation. The addition of a nitrogen source does not replace the need for methylated seed oil. Tank mixing *Valor* EZ Herbicide 3 with glyphosate will increase control of emerged weeds and aid in harvest.

To ensure thorough coverage, use a minimum of 10 gallons spray solution per acre by ground application and a minimum of 5 gallons per acre by aerial application. Select nozzle based on manufacturer's gallonage and pressure guidelines for postemergence application.

### **TIMING TO WHEAT**

Apply *Valor* EZ Herbicide 3, at 1.5 to 2 fl oz/A (0.047 to 0.063 lb ai), after wheat reaches the hard dough stage and grain has no more than 30% moisture. Wheat can be harvested 10 days after application. Valent recommends tank mixing with glyphosate.

## **DIRECTIONS FOR USE IN BUSHBERRIES, CANEBERRIES, CITRUS FRUIT, GRAPE, OLIVE, POME FRUIT, POMEGRANATE, STONE FRUIT, TREE NUTS AND NON-BEARING FRUIT TREES**

- Bushberries (Subgroup 13-07B): Aronia Berry; Blueberry, Highbush; Blueberry, Lowbush; Buffalo Currant; Chilean Guava; Cranberry, Highbush; Currant, Black; Currant, Red; Elderberry; European Barberry; Gooseberry; Honeysuckle, edible; Huckleberry; Jostaberry; Juneberry (Saskatoon Berry); Lingonberry; Native Currant; Salal; Sea Buckthorn; cultivars, varieties, and/or hybrids of these.
- Caneberries (Subgroup 13-07A): Blackberry; Loganberry; Raspberry, Black; Raspberry, Red; Raspberry, Wild; cultivars, varieties and/or hybrids of these.
- Citrus Fruit (Crop Group 10-10): Australian Desert Lime; Australian Finger-lime; Australian Round Lime; Brown River Finger Lime; Calamondin; Citron; Citrus hybrids; Grapefruit; Japanese Summer Grapefruit; Kumquat; Lemon; Lime; Mediterranean Mandarin; Mount White Lime; New Guinea Wild Lime; Orange, Sour; Orange, Sweet; Pummelo; Russell River Lime; Satsuma Mandarin; Sweet Lime; Tachibana Orange; Tahiti Lime; Tangelo; Tangerine (mandarin); Tangor; Trifoliate Orange; Uniq Fruit; cultivars, varieties and/or hybrids of these.
- Tree Nuts (Crop Group 14-12): African Nut-tree; Almond, Beechnut; Brazil Nut; Brazilian Pine; Bunya; Bur Oak; Butternut; Cajou Nut; Candlenut; Cashew; Chestnut; Chinquapin; Coconut; Coquito Nut; Dika Nut; Ginkgo; Guiana Chestnut; Hazelnut (Filbert); Heartnut; Hickory Nut; Japanese Horse-chestnut; Macadamia Nut; Mongongo Nut; Monkey-pot; Monkey Puzzle Nut; Okari Nut; Pachira Nut; Peach Palm Nut; Pecan; Pequi; Pili Nut; Pine Nut; Pistachio; Sapucaia Nut; Tropical Almond; Walnut, Black; Walnut, English; Yellowhorn, cultivars, varieties and/or hybrids of these.
- Pome Fruit (Crop Group 11-10): Apple; Azarole; Crabapple; Loquat; Mayhaw; Medlar; Pear; Pear, Asian; Quince; Quince, Chinese; Quince, Japanese; Tejocote; cultivars, varieties and/or hybrids of these.
- Stone Fruit (Crop Group 12-12): Apricot; Apricot, Japanese; Capulin; Cherry, Black; Cherry, Nanking; Cherry, Sweet; Cherry, Tart; Jujube, Chinese; Nectarine; Peach; Plum; Plum, American; Plum, Beach; Plum, Canada; Plum, Cherry; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plum, Klamath; Plum, Prune; Plumcot; Sloe and cultivars, varieties and/or hybrids of these.

## RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 12 fl oz (0.375 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application, except Caneberries, **DO NOT** apply more than 6 fl oz (0.188 lb flumioxazin) *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** apply more than 24 fl oz (0.750 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year, except Bushberries; for Bushberries, **DO NOT** apply more than 12 fl oz (0.375 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** make a sequential application within 30 days of the first application, except tree nuts, **DO NOT** make a sequential application within 60 days of the first application.
- **DO NOT** apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation.
- **DO NOT** apply within 300 feet of non-dormant pome or stone fruit, or 300 yards of non-dormant pears.
- Raise mower height during all mowing to reduce dust. Dust created by mowing can drift onto desirable vegetation resulting in injury.
- **DO NOT** apply to powdery soils or soils that are susceptible to wind displacement unless irrigation can be applied immediately after application.
- **DO NOT** mow treated areas between bud break and final harvest. Dust created by mowing may drift onto desirable vegetation resulting in injury.
- Follow the most restrictive label limitations and precautions of the tank mix product(s) being used.
- Avoid direct or indirect spray contact to foliage and green bark (non-barked trunk and non-barked vines with the exception of undesirable suckers).
- **DO NOT** apply to tree nuts established less than one year, unless protected from spray contact by non-porous wraps, grow tubes, or waxed containers.
- For non-bearing fruit trees (avocado and fig), **DO NOT** harvest fruit from treated trees within one year of application.
- Preharvest Interval (PHI)
  - Citrus Fruit: 3 days
  - Bushberries: 7 days
  - Caneberries: 7 days
  - Grape: 60 days
  - Tree Nuts: 60 days
  - Olive: 60 days
  - Pome Fruit: 60 days
  - Pomegranate: 60 days
  - Stone Fruit: 60 days

## PRECAUTIONS

- Use a maximum *Valor* EZ Herbicide 3 rate of 6 fl oz/A (0.188 lb ai) per application on any soil that has a sand plus gravel content over 80% if bushes, trees or vines are less than 3 years of age. (Two applications of 6 fl oz/A (0.188 lb ai) in a 12 month period can still be made as long as there have been 60 days between applications).
- Raise mower height during all mowing to reduce dust. Dust created by mowing can drift onto desirable vegetation resulting in injury.
- Follow the most restrictive label limitations and precautions of the tank mix product(s) being used.
- Avoid direct or indirect spray contact to foliage and green bark or canes (non-barked trunk and non-barked vines with the exception of undesirable suckers).
- Irrigate after application with minimum of 1/4 inch of water to activate the herbicide and to reduce wind displacement of soil.

## USE PRECAUTIONS FOR BUSHBERRIES

- If bushberries are established less than 2 years ensure that they are protected from spray contact by non-porous wrap, grow tubes or waxed containers.

## USE PRECAUTIONS FOR GRAPES

- If grapes are established less than 2 years ensure that they are trellised at least 3 ft from the soil surface or are protected from spray contact by non-porous wrap, grow tubes or waxed containers.
- Apply only to grapes that are trellised, staked or are free standing.
- Avoid direct or indirect spray contact to foliage and green bark (non-barked vines, with the exception of undesirable suckers).
- Plant new plantings of “own-rooted varieties”, for example Concord, so that all roots are a minimum 8 inches below the soil surface to be treated. In some situations, this may require hilling soil around newly planted vines so that the settled depth of the hill will be 4 to 5 inches above the vineyard floor.

## Juice, Raisin and Wine Grapes

- If applied during the period after bud break through final harvest, use shielded application equipment and applicator can ensure spray drift will not come in contact with crop fruit or foliage.

## Table Grapes

- Apply *Valor* EZ Herbicide 3 between final harvest up to bud break.

## USE PRECAUTIONS FOR CITRUS FRUIT, OLIVE, POME FRUIT, POMEGRANATE, STONE FRUIT AND TREE NUTS

- For pome fruit and stone fruit, *Valor* EZ Herbicide 3 can only be applied as a uniform band directed at the base of the trunk prior to silver tip in apples and bud break in stone fruit.
- For pome fruit and stone fruit make applications only to berms.
- For olive, pomegranate and tree nuts apply after bud break through final harvest using shielded application equipment if the applicator can ensure the spray drift will not come into contact with non-target vegetation, crop fruit and/or foliage. Shielded application equipment is not required if the following application parameters are followed:
  - Application pressure (at boom) < 30 PSI.
  - Application speed < 5 MPH.
  - Applicator can ensure the spray drift will not come into contact with non-target vegetation, crop fruit and/or foliage.
- If application is made to trees established less than one year, ensure they are protected from spray contact by non-porous wraps, grow tubes, paint or waxed containers.
  - For apples east of the Cascade Mountains in Washington, follow the restrictions above plus:
    - o Apply between final harvest and January 1.
    - o Apply only to apple blocks with an established (2 years or older) permanent cover crop that covers a minimum of 60% of the surface area in the block.
    - o Application must be incorporated with a minimum of one half inch of water within 48 hours after application.
    - o Apply only to orchard berms.

## USE PRECAUTIONS ON ALMOND AND STONE FRUIT IN DEFINED AREAS OF MERCED, SAN JOAQUIN AND STANISLAUS COUNTIES OF CALIFORNIA

The use of *Valor* EZ Herbicide 3 in soils common in parts of Merced, San Joaquin and Stanislaus counties in California is known to have resulted in injury to almonds under drought stress conditions. These soils are characterized by having been cut or filled, high sand content, low clay content and shallow profiles. Growers in the Defined Area must be aware and assume the risk of using *Valor* EZ Herbicide 3 on almond or stone fruit crops. The Defined Area can be seen on the Map or by the description that follows:

- Intersection of Highway 4 and Escalon-Bellota Road at Farmington in San Joaquin County;
- Directly South on Escalon-Bellota to the Santa Fe Avenue and railroad tracks at Escalon
- Southeast on Santa Fe Avenue down to the Merced River;
- East following the Merced River to the Merced/Mariposa County line;
- Northwest following the Merced County line through the intersection of Merced and Stanislaus County line following the Stanislaus/Tuolumne County and Calaveras County line to Highway 4;
- West on Highway 4 back to the Farmington intersection of Escalon – Bellota Road.



## USE PRECAUTIONS FOR NON-BEARING FRUIT TREES

Non-Bearing Avocado and Fig

- **DO NOT** apply more than 12 fl oz (0.375 lb ai) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** apply more than 24 fl oz (0.750 lb ai) of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** harvest fruit from treated trees within one year of application.
- **DO NOT** apply to trees established less than one year, unless protected from spray contact by non-porous wraps, grow tubes or waxed containers.
- **DO NOT** apply during the period after flowering through leaf drop, unless using shielded application
- equipment and the applicator can ensure spray drift will not come in contact with the crop foliage.

## USE DIRECTIONS

For bushberries, caneberries, citrus fruit, grape, olive, pomegranate, tree nuts, and non-bearing fruit trees, *Valor* EZ Herbicide 3 as a uniform broadcast application to the orchard or vineyard floor or as a uniform band directed at the base of the bush, cane, trunk or vine. For stone fruit and pear, *Valor* EZ Herbicide 3 can only be applied as a uniform band directed at the base of the trunk prior to “bud break”. For apple, *Valor* Herbicide can only be applied as a uniform band directed at the base of the trunk prior to “silver tip”. For other pome fruit, check with Valent personnel for application timing. The preferred application timing for *Valor* EZ Herbicide 3 is in the fall to maximize the potential for rainfall to activate and set the herbicide. **DO NOT** apply over the top of crop or allow spray to come in contact with crop as a result of application or drift.

## Preemergence Application

Apply 6 to 12 fl oz (0.188 to 0.375 lb ai) (maximum 6 fl oz/A for caneberries) of *Valor* EZ Herbicide 3 per broadcast acre as a preemergence application. Make preemergence (to weed emergence) applications of *Valor* EZ Herbicide 3 to a weed-free soil surface. Preemergence applications of *Valor* EZ Herbicide 3 must be completed prior to weed emergence. Moisture is necessary to activate *Valor* EZ Herbicide 3 on soil for residual weed control. Dry weather following application of *Valor* EZ Herbicide 3 may reduce effectiveness. However, when adequate moisture is received after dry conditions, *Valor* EZ Herbicide 3 will control susceptible germinating weeds.

### **Postemergence Application**

Apply 6 to 12 fl oz (0.188 to 0.375 lb ai) (maximum 6 fl oz/A for caneberries) of *Valor* EZ Herbicide 3 per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 qt/A crop oil concentrate). The addition of an adjuvant enhances *Valor* EZ Herbicide 3 activity on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of *Valor* EZ Herbicide 3.

Refer to Table 10 for weeds controlled by the residual activity of *Valor* EZ Herbicide 3. *Valor* EZ Herbicide 3 should be tank mixed with a labeled burndown herbicide for control of the emerged weeds listed in Table 13. *Weeds Controlled by Postemergence Activity of Valor EZ Herbicide 3 Tank Mixes*. Refer to tank mix partner's label for additional weed species and increased weed heights claimed. Refer to tank mix partner's label for additional restrictions, including minimum carrier volume and crops in which tank mix partner may be used. Burndown tank mix partners include glyphosate, paraquat, 2,4-D and Rely®. Tank mixes with glyphosate or 2,4-D containing products are not recommended during the period after bloom through final harvest to ensure crop safety from drift.

Residual weed control will be reduced if vegetation prevents the *Valor* EZ Herbicide 3 from reaching the soil surface. If vegetation is heavy, it is recommended to use a burndown herbicide with *Valor* EZ Herbicide 3 and make a sequential *Valor* EZ Herbicide 3 application prior to the emergence of new weeds.

### **Carrier Volume and Spray Pressure**

To ensure thorough coverage in burndown applications, use a minimum of 15 gallons of spray solution per acre. Use higher gallonage if dense vegetation or heavy crop residue is present.

Nozzle selection must meet manufacturer's gallonage and pressure guidelines.

### **Banded Application**

Rates listed in Table 13, *Weeds Controlled by Postemergence Activity of Valor EZ Herbicide 3 Tank Mixes*, refer to a broadcast application covering the entire acre. Refer to the Band Application table in Use Information section to calculate amount needed per acre when making a banded application.

**Table 13. Weeds Controlled by Postemergence Activity of Valor EZ Herbicide 3 Tank mixes**

| Broadleaf Weed Species                   |                                                   |                             |                                           |
|------------------------------------------|---------------------------------------------------|-----------------------------|-------------------------------------------|
| Common Name                              | Scientific Name                                   | Weed Height/Length (Inches) | Application Rates                         |
| Bindweed, Field <sup>(1)</sup>           | <i>Convolvulus arvensis</i>                       | 8                           | 6 to 12 fl oz/A<br>(0.188 to 0.375 lb ai) |
| Carpetweed                               | <i>Mollugo verticillata</i>                       | 4                           |                                           |
| Chickweeds                               |                                                   |                             |                                           |
| Common                                   | <i>Stellaria media</i>                            | 4                           |                                           |
| Mouseear                                 | <i>Cerastium vulgatum</i>                         | 4                           |                                           |
| Cocklebur, Common                        | <i>Xanthium strumarium</i>                        | 4                           |                                           |
| Evening primrose, Cutleaf <sup>(2)</sup> | <i>Oenothera laciniata</i>                        | 12                          |                                           |
| Filaree                                  |                                                   |                             |                                           |
| Broadleaf                                | <i>Erodium botrys</i>                             | 4                           |                                           |
| Redstem                                  | <i>Erodium cicutarium</i>                         | 4                           |                                           |
| Florida Beggarweed                       | <i>Desmodium tortuosum</i>                        | 2                           |                                           |
| Hemp Sesbania                            | <i>Sesbania exaltata</i>                          | 8                           |                                           |
| Jimsonweed                               | <i>Datura stramonium</i>                          | 4                           |                                           |
| Lambsquarters, Common                    | <i>Chenopodium album</i>                          | 4                           |                                           |
| Morningglories                           |                                                   |                             |                                           |
| Entireleaf                               | <i>Ipomoea hederacea</i> var. <i>integriscula</i> | 4                           |                                           |
| Ivyleaf                                  | <i>Ipomoea hederacea</i>                          | 4                           |                                           |
| Pitted                                   | <i>Ipomoea lacunosa</i>                           | 6                           |                                           |
| Red/Scarlet                              | <i>Ipomoea coccinea</i>                           | 4                           |                                           |
| Tall                                     | <i>Ipomoea purpurea</i>                           | 4                           |                                           |
| Mustard, Wild                            | <i>Brassica kaber</i>                             | 6                           |                                           |
| Pigweeds                                 |                                                   |                             |                                           |
| Palmer Amaranth                          | <i>Amaranthus palmeri</i>                         | 6                           |                                           |
| Redroot                                  | <i>Amaranthus retroflexus</i>                     | 6                           |                                           |
| Smooth                                   | <i>Amaranthus hybridus</i>                        | 6                           |                                           |
| Plantain, Broadleaf                      | <i>Plantago major</i>                             | 6                           |                                           |
| Prickly Sida (Teaweed)                   | <i>Sida spinosa</i>                               | 6                           |                                           |
| Purslanes                                |                                                   |                             |                                           |
| Common                                   | <i>Portulaca oleracea</i>                         | 4                           |                                           |
| Rock                                     | <i>Calandrinia</i> spp.                           | 2                           |                                           |
| Ragweeds                                 |                                                   |                             |                                           |
| Common                                   | <i>Ambrosia artemisiifolia</i>                    | 2                           |                                           |
| Giant                                    | <i>Ambrosia trifida</i>                           | 4                           |                                           |
| Rice Flatsedge                           | <i>Cyperus iria</i>                               | 4                           |                                           |
| Sicklepod                                | <i>Senna obtusifolia</i>                          | 4                           |                                           |
| Smartweeds                               |                                                   |                             |                                           |
| Ladysthumb                               | <i>Polygonum persicaria</i>                       | 4                           |                                           |
| Pale                                     | <i>Polygonum lapathifolium</i>                    | 4                           |                                           |
| Pennsylvania                             | <i>Polygonum pensylvanicum</i>                    | 4                           |                                           |
| Spotted Spurge                           | <i>Euphorbia maculata</i>                         | 4                           |                                           |
| Velvetleaf                               | <i>Abutilon theophrasti</i>                       | 4                           |                                           |
| Venice Mallow                            | <i>Hibiscus trionum</i>                           | 4                           |                                           |
| Waterhemp                                |                                                   |                             |                                           |
| Common                                   | <i>Amaranthus rudis</i>                           | 2                           |                                           |
| Tall                                     | <i>Amaranthus tuberculatus</i>                    | 2                           |                                           |

<sup>(1)</sup>Valor EZ Herbicide 3 will only provide control of the above ground portion of bindweed. Repeated applications will be needed to control regrowth.

<sup>(2)</sup>For acceptable control, cutleaf evening primrose must be 12 inches or less and in the rosette stage. Add crop oil concentrate, at 1 pt/A, or non-ionic surfactant at 0.25% v/v, to glyphosate tank mixes for cutleaf evening primrose control, including glyphosate formulations that contain a built-in adjuvant system.

## ADDITIONAL RESIDUAL WEED CONTROL

*Valor* EZ Herbicide 3 maybe tank mixed with oryzalin (Surflan®), simazine or diuron for additional residual weed control. Always read and follow label use directions for all products being used.

## FALLOWBED USE ON TRANSPLANTED MELON, PEPPER AND TOMATO BEDS [For Use in Arizona, California and Hawaii only]

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply more than 4 fl oz (0.125 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per application.
- **DO NOT** make more than 2 applications of *Valor* EZ Herbicide 3 per acre per year.
- **DO NOT** apply more than 8 fl oz (0.250 lb flumioxazin) of *Valor* EZ Herbicide 3 per acre per year.

Many weather-related factors, including high wind or heavy rains or cool conditions at or near crop transplanting, may result in crop injury in fields treated with *Valor* EZ Herbicide 3. On occasion this has resulted in a delay in maturity. Understand and accept these risks before using *Valor* EZ Herbicide.

| Application Rate                                                                                                                                                                                           | Adjuvant                              | GPA               | Transplanting Interval |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------|-------------------|------------------------|
| 4 fl oz/A                                                                                                                                                                                                  | Required by burndown tank mix partner | Ground – 20 to 40 | 2 Months               |
| <b>Application Method:</b> Apply with a burndown herbicide labeled for the control of emerged weeds. <i>Valor</i> EZ Herbicide 3, when used alone, will not provide satisfactory control of emerged weeds. |                                       |                   |                        |

### Use for Preemergence Fallowbed Weed Control Prior To Transplanting


- Always read and follow all label directions when using any pesticide alone or in tank mix combinations.
- The top 4 inches of the bed, from a horizontal and vertical perspective, where the crop will be transplanted, must be removed prior to transplanting.
- Irrigate treated field after application and prior to transplanting with minimum of 1/4 inch of water if rainfall does not occur between application and transplanting.
- Use only healthy transplants. **DO NOT** use on direct seeded crops.
- [On flat beds (tomato only), the soil must be incorporated to a depth of at least 4 inches, twice, prior to transplanting. Failure to incorporate may result in stand reduction and/or crop injury.]
- This use pattern makes no claim for in-season weed control after the beds have been disturbed.



Beds are formed and *Valor* EZ Herbicide<sup>2</sup> is applied with a burndown herbicide.



A minimum of 2 months after *Valor* EZ Herbicide<sup>2</sup> application, the tops of the beds are removed and the soil from the tops of the beds is placed in the area between the beds.



Crops are transplanted into beds.

## DIRECTIONS FOR USE TO MAINTAIN BARE GROUND ON NON-CROP AREAS OF FARMS, ORCHARDS AND VINEYARDS

### RESTRICTIONS AND LIMITATIONS

- **DO NOT** apply to farm alleys or roads where traffic may result in treated dust settling onto crops or other desirable vegetation.
- **DO NOT** apply to ditch banks.

*Valor* EZ Herbicide 3, when used as directed, can be used on farms, orchards and vineyards for non-selective vegetation control to maintain bare ground on non-crop areas that must be kept weed free. Follow all applicable directions as outlined above under "USE INFORMATION".

*Valor* EZ Herbicide 3 offers residual and postemergence control of susceptible broadleaf and grass weeds as well as an additional mode of action to assist in the control of ALS (acetolactate synthase) resistant weeds. *Valor* EZ Herbicide 3 can be tank mixed with the herbicides listed in Table 14. *Tank Mix Combinations to Maintain Bare Ground Non-Crop Areas* for increased residual or postemergence control. The length of residual control is dependent on the rate applied as well as on rainfall and temperature conditions. Length of residual control will decrease as temperature and precipitation increase. *Valor* EZ Herbicide 3 rates of 6 to 12 fl oz/A (0.188 to 0.375 lb ai) are required to provide residual control of the weeds listed in Table 10.

### PREEMERGENCE APPLICATION

Apply 6 to 12 fl oz (0.188 to 0.375 lb ai/A) of *Valor* EZ Herbicide 3 per broadcast acre as a preemergence application. Make preemergence (to weed emergence) applications of *Valor* EZ Herbicide 3 to a weed-free soil surface. Preemergence applications of *Valor* EZ Herbicide 3 must be completed prior to weed emergence. Moisture is necessary to activate *Valor* EZ Herbicide 3 on soil for residual weed control. Dry weather following application of *Valor* EZ Herbicide 3 may reduce effectiveness. However, when adequate moisture is received after dry conditions, *Valor* EZ Herbicide 3 will control susceptible germinating weeds.

### POSTEMERGENCE APPLICATION

Apply 6 to 12 fl oz (0.188 to 0.375 lb ai/A) of *Valor* EZ Herbicide 3 per broadcast acre plus an adjuvant (0.25% v/v non-ionic surfactant or 1 qt/A crop oil concentrate). The addition of an adjuvant enhances *Valor* EZ Herbicide 3 activity on emerged weeds. Thorough spray coverage is necessary to maximize the postemergence activity of *Valor* EZ Herbicide 3. Emerged weeds are controlled postemergence with *Valor* EZ Herbicide 3, however, translocation of *Valor* EZ Herbicide 3 within a weed is limited, and control is affected by spray coverage and by the addition of an adjuvant. The most effective postemergence weed control with *Valor* EZ Herbicide 3 occurs when applied in combination with a surfactant to weeds less than 2 inches in height. Use a tank mix partner in combination with *Valor* EZ Herbicide 3 for the postemergence control of weeds larger than 2 inches. Specified tank mix partners are listed in Table 14. *Tank Mix Combinations to Maintain Bare Ground Non-Crop Areas*.

**IMPORTANT:** Completely read and follow the label of any potential tank mix partner with *Valor* EZ Herbicide 3. When using tank mixtures, use conditions must be in accordance with the most restrictive of the label limitations and precautions on either herbicide label.

**Table 14. Tank Mix Combinations to Maintain Bare Ground on Non-Crop Areas**

|            |       |      |          |
|------------|-------|------|----------|
| glyphosate | 2,4-D | Rely | paraquat |
|------------|-------|------|----------|

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage, disposal or cleaning of equipment.

### PESTICIDE STORAGE

Keep pesticide in original container.

Store in a cool, dry, secure place.

**DO NOT** put formulation or dilute spray solution into food or drink containers.

**DO NOT** contaminate food or foodstuffs.

**DO NOT** store or transport near feed or food.

Not for use or storage in or around the home.

For help with any spill, leak, fire or exposure involving this material, call day or night **(800) 892-0099**.

### PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING

**[Nonrefillable Container: DO NOT** reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Then offer for recycling, if available or puncture and dispose of in a sanitary landfill.]

#### **[Nonrefillable containers larger than 5 gallons]**

**[Nonrefillable container. DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

#### **[Container statement for refillable container]**

**[Refillable Container:** Refilling Container: Refill this container with Flumioxazin only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When empty, return to point of sale.]

### RECYCLING

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 877-952-2272 (toll free) or [www.acrecycle.org](http://www.acrecycle.org).

**RISKS OF USING THIS PRODUCT,  
LIMITED WARRANTY AND DISCLAIMER,  
AND LIMITATION OF LIABILITY**

**IMPORTANT:** Read the entire Label including this section titled Risks of Using this Product, Limited Warranty and Disclaimer, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

**RISKS OF USING THIS PRODUCT**

The buyer and user (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

The Directions for Use of this product must be followed carefully. Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential, or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. To the extent consistent with applicable law, Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

**LIMITED WARRANTY AND DISCLAIMER**

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label **and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

**LIMITATION OF LIABILITY**

**To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

**PROMPT NOTICE OF CLAIM**

To the extent consistent with applicable law allowing such requirements, Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is later, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law, if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

**NO AMENDMENTS**

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Risks of Using This Product, Limited Warranty and Disclaimer, and Limitation of Liability**, which may not be modified by any oral or written agreement.

**TANK MIXES**

**NOTICE:** Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law.

**Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling of any product used applies in tank mixtures, including all crop rotational and other crop restrictions.**

**[NOTES TO REVIEWER:]**

*[If this product is marketed or distributed by a third party, their logo and Warranty statement will be included on the label.]*

**©2025 Valent U.S.A. LLC**

*Cobra, Phoenix, Select Max and Valor* are trademarks and registered trademarks of Valent U.S.A. LLC

All other trademarks are the property of their respective owners.

*Valent Tank Cleaner* is a product of Valent U.S.A. LLC

Manufactured for:

**Valent U.S.A. LLC**

P.O. Box 5075

San Ramon CA 94583

**Made in U.S.A.**

EPA Reg. No. 59639-273

EPA Est.

059639-00273.20250730.ValorEZ3.AMEND.Cotton.Non-dorm.Pears.+Shack.Clean