

U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460 EPA Reg. Number:

Date of Issuance:

59807-15

DEC 1 7 2013

NOTICE OF PESTICIDE:

X Registration
Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Marathon 1% Granular Greenhouse and Nursery Insecticide

Name and Address of Registrant (include ZIP Code):

OHP, Inc.

P.O. Box 51230

Mainland, PA 19451

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data. You must comply with DCI ID# GDCI-129099-951, issued on November 10, 2010. If you have questions about the Generic DCI issued, you may contact Mr. Russell Wasem from the Pesticide Reevaluation Division at Wasem.Russell@epa.gov.
- 2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You have 18 months from the date of registration to provide these data.

Signature of Approving Official:

Date:

Venus Eagle, Product Manager 01

Insecticide-Rodenticide Branch, Registration Division (7505P)

DEC 1 7 2013

Page 2 EPA Reg. No. 59807-15

- 3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 59807-15."
- 4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

• Basic CSF, dated July 22, 2013

If you have any questions, please contact Julie Chao at 703-308-8735 or chao.julie@epa.gov.

Venus Eagle Product Manager 01 Insecticide-Rodenticide Branch Registration Division (7505P)

Enclosure

GROUP 4A INSECTICIDE

Marathon 1% Granular Greenhouse and Nursery Insecticide

OTHER INGREDIENTS:
TOTAL:
EPA Reg. No. 59807-XX

EPA Est. No.

STOP - READ THE LABEL BEFORE USE KEEP OUT OF REACH OF CHILDREN CAUTION

FOR TRANSPORTATION EMERGENCIES ONLY CALL CHEMTREC 1-800-424-9300 FOR HEALTH and SAFETY and PRODUCT USE INFORMATION CALL 1-800-356-4647

FIRST AID				
IF IN EYES:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, i present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. 			
IF SWALLOWED:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. 			
IF ON SKIN OR CLOTHING:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. 			
	toll free the OHP, Inc. Emergency Telephone No. 1-800-356-4647. Have a product container or label with control center or doctor, or going for treatment.			

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Keep children and pets off treated area until dry.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- · Shoes plus socks
- · Protective eye wear

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

DEC 1 7 2013

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under:

EPA. Reg. No: 59807-15

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker
Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or
modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should:

- · Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- · Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops/plants or weeds. Do not apply this product or allow it to drift to blooming crops/plants or weeds if bees are foraging. Do not apply this product, by any application method, to linden, basswood or other Tilia species.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- · Shoes plus socks
- · Protective eye wear c

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available or place in trash or in a sanitary landfill.

Handle and open container in a manner as to prevent spillage. If material is spilled for any reason or cause, carefully contain any spilled material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides above. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away. You may contact the OHP, Inc. Emergency telephone number at 1-800-356-4647 or contact Chemtrec at 1-800-424-9300.

APPLICATION INFORMATION

Marathon 1% Granular Greenhouse and Nursery Insecticide is for systemic insect control on ornamentals, fruit and nut trees, and vegetable plants in greenhouses, nurseries and interior plantscapes. Marathon 1% Granular Greenhouse and Nursery Insecticide is a systemic product and will be translocated upward within the plant. To assure optimum effectiveness, the product must be placed by incorporation where the roots of the plant can absorb the active ingredient. Irrigate moderately but thoroughly after application, allowing no leaching and runout from containers for at least three irrigations or 10 days whichever is longer.

Resistance: Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area.

For resistance management purposes, a foliar application of any chloronicotinyl insecticide following a Marathon 1% Granular Greenhouse and Nursery Insecticide soil application in the same crop is not recommended.

Incorporation: Incorporation of Marathon 1% Granular Greenhouse and Nursery Insecticide can be achieved by cultivation, irrigation, rainfall, mechanical placement or by using mechanical soil or media mixing equipment.

Woody Perennials: Onset of protection is slower than in herbaceous species. A delay of 2 or more weeks should be expected. Longer delays may be expected with larger plants. Application should therefore be made in advance of expected insect activity.

Bark Media: Media with 30% or more bark content when treated with Marathon 1% Granular Greenhouse and Nursery Insecticide may confer a shorter period of protection.

APPLICATION TO GRASSY AREAS IN NURSERIES

Marathon 1% Granular Greenhouse and Nursery Insecticide can be used for the control of soil inhabiting pests of grassy areas of nurseries, such as northern and southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizotrogus majalis*; green June beetle, *Cotinis nitida*; May or June beetle, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; *Euetheola humilus rugiceps*; billbugs, *Sphenophorus* spp.; annual bluegrass weevil, *Listronotus maculicollis*; black turfgrass ataenius, *Ataenius spretulus*; *Aphodius* spp.; and mole crickets, *Scapteriscus* spp. Marathon 1% Granular Greenhouse and Nursery Insecticide can also be used for suppression of cutworms and chinchbugs. Marathon 1% Granular Greenhouse and Nursery Insecticide can be used as directed on nursery grass in sites such as under or around field or container grown plants, on roadways or other grassy areas in or around nurseries.

The active ingredient in Marathon 1% Granular Greenhouse and Nursery Insecticide has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made proceeding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Do not make applications when grassy areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.

Restrictions:

- Do not use on commercial sod farms.
- Do not apply more than a total of 40 lbs of product (0.4 lbs of active ingredient) per acre per year.

For systemic insect control on fruit and nut trees, and vegetable plants, greenhouse, interior plantscapes, and nursery ornamentals grown in flats, benches, beds and containers. DOSAGE **BULK APPLICATION** TOPICAL APPLICATION CONTAINER **USE SITE PEST** BULK TOPICAL **APPLICATION DIRECTIONS** SIZE POTS/CU YD RATE RATE (INCHES) OF MIX LBS/CU LEVEL TEASP GRAMS/POT OON YD 2.0 to 3.0 4500 to 3200 1/8 to 1/4 Herbaceous 7.0 to 3.5 0.5 to 1.0 Bulk Rate Application: Mix the Adelgids required amount of Marathon 1600 to 1300 Species-Aphids 3.5 to 5.0 3.9 to 3.0 1/4 to 1/3 1.0 to 1.4 1% Granular Greenhouse and including 5.5 to 7.0 900 to 400 Armored scale 1.8 to 1.2 1/3 to 1/2 1.4 to 2.0 Nursery Insecticide per cubic vegetable (suppression) 8.0 to 10.0 200 to 160 1.0 to 0.8 1/2 to 2/3 2.0 to 2.7 yard of medium. Mix plants thoroughly before filling pots Fungus gnats 11.0 to 12.0 80 to 60 0.8 to 0.5 2/3 to 1.0 2.7 to 4.1 (one or two (larvae only)1 and transplanting. plants per pot) Japanese Topical Application: May be Herbaceous 2.0 to 3.0 4,500 to 3,200 9.0 to 7.0 1/4 of 1/3 1.0 to 1.3 made at transplanting, beetles Species (adults) 3.5 to 5.0 1,600 to 1,300 1.3 to 2.0 repotting, or on established 6.0 to 4.0 1/3 to 1/2 including containerized plants. Irrigate Lacebugs 5.5 to 7.0 900 to 400 2.5 to 2.0 1/2 to 2/3 20 to 27 moderately after application to vegetable Leaf beetles plants5 (three 8.0 to 10.0 200 to 160 1.5 to 1.0 2/3 to 1.0 2.7 to 4.1 move the active ingredient into (including Elm or more plants the root zone. Do not allow 11.0 to 12.0 80 to 60 10to08 to 1-1/2 4.1 to 6.1 and Viburnam per pot) significant leaching and runout leaf beetles) for at least 3 irrigations or 10 Woody Leafhoppers Perennial days, whichever is longer. (including Species glassy-winged 5 to 10 1 Gallon 260 to 230 Containerized 5.0 to 2.5 1-1/4 to 2-1/2 For optimum control, make sharpshooter) greenhouse applications prior to egg hatch 2 Gallon 125 to 100 4.4 to 2.2 2-1/2 to 5.0 10 to 20 Leafminers of the target pest. Irrigate and nursery 70 to 60 3-2/3 to 7-1/3 3 Gallon 3.3 to 1.7 15 to 30 Root Mealybugs² ornamentals, moderately after application to 5 Gallon 50 to 30 18.5 to 37 vegetables5 2.0 to 1.0 4-1/2 to 9.0 move the active ingredient into Psyllids and the root zone Root Weevil interiorscapes Complex plants (such as Black Fruit and nut 1 Gallon 260 to 230 5.0 to 2.5 1-1/4 to 2-1/2 5 to 10 For optimum control, make Vine Weevil. applications prior to egg hatch Apopka 2 Gallon 125 to 100 4.4 to 2.2 2-1/2 to 5.0 10 to 20 trees Weevil, Citrus of the target pest. Irrigate 3 Gallon 70 to 60 3.3 to 1.7 3-2/3 to 7-1/3 15 to 30 moderately after application to Root Weevil3) 5 Gallon 50 to 30 2.0 to 1.0 4-1/2 to 9.0 18.5 to 37 move the active ingredient into Soft scale the root zone. Thrips (suppression)4 Apply as a broadcast treatment Ornamental containerized Apply 15 ounces per 1,000 sq ft and vegetable plants grown and incorporate into the soil or Whiteflies in flats, on plants⁵ media before planting. On White Grub benches, or established plants, irrigate larvae in beds moderately after application to (such as move the active ingredient into Japanese the root zone. Minimize beetle. leaching and runout for at least Chafers, 3 irrigations or 10 days, European whichever is longer. chafer, 1 level tsp of Marathon 1% Granular Greenhouse and Nursery Insecticide = 4.1 grams 3 teaspoons = 1 tablespoon Oriental beetle, Asiatic Fungus gnat larvae in the soil will be controlled by incorporation. No adult Fungus Gnat control. Other foliar insect control is garden beetle) achieved by the uptake of Merit 1 G from a healthy root system translocating the active ingredient up into the plant. ² Root Mealybug control will require a thorough incorporation of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: per bulk application. Citrus Root Weevil: For use on non-bearing citrus nursery stock. Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed. Note: for use on vegetable plants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.

For use only on nursery ornamentals.				
PEST	USE SITE	DOSAGE	APPLICATION DIRECTIONS	
White grub larvae (such as Japanese beetle, Masked chafers, European chafer, Oriental beetle, Asiatic garden beetle)	Field and Forest Nurseries	Apply as a uniform band on either side of row using a band six (6) inches wider than the actual root ball diameter to be dug. Do not allow bands in adjacent rows to overlap. Use 45 oz. per 1,000 ft of row or 3,000 sq ft. For grub control in areas of turf, apply as a broadcast application using 45 oz per 3,000 sq. ft.	Vegetation in the area to be treated must be mowed to a height of 3 inches or less prior to application. Mowing to the lowest possible height will insure greater consistency of control. Apply May through July. For optimum control, treatment must be followed by rainfall or irrigation.	

RESTRICTIONS

Do not graze treated areas or use clippings from treated areas for feed or forage.

Do not apply to soils which are water-logged, or saturated. This condition will not allow the penetration of water into the root zone of the plants.

Do not over-irrigate or allow excessive runoff to occur following an application of Marathon 1% Granular Greenhouse and Nursery Insecticide.

Do not apply this product, by any application method, to linden, basswood or other Tilia species.

Do not exceed a total of 40 lbs of product (0.4 lbs of active ingredient) per acre per year.

Food Crops: Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12 month plant-back interval must be observed.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond OHP, Inc. control it is impossible for OHP, Inc. to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, OHP, INC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of OHP, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, OHP, INC. DISCLAIMS ANY liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT OHP, INC.'S ELECTION, THE REPLACEMENT OF PRODUCT.

Marathon® is a registered trademark of OHP, Inc.

OHP, Inc. PO Box 51230 Mainland, PA 19451 (800) 356-4647