



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 05, 2026

SENT BY EMAIL

Jordan Moseley
JordanM@amvac.com
OHP, INC

Subject: Labeling Notification per Pesticide Registration Notice (PRN) 98-10 - Minor Label Revisions
Product Name: MARATHON 1% GRANULAR GREENHOUSE AND NURSERY
INSECTICIDE
Admin Number: 59807-15
EPA Receipt Date: 10/08/2025
Action Case Number: 00672376

Dear Jordan Moseley:

The U.S. Environmental Protection Agency is in receipt of your application for notification under Pesticide Registration Notice 98-10 for the above referenced product. The EPA has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The labeling submitted with this application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains statements or claims substantially differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

If you have questions, please contact Scott Campbell via email at campbell.scott@epa.gov.

Sincerely,

Scott Campbell, Biologist
IVB3, RD
Office of Pesticide Programs

Marathon 1% Granular Greenhouse and Nursery Insecticide

[Alternate Brand Name: Marathon 1% G Insecticide]

For systemic insect control on ornamentals, non-bearing fruit and nut trees, and vegetable transplants in greenhouses, nurseries, and interior landscapes.

ACTIVE INGREDIENT:

Imidacloprid1.0%

OTHER INGREDIENTS:99.0%

TOTAL: 100.0%

EPA Reg. No. 59807-15

EPA Est. No. _____

NOTIFICATION

59807-15

STOP - READ THE LABEL BEFORE USE KEEP OUT OF REACH OF CHILDREN

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

01/05/2026

CAUTION

FOR TRANSPORTATION EMERGENCIES ONLY CALL CHEMTREC 1-800-424-9300

FOR MEDICAL AND PRODUCT USE INFORMATION CALL 1-800-356-4647

FIRST AID

| | |
|--------------------------------|--|
| IF IN EYES: | <ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| IF SWALLOWED: | <ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice. |

In case of emergency call toll free the OHP, Inc. Emergency Telephone No. 1-800-356-4647.

Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) Call 1-800-356-4647

For Chemical Emergency, Spill, Leak, Fire or Accident, Call CHEMTREC® 1-800-424-9300

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.

See [inside][side][back][booklet][leaflet] for [complete] [additional] [First Aid,] [Precautionary Statements] [,Directions for Use] [and][,Storage and Disposal].

EPA Reg. No. 59807-15

EPA Est. No. _____

NET CONTENTS: _____



An American Vanguard Company

OHP, Inc.
5151 McCrimmon Parkway, Suite 275
Morrisville, NC 27560
PO Box 51230
Mainland, PA 19451
(800) 356-4647

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Keep children and pets off treated area until dry.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or ~~Viton~~ **Viton**®.
- Shoes plus socks
- Protective eye wear

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

- When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops/plants or weeds. Do not apply this product or allow it to drift to blooming crops/plants or weeds if bees are foraging.

Do not apply this product, by any application method, to linden, basswood or other Tilia species.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It

contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or Viton.
- Shoes plus socks
- Protective eye wear

APPLICATION INFORMATION

Marathon 1% Granular Greenhouse and Nursery Insecticide is for systemic insect control on ornamentals, non-bearing fruit and nut trees, and vegetable transplants in greenhouses, nurseries and interior plantscapes. **Marathon 1% Granular Greenhouse and Nursery Insecticide** is a systemic product and will be translocated upward within the plant. To assure optimum effectiveness, the product must be placed by incorporation where the roots of the plant can absorb the active ingredient. Irrigate moderately but thoroughly after application, allowing no leaching and runoff from containers for at least three irrigations or 10 days whichever is longer.

RESISTANCE MANAGEMENT RECOMMENDATIONS

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, conform the use of this product to resistance management strategies established for the use area. Consult your Cooperative Extension Service for resistance management strategies and recommended pest management practices for your area.

For resistance management purposes, a foliar application of any chloronicotinyl insecticide following a **Marathon 1% Granular Greenhouse and Nursery Insecticide** soil application in the same crop is not recommended.

Incorporation: Incorporation of **Marathon 1% Granular Greenhouse and Nursery Insecticide** can be achieved by cultivation, irrigation, rainfall, mechanical placement or by using mechanical soil or media mixing equipment.

Woody Perennials: Onset of protection is slower than in herbaceous species. A delay of 2 or more weeks is expected. Longer delays may be expected with larger plants. Make application in advance of expected insect activity.

Bark Media: Media with 30% or more bark content when treated with **Marathon 1% Granular Greenhouse and Nursery Insecticide** may confer a shorter period of protection.

APPLICATION TO GRASSY AREAS IN NURSERIES

Marathon 1% Granular Greenhouse and Nursery Insecticide can be used for the control of soil inhabiting pests of grassy areas of nurseries, such as northern and southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizotrogus majalis*; green June beetle, *Cotinis nitida*; May or June beetle, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; *Euethola humilis rugiceps*; billbugs, *Sphenophorus* spp.; annual bluegrass weevil, *Listronotus maculicollis*; black turfgrass atenius, *Ataenius spretulus*; *Aphodius* spp.; and mole crickets, *Scapteriscus* spp. **Marathon 1% Granular Greenhouse and Nursery Insecticide** can also be used for suppression of cutworms and chinchbugs. **Marathon 1% Granular Greenhouse and Nursery Insecticide** can be used as directed on nursery grass in sites such as under or around field or container grown plants, on roadways or other grassy areas in or around nurseries.

The active ingredient in **Marathon 1% Granular Greenhouse and Nursery Insecticide** has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Do not make applications when grassy areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.

RESTRICTIONS:

- **Do not** use on commercial sod farms.
- **Do not** apply more than a total of 40 lbs of product (0.4 lbs of active ingredient) per acre per year.

DIRECTIONS FOR APPLICATIONS TO ORNAMENTALS AND VEGETABLE TRANSPLANTS

For systemic insect control on ornamentals, non-bearing fruit and nut trees, and vegetable transplants in greenhouses, nurseries and interior plantscapes. Ornamentals grown in flats, benches, beds and containers.

Restriction For Outdoor Use: Do not apply this product, by any application method, to linden, basswood, or other Tilia species.

| PEST | USE PATTERN | DOSAGE | | | | | APPLICATION DIRECTIONS |
|---|--------------------------------|-------------------------|---------------------------------|---------------------|-----------------------------|------------|---|
| | | CONTAINER SIZE (INCHES) | BULK APPLICATION | | TOPICAL APPLICATION | | |
| | | | POTS/CU YD OF MIX | BULK RATE LBS/CU YD | TOPICAL RATE LEVEL TEASPOON | GRAMS/POT | |
| Adelgids | Herbaceous | 2.0 to 3.0 | 4,500 to 3,200 | 7.0 to 3.5 | 1/8 to 1/4 | 0.5 to 1.0 | Bulk Rate Application: Mix the required amount of Marathon 1% Granular Greenhouse and Nursery Insecticide per cubic yard of medium. Mix thoroughly before filling pots and transplanting. Topical Application: May be made at transplanting, repotting, or on established containerized plants. Irrigate moderately after application to move the active ingredient into the root zone. Do not allow significant leaching and runoff for at least 3 irrigations or 10 days, whichever is longer. |
| Aphids | Species- including | 3.5 to 5.0 | 1,600 to 1,300 | 3.9 to 3.0 | 1/4 to 1/3 | 1.0 to 1.4 | |
| Armored scale | vegetable | 5.5 to 7.0 | 900 to 400 | 1.8 to 1.2 | 1/3 to 1/2 | 1.4 to 2.0 | |
| (suppression) | transplants ⁵ (one | 8.0 to 10.0 | 200 to 160 | 1.0 to 0.8 | 1/2 to 2/3 | 2.0 to 2.7 | |
| Fungus gnats | or two plants per | 11.0 to 12.0 | 80 to 60 | 0.8 to 0.5 | 2/3 to 1.0 | 2.7 to 4.1 | |
| (larvae only) ¹ | pot) | | | | | | |
| Japanese beetles | Herbaceous | 2.0 to 3.0 | 4,500 to 3,200 | 9.0 to 7.0 | 1/4 of 1/3 | 1.0 to 1.3 | |
| (adults) | Species including | 3.5 to 5.0 | 1,600 to 1,300 | 6.0 to 4.0 | 1/3 to 1/2 | 1.3 to 2.0 | |
| Lacebugs | vegetable | 5.5 to 7.0 | 900 to 400 | 2.5 to 2.0 | 1/2 to 2/3 | 2.0 to 2.7 | |
| Leaf beetles | transplants ⁵ | 8.0 to 10.0 | 200 to 160 | 1.5 to 1.0 | 2/3 to 1.0 | 2.7 to 4.1 | |
| (including Elm and | (three or more | 11.0 to 12.0 | 80 to 60 | 1.0 to 0.8 | 1.0 to 1-1/2 | 4.1 to 6.1 | |
| Viburnum leaf | plants per pot) | | | | | | |
| beetles) | Woody Perennial | | | | | | |
| Species | | | | | | | |
| Leafhoppers | | | | | | | |
| (including glassy-winged | | | | | | | |
| sharpshooter) | | | | | | | |
| Leafminers | | | | | | | |
| Root Mealybugs ² | | | | | | | |
| Psyllids | Containerized | | 260 to 230 | 5.0 to 2.5 | 1-1/4 to 2-1/2 | 5 to 10 | For optimum control, make applications prior to egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone. |
| Root Weevil Complex | greenhouse and | | 125 to 100 | 4.4 to 2.2 | 2-1/2 to 5.0 | 10 to 20 | |
| (such as Black Vine | nursery | 1 Gallon | 70 to 60 | 3.3 to 1.7 | 3-2/3 to 7-1/3 | 15 to 30 | |
| Weevil, Apopka | ornamentals, | 2 Gallon | 50 to 30 | 2.0 to 1.0 | 4-1/2 to 9.0 | 18.5 to 37 | |
| Weevil, Citrus Root | vegetable | 3 Gallon | | | | | |
| Weevil ³) | transplants ⁵ , and | 5 Gallon | | | | | |
| Soft scale Thrips | interiorscapes | | | | | | |
| (suppression) ⁴ | plants | | | | | | |
| Whiteflies | Non-bearing fruit | 1 Gallon | 260 to 230 | 5.0 to 2.5 | 1-1/4 to 2-1/2 | 5 to 10 | For optimum control, make applications prior to egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone. |
| | and nut trees ⁶ | 2 Gallon | 125 to 100 | 4.4 to 2.2 | 2-1/2 to 5.0 | 10 to 20 | |
| | | 3 Gallon | 70 to 60 | 3.3 to 1.7 | 3-2/3 to 7-1/3 | 15 to 30 | |
| | | 5 Gallon | 50 to 30 | 2.0 to 1.0 | 4-1/2 to 9.0 | 18.5 to 37 | |
| White Grub larvae | | | | | | | |
| (such as Japanese | | | | | | | |
| beetle, Chafer, | | | | | | | |
| European chafer, | | | | | | | |
| Oriental beetle, | | | | | | | |
| Asiatic garden | | | | | | | |
| beetle) | Ornamental and | Containerized | Apply 15 ounces per 1,000 sq ft | | | | Apply as a broadcast |
| | vegetable | plants grown in | | | | | treatment and incorporate |
| | transplants ⁵ | flats, on | | | | | into the soil or media |
| | | benches, or in | | | | | before planting. On |
| | | beds | | | | | established plants, irrigate |
| | | | | | | | moderately after |
| | | | | | | | application to move the |
| | | | | | | | active ingredient into the |
| | | | | | | | root zone. Minimize |
| | | | | | | | leaching and runoff for at |
| | | | | | | | least 3 irrigations or 10 |
| | | | | | | | days, whichever is longer. |
| 1 level tsp of Marathon 1% Granular Greenhouse and Nursery Insecticide = 4.1 grams 3 teaspoons =1 tablespoon | | | | | | | |
| ¹ Fungus gnat larvae in the soil will be controlled by incorporation. No adult Fungus Gnat control. Other foliar insect control is achieved by the uptake of Marathon 1% Granular Greenhouse and Nursery Insecticide from a healthy root system translocating the active ingredient up into the plant. | | | | | | | |
| ² Root Mealybug control will require a thorough incorporation of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: per bulk application. | | | | | | | |
| ³ Citrus Root Weevil: For use on non-bearing citrus nursery stock. | | | | | | | |
| ⁴ Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed. | | | | | | | |
| ⁵ Note: for use on vegetable transplants intended for resale only including: Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato. | | | | | | | |
| ⁶ Non-bearing fruit and nut trees are those trees that will not bear fruits or nuts for one year after application | | | | | | | |

FOR USE ONLY ON NURSERY ORNAMENTALS.

| PEST | USE | DOSAGE | APPLICATION DIRECTIONS |
|--|----------------------------|--|---|
| White grub larvae (such as Japanese beetle, Masked chafers, European chafer, Oriental beetle, Asiatic garden beetle) | Field and Forest Nurseries | Apply as a uniform band on either side of row using a band six (6) inches wider than the actual root ball diameter to be dug. Do not allow bands in adjacent rows to overlap. Use 45 oz. per 1,000 ft of row or 3,000 sq ft. For grub control in areas of turf, apply as a broadcast application using 45 oz per 3,000 sq ft. | Vegetation in the area to be treated must be mowed to a height of 3 inches or less prior to application. Mowing to the lowest possible height will insure greater consistency of control. Apply May through July. For optimum control, treatment must be followed by rainfall or irrigation. |

RESTRICTIONS

Do not graze treated areas or use clippings from treated areas for feed or forage.

Do not apply to soils which are water-logged, or saturated. This condition will not allow the penetration of water into the root zone of the plants.

Do not over-irrigate or allow excessive runoff to occur following an application of **Marathon 1% Granular Greenhouse and Nursery Insecticide**.

Do not exceed a total of 40 lbs of product (0.4 lbs of active ingredient) per acre per year.

Restriction for Outdoor Use: Do not apply this product, by any application method, to linden, basswood or other Tilia species.

Food Crops: Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient.

For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a ~~12-month~~12-month plant-back interval must be observed.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: NONREFILLABLE CONTAINER. Do not reuse or refill this container. Offer for recycling, if available or place in trash or in a sanitary landfill. Handle and open container in a manner as to prevent spillage. If material is spilled for any reason or cause, carefully contain any spilled material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides above. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away. You may contact the OHP, Inc. Emergency telephone number at 1-800-356-4647 or contact Chemtrec at 1-800-424-9300.

IMPORTANT: READ BEFORE USE

~~Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.~~

~~If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.~~

~~**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond OHP, Inc. control it is impossible for OHP, Inc. to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.~~

~~**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, OHP, INC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of OHP, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, OHP, INC. DISCLAIMS ANY liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.~~

~~**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT OHP, INC.'S ELECTION, THE REPLACEMENT OF PRODUCT.~~

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; and (b) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS WARRANTY DOES NOT EXTEND TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS, OR UNDER CONDITIONS NOT REASONABLY FORESEEABLE.

THERE ARE NO EXPRESS OR IMPLIED WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN TORT, CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

© 20[YY] All Rights Reserved. OHP, Inc. is a wholly owned subsidiary of American Vanguard Corporation. AMERICAN VANGUARD is a trademark owned by AMVAC Chemical Corporation. OHP, MARATHON and their respective logos are trademarks owned by OHP, Inc. All other trademarks are the property of their respective owners.

~~Marathon® is a registered trademark of OHP, Inc. OHP, Inc.~~

~~PO Box 51230~~

~~Mainland, PA 19451~~

~~(800) 356-4647~~

~~March 16, 2015~~