

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes irreversible eye damage. Causes skin irritation. Harmful if inhaled. Harmful if swallowed. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Causes asthmatic signs and symptoms in hyper-reactive individuals.

- Do not get in eyes, on skin, on clothing.
- Avoid breathing vapor.
- Do not swallow.
- Wear goggles, protective clothing, and rubber gloves.
- Wash thoroughly with soap and water after handling.
- Remove contaminated clothing and wash before reuse.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry, well ventilated climate controlled area.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or your Environmental Control Agency, or the Hazardous Waste representative at nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For single use only
U.S. Patent 6032705
U.S. Patent No. 5,635,196

PremiCide
Sanitizer/Solidifier

THE EFFICACY OF THIS PRODUCT HAS BEEN DEMONSTRATED IN THE PRESENCE OF 100% BLOOD SERUM.

Active Ingredient:

Glutaraldehyde.....	9.61%
Other Ingredients.....	90.39%
Total.....	100%

KEEP OUT OF REACH OF CHILDREN

DANGER

FIRST AID

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call poison control center or doctor for treatment advice.

IF ON SKIN:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call poison control center or doctor for treatment advice.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance. Then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Questions? 1-800-424-9300

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

See side panel for additional precautionary statements

Fast Acting Encapsulant which contains "SCENTED DRY Glutaraldehyde" to Sanitize and Solidify Liquid Medical Waste and Laboratory, Human and Animal Wastes. Antibacterial. Controls odor formation by destroying odor causing bacteria. THE EFFICACY OF THIS PRODUCT HAS BEEN DEMONSTRATED IN THE PRESENCE OF 100% BLOOD SERUM.

Reorder No: PG500

Solidifies up to 500cc

Directions for use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To help prevent the blood from clotting inside the suction canister, immediately dispense PremiCide following the end of surgery or after the suction canister has been completely filled.

- Wear hand, eye and face protection when handling infectious waste and using PremiCide.
- Remove plastic safety seal as per instructions.
- Prior to or following surgery, open the appropriate port and place the PremiCide top on to the open port and push down to secure proper fitting.
- Following surgery turn off suction and remove the patient tubing. Close all ports and activate suction. Firmly grasp the base of the cap and pull upwards on the bottle. Suction will pull the PremiCide powder into the canister.
- Once the bottle is empty, tap the side of the bottle to remove any remaining powder and then push down on the bottle to close the system. Once the liquid waste is solidified, remove the entire bottle by gripping the base of the PremiCide cap and lift up to remove. Replace the port cap on the suction canister and lid, invert Solidified canister and dispose of according to institutional procedures and local State and Federal regulations.

LIMITED WARRANTY AND DISCLAIMER

See enclosed sheet.

OBF TECHNOLOGIES, INC.

2719 Curtiss Street • Downers Grove, IL 60515

(630) 515-1683

EPA Est. No. 072075-WI-001 EPA Reg. No. 59839-1

Net Contents: .72 ounces

ACCEPTED
MAR 21 2001
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 59839-1

59839-1
3/21/2001
page 1 of 1