

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <u>  x  </u> Registration <u>      </u> Reregistration</p> <p>(under FIFRA, as amended)</p>	EPA Reg. Number: 59913-11	Date of Issuance: MAY 12 2004
	Term of Issuance: Conditional	
	Name of Pesticide Product: Safer Brand Garden Fungicide RTUII	
Name and Address of Registrant (include ZIP Code): Safer Inc. c/o Lewis & Harrison 122 C Street, N.W., Suite 740 Washington, DC 20001		
<p><b>Note:</b> Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(C) provided that you:</p> <ol style="list-style-type: none"> <li>1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.</li> <li>2. Make the following label change:  Revise the EPA Registration Number to read, "EPA Reg. No. 59913-11".</li> <li>3. Storage stability and corrosion characteristics data must be submitted upon completion of the studies.</li> </ol>		
Signature of Approving Official: <i>Mary L. Waller</i> Mary L. Waller, Product Manager (21)	Date: MAY 12 2004	

Submit one copy of the final printed label before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Mary L. Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7505C)

Enclosure

7505C:CGrable:cg:5/12/04

EPA Reg. No. 059913-\_\_\_\_\_

Draft Master Label

November 26, 2003

Page 1 of 5

MASTER LABEL- FRONT PANEL (PAGE #1 of 4)

**SAFER® BRAND GARDEN FUNGICIDE RTU II**

 For Organic Production

[Optional Logo]



[Optional Logo]

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

MAY 12 2004

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.

59913-11

Controls  
POWDERY MILDEW, BLACK SPOT, LEAF SPOT and RUST  
On  
Fruits, Vegetables, Flowers and Foliage

[Additional Brand Names]

**Garden Fungicide For Flowers, Fruit, and Vegetables, Safer® Brand**

ACTIVE INGREDIENT:	BY WT.
Sulfur.....	0.40%
OTHER INGREDIENTS:.....	99.60%
TOTAL.....	100.00%

**KEEP OUT OF REACH OF CHILDREN**

**See back panel for additional Precautionary Statements**

NET CONTENTS 24 Fl. Oz., 32 Fl. Oz., 128 Fl. Oz.

**PRECAUTIONARY STATEMENTS**  
Hazards to Humans and Domestic Animals

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**WHAT IT IS:** Safer® Brand Garden Fungicide RTU II is specially formulated with sulfur which inhibits the attack of fungal diseases by creating an environment which is not optimal for disease growth.

**WHAT IT CONTROLS:** Powdery mildew, black spot, leaf spot and rust.

**WHERE TO USE IT:** Roses, Flowers, Foliage Plants, Ornamentals, Beans, Peas, Broccoli, Cabbage, Cauliflower, Collards, Kale, Peppers, Strawberries, Apples, Pears, Potatoes, Cucumbers, Squash, Grapes, Blackberries, Blueberries, Raspberries, Loganberries, Currants, Melons and Tomatoes.

**HOW TO USE IT:** Spray at first sign of infection. Spray plants thoroughly to wet the upper and lower surfaces of foliage.

**ROSES, FLOWERS, FOLIAGE PLANTS & ORNAMENTALS:** Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew, leaf spot, black spot, and rust. Be sure to treat new growth and buds.

**AZALEA, BUTTONBUSH, CHRYSANTHEMUM, HYDRANGEA, MATRIMONY VINE, RHODODENDRON, AND SNAPDRAGON:** Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew, leaf spot, and rust. Be sure to treat new growth and buds.

**CATALPA, DELPHINIUM, HONEYSUCKLE, HORSE CHESTNUT, LILAC, LINDEN, PHLOX, SWEETPEA, SYCAMORE, TRUMPET VINE, WITCH HAZEL, AND ZINNIA:** Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew and leaf spot. Be sure to treat new growth and buds.

**CARNATION AND STAGHORN SUMAC:** Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control leaf spot and rust. Be sure to treat new growth and buds.

**ALDER AND ASTER:** Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew and rust. Be sure to treat new growth and buds.

**DAHLIA:** Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew. Be sure to treat new growth and buds.

**BEANS, PEAS, BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE AND PEPPERS:** Apply at first sign of disease infestation. Spray at 7 day intervals, following rain, or as needed to control powdery mildew. Can be used up to the day before harvest.

**STRAWBERRIES:** Apply at first sign of disease infestation. Repeat at 3 week intervals to control powdery mildew. Do not use on strawberry fruit that will be used for canning. Can be used up to the day before harvest.

**APPLES & PEARS:** Apply at pre-bloom stage or after petal fall as needed to control powdery mildew. NOTE: Some apple & pear varieties are sulfur sensitive; test for varietal sensitivity prior to full scale treatment.

**POTATOES:** Apply before disease infestation appears and repeat as necessary to control powdery mildew.

**CUCUMBERS & SQUASH:** Apply at first sign of disease infestation and repeat as necessary to control powdery mildew. Do not spray young plants.

**GRAPES:** For control of powdery mildew, apply when new shoots are 6-10 inches. Repeat before blooms open and at 14 day intervals as needed. NOTE: Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord & other labrusca varieties.

**BLACKBERRIES, BLUEBERRIES, RASPBERRIES, LOGANBERRIES AND CURRANTS:** Apply each week starting at first bloom color to control powdery mildew. Continue to spray at 10 to 14 day intervals until fruit is set, then repeat as needed or following rain.

**MELONS:** Apply at first sign of disease infestation. Spray at 7 to 10 day intervals following rain, or as needed to control powdery mildew. NOTE: Melon foliage is extremely sensitive to sulfur burn. See paragraph on Use Cautions.

**TOMATOES:** Apply at first sign of disease infestation and repeat as necessary to control powdery mildew. Do not spray young plants.

**USE CAUTIONS:** Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not use on Boston fern, spinach, apricots, filberts, walnuts and viburnum. Do not apply when temperature exceeds 85° F or in full sun. Do not apply for at least 4 weeks following application of oil spray. Avoid contact with plants until spray has dried.

EPA Reg. No. 059913-\_\_\_\_\_

Draft Master Label

November 26, 2003

Page 4 of 5

MASTER LABEL- BACK PANEL (Page #3 of 4)

6/7

### STORAGE AND DISPOSAL

**STORAGE:** Store only in original container in a cool, dry area inaccessible to children and pets. Protect from freezing and heat. Keep container tightly closed to prevent evaporation.

**DISPOSAL:**

If Empty: Do not reuse this container. Place in trash or offer for recycling if available.

If Partly Filled: Call your local solid waste agency or (800)-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Woodstream Corp. and its subsidiaries will not accept liability for damages or injury resulting from misuse. For information on this product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Distributed by

Safer®, Inc., a wholly owned subsidiary of Woodstream Corporation

69 North Locust Street

Lititz PA 17543

1-800-800-1812 for information

EPA Reg. No. 59913-\_\_\_\_\_

EPA Est. No. 44616-MO-001

All ® Property of Woodstream Corporation

© Copyright Woodstream Corporation, 2003

EPA Reg. No. 059913-\_\_\_\_\_

Draft Master Label

November 26, 2003

Page 5 of 5

MASTER LABEL- BACK PANEL (Page #4 of 4)

7/7

[Optional Marketing Claims]

*Effective*

*Water-based*

*Made with elemental sulfur*

*Use as a preventative treatment.*

*Use up to day before harvest.*

*Always shake well before using.*

*Shake well*

*Made in USA*

*Printed in USA*

*Si no usted entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)*

*The Right Solution™. Safer® Brand Garden Fungicide Ready-To-Use is used for prevention and control of fungal diseases, but is gentle on flowers, plants and produce. It treats foliage with the naturally occurring essential element sulfur, to create an environment where damaging fungi cannot live.*

*Its non-burning, easy to use spray formulation can be used on flowers, fruits and vegetables up until the day before harvest.*

