

59913-11

05/02/2005

1/4

EPA Reg. No. 059913-11
Amendment -Master Label
April 28, 2005
MASTER LABEL- FRONT PANEL (PAGE #1 of 1)

SAFER® BRAND GARDEN FUNGICIDE RTU II

For Organic Production [Optional Logo]
For Organic Gardening [Optional Logo]
OMRI™ Listed- Organic Materials Review Institute [Optional Logo]

Controls
POWDERY MILDEW, BLACK SPOT, LEAF SPOT and RUST
On
Fruits, Vegetables, Flowers and Foliage

[Additional Brand Names]


*Garden Fungicide For Flowers, Fruit, and Vegetables, Safer® Brand
Garden Fungicide, For Flowers, Fruits & Vegetables-Safer® Brand
Safer® Brand – Garden Fungicide, For Flowers, Fruits & Vegetables*

| | |
|-------------------------|---------|
| ACTIVE INGREDIENT: | BY WT. |
| Sulfur..... | 0.40% |
| OTHER INGREDIENTS:..... | 99.60% |
| TOTAL..... | 100.00% |

KEEP OUT OF REACH OF CHILDREN

See back panel for additional Precautionary Statements

NET CONTENTS 24 Fl. Oz., 32 Fl. Oz., 128 Fl. Oz.



2/4

EPA Reg. No. 059913-11
Amendment -Master Label
April 28, 2005

CARNATION AND STAGHORN SUMAC: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control leaf spot and rust. Be sure to treat new growth and buds.

ALDER AND ASTER: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew and rust. Be sure to treat new growth and buds.

DAHLIA: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals, following rain, or as needed to control powdery mildew. Be sure to treat new growth and buds.

BEANS, PEAS, BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, KALE AND PEPPERS: Apply at first sign of disease infestation. Spray at 7 day intervals, following rain, or as needed to control powdery mildew. Can be used up to the day before harvest.

STRAWBERRIES: Apply at first sign of disease infestation. Repeat at 3 week intervals to control powdery mildew. Do not use on strawberry fruit that will be used for canning. Can be used up to the day before harvest.

APPLES & PEARS: Apply at pre-bloom stage or after petal fall as needed to control powdery mildew. NOTE: Some apple & pear varieties are sulfur sensitive; test for varietal sensitivity prior to full scale treatment.

POTATOES: Apply before disease infestation appears and repeat as necessary to control powdery mildew.

CUCUMBERS & SQUASH: Apply at first sign of disease infestation and repeat as necessary to control powdery mildew. Do not spray young plants.

GRAPES: For control of powdery mildew, apply when new shoots are 6-10 inches. Repeat before blooms open and at 14 day intervals as needed. NOTE: Do not apply to wine grapes within 21 days of harvest. Sulfur may damage concord & other labrusca varieties.

BLACKBERRIES, BLUEBERRIES, RASPBERRIES, LOGANBERRIES AND CURRANTS: Apply each week starting at first bloom color to control powdery mildew. Continue to spray at 10 to 14 day intervals until fruit is set, then repeat as needed or following rain.

MELONS: Apply at first sign of disease infestation. Spray at 7 to 10 day intervals following rain, or as needed to control powdery mildew. NOTE: Melon foliage is extremely sensitive to sulfur burn. See paragraph on Use Cautions.

TOMATOES: Apply at first sign of disease infestation and repeat as necessary to control powdery mildew. Do not spray young plants.

USE CAUTIONS: Some plants are susceptible to injury from sulfur under certain climatic conditions. Do not use on Boston fern, spinach, apricots, filberts, walnuts and viburnum. Do not apply when temperature exceeds 85° F or in full sun. Do not apply for at least 4 weeks following application of oil spray. Avoid contact with plants until spray has dried.

EPA Reg. No. 059913-11
Amendment -Master Label
April 28, 2005
MASTER LABEL- BACK PANEL (Page #3 of 4)

STORAGE AND DISPOSAL

STORAGE: Store only in original container in a cool, dry area inaccessible to children and pets. Protect from freezing and heat. Keep container tightly closed to prevent evaporation.

DISPOSAL:

If Empty: Do not reuse this container. Place in trash or offer for recycling if available.
If Partly Filled: Call your local solid waste agency or (800)-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Woodstream Corp. and its subsidiaries will not accept liability for damages or injury resulting from misuse. For information on this product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Distributed by
Safer®, Inc., a wholly owned subsidiary of Woodstream Corporation
69 North Locust Street
Lititz PA 17543
1-800-800-1812 for information

EPA Reg. No. 59913-11
EPA Est. No. 44616-MO-001

All ® Property of Woodstream Corporation
© Copyright Woodstream Corporation, 2005

4/4
EPA Reg. No. 059913-11
Amendment -Master Label
April 28, 2005

[Optional Marketing Claims]

*### oz. Ready-To-Use
Effective
Water-based
Made with elemental sulfur
Use as a preventative treatment.
Use up to day before harvest.
Always shake well before using.
Shake well
Made in USA
Printed in USA*

Si no usted entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

The Right Solution™. Safer® Brand Garden Fungicide Ready-To-Use is used for prevention and control of fungal diseases, but is gentle on flowers, plants and produce. It treats foliage with the naturally occurring essential element sulfur, to create an environment where damaging fungi cannot live.

Its non-burning, easy to use spray formulation can be used on flowers, fruits and vegetables up until the day before harvest.