

ENVIRONMENTAL PROTECTION AGENCY

D 368668

John French, Ph.D.
Sipcam Agro, USA Inc.
300 Colonial Center Parkway
Suite 230
Roswell, GA 30076

APR 04 2007

Dear Dr. French:

SUBJECT: Eminent 125SL Fungicide
EPA Registration Number 60063-12
Your Submissions of June 23, 2006 and August 23, 2006 adding the new uses
pecans and peanuts

The amendment referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) is acceptable.

In the Agency's risk assessment for 1,2,4-triazole and its metabolites triazole alanine, and triazole acetic acid, dated February 7, 2006 (see docket number 2005-0497 at <http://www.regulations.gov>), the Agency identified additional data that are needed for triazole-derivative active ingredients which have the 1,2,4,-triazole ring. The Agency plans to issue a Data Call-In for all triazole-derivative active ingredients which will identify all data needs for these active ingredients listed in the February 7, 2006 risk assessment. The Data Call-In will not address data needs that are related to parent-only data requirements.

Please submit one copy of the final printed labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records. If you have any questions, please contact Lisa Jones of my team at (703) 308-9424 or at jones.lisa@epa.gov.

Sincerely,

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505P)

Enclosure: Stamped Label

CONCURRENCES

SYMBOL <	7505C						
SURNAME <	Lisa Jones						
DATE <	Apr 4, 2007						



2 8 7

SIPCAM AGRO USA, INC.

EMINENT® 125SL FUNGICIDE

ACTIVE INGREDIENT:

Tetraconazole {1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2-tetrafluoroethoxy) propyl]-1H-1,2,4-triazole}... 11.6%

OTHER INGREDIENTS: 88.4%

TOTAL: 100.0%

Contains 1 lb. active ingredient (tetraconazole) per gallon.

KEEP OUT OF REACH OF CHILDREN


CAUTION

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> ➤ Call a poison control center or doctor immediately for treatment advice. ➤ Have affected person sip a glass of water if able to swallow. ➤ Do not induce vomiting unless told by a poison control center or doctor. ➤ Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> ➤ Take off contaminated clothing. ➤ Rinse skin immediately with plenty of water for 15-20 minutes. ➤ Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> ➤ Hold eye open and rinse slowly and gently with water for 15-20 minutes. ➤ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ➤ Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 900-4044 Poison Control Center (human health) (800) 345-4735 ASPCA (animal health)

EPA Registration No. 60063 - 12

EPA Establishment No. _____

NET CONTENTS: _____



Manufactured for:
SIPCAM AGRO USA, INC.
300 Colonial Center Parkway, Suite 230
Roswell, GA 30076 USA

EMINENT® 125SL FUNGICIDE

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, and viton ≥14 mils. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product may be toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms adjacent to treatment areas. Exercise care when making applications of Eminent, and do not apply when atmospheric conditions favor drift or runoff. Do not contaminate water when disposing of equipment washwaters or rinsate.

In order to mitigate concern for reproductive effects to endangered bird and mammal species which may occur incidentally in sugarbeet growing areas, you are required to ascertain through the state Department of Agriculture, or Cooperative Extension Service, whether the treatment area may contain habitat of Federally listed bird and mammal species; if so, treatment must be avoided in these areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instruction and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

GENERAL USE INFORMATION:

Apply EMINENT in water carrier by spraying onto crop surfaces that are intended to be protected from disease. To obtain adequate coverage of typical agricultural crops, total spray volume usually will range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute sprays, and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended.

EMINENT should be added to the spray tank while filling with water. Keep the agitator running when filling spray tank and during spray operations. When tank mixing this product with other pesticides observe the more restrictive label imitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Combination in the spray tank with other pesticides, fertilizers or surfactants is not recommended unless prior use has shown the combination to be physically compatible, effective and noninjurious under your conditions of use.

EMINENT contains a sterol demethylation inhibitor (sterol DMI) fungicide. EMINENT should be applied in tank mix or in alternate applications with non-DMI fungicides as a resistance management strategy. Do not mix or alternate EMINENT with other DMI fungicides.

AGRICULTURAL CROPS:

Crop	Diseases Controlled	EMINENT 125SL Rate per Acre	Pre-harvest Interval	Application Directions
Sugarbeets	Cercospora leafspot, Powdery Mildew	13 fluid ounces	14 days	<p>Apply EMINENT at intervals of 21 to 28 days, beginning when conditions are favorable for Cercospora leafspot or powdery mildew. Applications of EMINENT must be alternated with a non-triazole fungicide which is registered for use on sugarbeets. If the alternate fungicide is known to provide protection for up to 14 days, applications of EMINENT should be no longer than 28 days apart. Do not make more than two (2) applications of EMINENT per growing season. Crops other than sugarbeet must not be grown within 120 days following the last application of EMINENT 125SL.</p>
Peanut	Early leafspot, Late leafspot, Web blotch, Rust	13 fluid ounces	14 days before digging	<p>Apply EMINENT at intervals of 14 days, beginning when conditions become favorable for disease, generally around 30 days after planting.</p> <p>Applications of EMINENT should be alternated with a non-triazole fungicide which is registered for use on peanuts. If the alternate fungicide is known to provide protection for up to 14 days, applications of EMINENT should be no longer than 28 days apart.</p> <p>Do not make more than four (4) applications of EMINENT per growing season. Do not feed treated peanut hay to livestock.</p> <p>Crops other than peanuts must not be grown within 120 days following the last application of EMINENT 125SL.</p>

Pecan	Scab, Downy spot , Leaf blotch (Mycosphaerella spp.), Vein spot and/or Leafspot (Gnomonia spp), Leaf scorch, Powdery mildew	16 fluid ounces	30 days	Apply EMINENT 125SL in adequate water to provide complete coverage. Spray volumes of at least 100 gallons per acre should be used for ground applications and at least 10 gallons per acre for aerial applications. Apply of EMINENT 125SL at intervals of 14 to 21 days, beginning when conditions are favorable for scab or other foliage and nut hull diseases. Applications of EMINENT 125SL may be alternated or tank mixed with a non-triazole fungicide which is registered for use on pecans. Do not make more than eight applications of EMINENT 125SL per growing season.
-------	--	-----------------	---------	---

GENERAL CHEMIGATION INSTRUCTIONS:

Apply this product only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation system. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

For specific information about calibration, contact State Extension Service specialists, equipment manufacturers, or other irrigation experts.

Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Sprinkler Chemigation:

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add EMINENT slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, etc., should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank.

Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

EMINENT should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry, temperature-controlled, secure, place

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND LIMITATION OF DAMAGES

Conditions of sale: To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. SIPCAM AGRO USA, INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED. To the extent consistent with applicable law, SIPCAM AGRO USA, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND SIPCAM AGRO USA, INC.'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. SIPCAM AGRO USA, INC. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.

3/29/2007