



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 29, 2019

Patricia McFadden
Registration Manager
Sipcam Agro USA, Inc.
2525 Meridian Parkway, Suite 350
Durham, NC 27713

Subject: Label Notification per PRN 98-10 – Correct typographical errors for soybean tank
mix directions and resistance management section
Product Name: Stalwart C
EPA Registration Number: 60063-22
Application Date: 12/06/2018
Decision Number: 551431

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact Marc Sheahin at 703-347-8639 or via email at sheahin.marc@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Mindy Ondish".

Mindy Ondish
Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

NOTIFICATION

60063-22

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

08/29/2019

Metolachlor	Group	15	Herbicide
-------------	-------	----	-----------

Stalwart® C

For weed control in [Corn (field, pop, sweet, grown for seed)], [Cotton,] [Peanuts,] [Crop Group 6 Legume Vegetables (succulent and dried)] [Potatoes,] [Safflowers,] [Sorghum,] [Soybeans,] and [Tomato]

Active Ingredient:

Metolachlor: 2-chloro- N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl) acetamide 84.1%

Other Ingredients: 15.9%

Total: 100.0%

Contains 7.8 lbs. of metolachlor active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> •Call a poison control center or doctor immediately for treatment advice. •Have affected person sip a glass of water if able to swallow. •Do not induce vomiting unless told by a poison control center or doctor. •Do not give anything by mouth to an unconscious person.
IF IN EYES	<ul style="list-style-type: none"> •Hold eye open and rinse slowly and gently with water for 15-20 minutes. •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. •Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> •Move person to fresh air. •If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. •Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> •Take off contaminated clothing. •Rinse skin immediately with plenty of water for 15-20 minutes. •Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 900-4044 Poison Control Center

See additional precautionary statements and directions for use inside booklet.
 Read the entire label carefully before using this product.

OPTIONAL LANGUAGE FOR LABEL

[Peel back [label] [book] here] [Open book [label] here]

[Application Type] [AG] [Agriculture]

[Not for sale, use, or distribution in Nassau County or Suffolk County, New York.]

[Formulated in the United States of America, with U.S. and imported ingredients.]

[Product of _____] [if manufactured in a country other than U.S., country name will appear here]

[Herbicide]

EPA Reg. No. 60063-22
EPA Est. _____ [Lot number begins with xxx]
Net Contents: _____ [Gallons] [gal.][Liters]

[label date/lot code]

Manufactured For:
Sipcam Agro USA, Inc.
2525 Meridian Parkway, Suite 350
Durham, NC 27713

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eye wear.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt and long pants, socks, shoes, and gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

Coveralls over short-sleeved shirt and short pants

Chemical-resistant gloves: made of Barrier Laminate, Butyl Rubber ≥ 14 mils, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, Natural Rubber ≥ 14 mils, Polyethylene, Polyvinyl Chloride (PVC) ≥ 14 mils, or Viton ≥ 14 mils.

Chemical-resistant footwear plus socks

Chemical-resistant headgear for overhead exposure

Chemical-resistant apron when cleaning equipment mixing or loading.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

Mixers and loaders supporting aerial applications are required to use closed systems. The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4)). When using the closed system, the mixers and loaders PPE requirements may be reduced or modified as specified in the WPS. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

GROUND WATER ADVISORY

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

SURFACE WATER ADVISORY

Metolachlor can contaminate surface water through ground spray drift. Under some conditions, metolachlor may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

MIXING / LOADING INSTRUCTIONS

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates. Check-valves or antisiphoning devices must be used on all mixing and/or irrigation equipment.

This product may not be mixed or loaded within 50 ft. of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs. This product may not be mixed/loaded or used within 50 ft. of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FOR ALL TANK MIXTURES: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains

specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves: barrier laminate or viton
- Shoes plus socks

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES

Not for sale, use, or distribution in Nassau County or Suffolk County, New York.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

PRODUCT INFORMATION

Stalwart C is a selective herbicide registered for use as a preplant surface-applied, preplant incorporated, or preemergence treatment in water or fluid fertilizer for control of most annual grasses and certain broadleaf weeds in corn (field, pop, sweet, grown for seed), cotton, peanuts, crop group 6 legume vegetables (succulent and dried), potatoes, safflowers, sorghum, soybeans, and tomatoes. Stalwart C is also registered as a postemergent treatment on corn, cotton, potato, soybean, and tomato.

RESTRICTION:

DO NOT use in nurseries, turf, or landscape plantings.

DO NOT apply to frozen ground.

DO NOT apply under conditions which favor runoff or wind erosion of soil containing this product to nontarget areas. To prevent off-site movement due to runoff or wind erosion:

1. DO NOT treat powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface must first be settled by rainfall or irrigation.
2. DO NOT apply to impervious substrates, such as paved or highly compacted surfaces.
3. DO NOT use tailwater from the first flood or furrow irrigation of treated fields to treat nontarget crops, unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

Observe all use restrictions and precautions on the labels of each product used in tank mixtures. Tank mixtures are permitted only in those states where the tank mix partner is registered.

FOR ALL TANK MIXTURES: It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Where directions on this label specify a Stalwart C tank mixture with atrazine, follow the rates, restrictions, and use precautions on the labeling of the atrazine product used. Certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state

lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

If this product is incorporated, any supplemental tillage before planting must not exceed the depth of incorporation.

Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial weed control.

USE PRECAUTIONS

1. Injury may occur following the use of this product under abnormally high soil moisture conditions during early development of the crop.
2. Dry weather following preemergence application of this product or a tank mixture may reduce effectiveness. Cultivate if weeds develop.

RESISTANCE MANAGEMENT

For resistance management, **clomazone metolachlor** is a Group **15 43** herbicide. Any weed population may contain or develop plants naturally resistant to **clomazone metolachlor** and other Group **15 43** herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group **15 43** herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

INTEGRATED WEED PEST MANAGEMENT

Integrate this product into an overall weed pest management strategy whenever the use of an herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Consult local agricultural and weed authorities for additional IPM strategies established for your area.

SOIL TEXTURES AND HERBICIDE RATES

Where rates are based on *coarse-*, *medium-*, or *fine-textured soils*, it is understood that soil textural classes are generally categorized as follows:

Coarse	Medium	Fine	
Sand	Loam	Sandy clay loam	Sandy clay
Loamy sand	Silt loam	Silty clay loam	Silty clay
Sandy loam	Silt	Clay loam	Clay

Within rate ranges in the rate tables and elsewhere on this label, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter.

Stalwart C may be applied preemergence alone, or in combination with tank mix partners specified on this label, following preplant incorporated herbicides when used according to their label directions, provided that such use is not prohibited on the respective labels.

Thoroughly clean sprayer or other application device before using. Dispose of cleaning solution in a responsible manner. Do not use a sprayer or applicator contaminated with any other materials, or crop damage or clogging of the application device may result.

MIXING INSTRUCTIONS

Stalwart C Alone: Mix Stalwart C with water or fluid fertilizer (as specified in the individual crop sections) and apply as a spray. Fill the spray tank 1/2 - 3/4 full with water or fluid fertilizer, add the proper amount of this product, then add the rest of the water or fluid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

Tank Mixtures: Fill the spray tank one-quarter full with water, and start agitation; add tank mix partners (with the exception of paraquat or glyphosate), allow it to become dispersed; then add Stalwart C; then add paraquat, glyphosate/2,4-D, or glyphosate if these products are being used; and finally the rest of the water.

In some tank mixtures with atrazine, dicamba, linuron, simazine, pendimethalin, simazine; fluid fertilizers may replace all or part of the water as carrier, except in the atrazine postemergence and the dicamba post-emergence tank mixes. For tank mixtures with atrazine, see additional mixing instructions on the atrazine label. For each tank mixture, check compatibility with fluid fertilizer before mixing in spray tank. For all tank mixtures, conduct a compatibility test as described in the Compatibility Test Section of this label. For all tank mixtures, agitate during mixing and application to maintain a uniform suspension.

*See **Mixing Instructions** for tank mixtures with flumeturon, atrazine, or simazine+pendimethalin under the appropriate tank mixture section.

APPLICATION PROCEDURES

Application Timing

Stalwart C alone or in tank mixtures with other labeled herbicides may be applied for weed control in crops listed on this label. Refer to the individual crop sections of the label to determine if application timings listed below are applicable.

- a. Preplant Surface-Applied:** For minimum-tillage or no-tillage systems only, Stalwart C alone and some Stalwart C tank mixtures may be applied up to 45 days before planting. Use only split applications for treatments made 30-45 days before planting, with 2/3 the specified broadcast rate for the crop and soil texture applied initially and the remaining 1/3 at planting. Treatments less than 30 days before planting may be made either as a split or a single application. Refer to individual crop sections to determine if early preplant surface application is allowed.
If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (for example, paraquat or glyphosate). Observe directions for use, precautions, and restrictions on the label of the contact herbicide. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.
- b. Preplant Incorporated:** Apply Stalwart C to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If the crop will be planted on beds, apply and incorporate this product after bed formation, unless specified otherwise.
- c. Preemergence:** Apply this product during planting (behind the planter) or after planting, but before weeds or crops emerge.
- d. Postemergence (cotton and soybean use only):** For pre-emergence or partial control of the weeds listed in the WEEDS CONTROLLED – STALWART C APPLIED ALONE and in the WEEDS PARTIALLY CONTROLLED sections of this label, use one application of this product at the rate specified in the cotton or soybean sections of this label. This product alone will not control emerged weeds, so it must be applied to a weed-free surface or in a tank mixture with products that provide postemergence weed control. If weeds are present at the time of application, tank mix with a labeled postemergence herbicide and observe all directions for use, precautions, limitations, and restrictions on the label of the tank mix partner. For additional postemergence information, follow the crop specific label requirements identified on this label.

SPECIAL APPLICATION PROCEDURES

1. CA Only (Corn, Safflowers, Crop Group 6 Legume Vegetables (succulent and dried)):

Preplant Incorporated: Broadcast this product alone or with tank mix partners listed on this label to the soil and thoroughly incorporate with a disk or similar implement set to till 4-6 inches deep. For more thorough incorporation, till the soil in 2 different directions (cross-till). Crops may be planted on flat surface or on beds. Caution should be used when forming the beds that only soil from the Stalwart C treated zone is used (i.e., untreated soil should not be brought to soil surface). If the application is made to pre-formed beds, incorporate this product with a tillage implement set to till 2-4 inches deep. Care should be taken during tilling to keep the tilled (Stalwart C treated) soil on the beds.

2. Preemergence: Apply this product after planting. Water with sprinkler or flood irrigation within 7-10 days.

3. Fall Application (Only in IA, MN, ND, SD, WI, and North of Route 20 in the state of NE, and North of Route 136 in the state of IL – See specific instructions in the individual crop sections of this label for timing of application and other information): Use on medium and fine soils with greater than 2.5% organic matter that will be planted to corn or soybeans the next spring. Ground may be tilled before or after application.

RESTRICTIONS – Fall applications

1. DO NOT apply to frozen ground.
2. DO NOT exceed a 2 to 3-inch incorporation depth if tilled after treatment.
3. If a spring application is made, the total rate of the fall plus spring applications MUST NOT exceed the maximum total rate for the specific crop, or illegal residues may result.

4. Ground Application: Apply this product alone or in tank mixtures by ground equipment in a minimum of 10 gals. of spray mixture per acre, unless otherwise specified. Use sprayers that provide accurate and uniform application. For tank mixtures of this product with wettable powder or dry flowable formulations, screens and strainers should be no finer than 50-mesh. Rinse sprayer thoroughly with clean water immediately after use.

Calculate the amount of herbicide needed for band treatment by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

For information on applying in lower volumes of carrier, see the **Low Carrier Application Section**.

For application by air or through center pivot systems, see the **Aerial Application** section or **Center Pivot Irrigation Application** section.

For information on impregnating dry fertilizer, see the **DRY BULK GRANULAR FERTILIZERS** section.

WEEDS CONTROLLED – STALWART C APPLIED ALONE

Barnyardgrass (watergrass)	green foxtail	woolly cupgrass*	carpetweed
bristly foxtail	prairie cupgrass	yellow foxtail	common waterhemp
crabgrass	red rice	yellow nutsedge	Eastern black nightshade
crowfootgrass	robust foxtails (purple, white)		Florida pusley
fall panicum	signalgrass (<i>Brachiaria</i>)		galinsoga
foxtail millet	southwestern cupgrass		pigweed
giant foxtail	wild proso millet *		tall waterhemp
goosegrass	witchgrass		

*For control of these weeds in corn only, refer to the **Corn – Woolly Cupgrass and Wild Proso Millet Control Program** section of this label.

Weeds Partially Controlled*:

Common purslane	Eclipta,	Florida beggarweed**
Hairy nightshade	Sandbur	Seedling johnsongrass
Shattercane	Texas panicum***	Volunteer sorghum
Wild proso millet	Woolly cupgrass.	

** For partial control of this weed, use a minimum of 2.0 pts./A and apply preemergence.

*** For partial control of this weed, use a minimum of 2.0 pts./A and apply through a center pivot irrigation system.

Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial weed control. Control of these weeds can be erratic, due partially to variable weather conditions. Control may be improved by following these suggested procedures:

1. **Thoroughly till moist soil** to destroy germinating and emerged weeds. If this product is to be applied preplant incorporated, this tillage may be used to incorporate this product if uniform 2-inch incorporation is achieved as recommended under **Application Procedures**.
2. Plant crop into moist soil **immediately after tillage**. If this product is to be used preemergence, apply at planting or immediately after planting.
3. If available, **sprinkler irrigate** within 2 days after application. Apply one-half to 1 inch of water. Use lower water volume (one-half inch) on *coarse-textured* soils and higher volume (1 inch) on *fine-textured soils*. Also, refer to the section on **Center Pivot Irrigation Application** for this method of applying Stalwart C.
4. If irrigation is not possible and rain does not occur within 2 days after planting and application, weed control may be decreased. Under these conditions, a uniform, shallow cultivation is recommended as soon as weeds emerge.

ROTATIONAL CROPS

STALWART C ALONE

Replanting if a crop is lost

If a crop treated with this product alone is lost, any crop on this label may be replanted immediately if the rate from the previous crop does not exceed the rate for the crop to be planted. Do not make a second broadcast application of this product.

Rotational Crop Directions

(1) Barley, oats, rye, or wheat may be planted 4.5 months following treatment.

(2) Alfalfa may be planted 4 months following application. Clover may be seeded 9 months following application.

RESTRICTIONS: To avoid injury to rotational alfalfa or clover:

1. DO NOT apply more than 2.0 lbs. a.i. of metolachlor per acre (2.0 pts. of Stalwart C) preemergence (including preplant surface, preplant incorporated, postplant incorporated, etc.)
 2. DO NOT make lay-by or other postemergent applications of Stalwart C.
 3. DO NOT graze or feed forage or fodder from cotton to livestock.
- (3) Any crop on this label, in addition to root crops, tobacco, barley, buckwheat, milo, oats, rice, rye, wheat, cabbage, or peppers may be planted in the next spring following treatment.
- (4) Following a lay-by treatment or multiple treatments applied the previous season, any crop on this label, in addition to tobacco, cabbage, or peppers, may be planted in the spring. All other rotational crops may be planted 12 months after a lay-by application.
- (5) Do not graze or feed forage or fodder from cotton to livestock.

Stalwart C Tank Mixtures

For **Rotational Crops** restrictions for this product used in tank mixtures, refer to the statements/restrictions above for this product and to the respective product labels of any mixing partner(s) for additional statements/restrictions.

SPRAY EQUIPMENT

LOW CARRIER APPLICATION

For Broadcast Ground Application Only

Use sprayers that provide accurate and uniform application. **Only water may be used as a carrier.** Screens in suction and in-line strainers should be 50-mesh. Manufacturers may require that tip screens as fine as 100-mesh be used with some nozzles. Use a pump with capacity to: (1) maintain up to 35-40 psi at the nozzles, and (2) provide sufficient agitation in tank to keep mixture in

suspension. Use a minimum of 5.0 gals. of spray mixture per acre. Maximum recommended sprayer speed is 15 mph. Rinse sprayer thoroughly with clean water immediately after each use.

Low pressure nozzles are recommended to reduce drift and increase application accuracy. Care should be taken when using automatic rate-controlling devices to spray the material within the rated working pressure and flow ranges of the nozzles selected. Nozzle screens should be used when recommended by the manufacturer. All nozzles should be placed on 20-inch centers, except flooding types which should be placed on 40-inch centers. When Flat Fan-type nozzles are used, angles of 80° or 110° are recommended. Always read and follow the manufacturer's directions for optimum setup and performance of their nozzles or tips.

AERIAL APPLICATION

Apply Stalwart C in water alone or in tank mixtures with atrazine, or linuron, metribuzin in a minimum total volume of 2.0 gals./A by aircraft. This product may also be applied by air in combination with pendimethalin, or trifluralin. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft., using low-drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply Stalwart C alone or Stalwart C + atrazine by aircraft at a minimum upwind distance of 400 ft. from sensitive plants, or apply Stalwart C + linuron, or metribuzin at a minimum upwind distance of 300 ft. from sensitive plants.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

Aerial Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses, or to applications using dry formulations.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downward more than 45 degrees.

Where states have more stringent regulations, they must be observed.

The applicator must be familiar with and take into account the information covered in the **Aerial Drift Reduction Advisory Information** section below.

Aerial Drift Reduction Advisory Information Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the **Wind, Temperature and Humidity**, and **Temperature Inversions** sections).

Controlling Droplet Size

- **Volume** – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

- **Pressure** – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles** – Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 ft. above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential.

Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

CENTER PIVOT IRRIGATION APPLICATION

Stalwart C alone or in tank mixture with other herbicides on this label, which are registered for center pivot application, may be applied in irrigation water preemergence (after planting, but before weeds or crop emerge) at rates specified on this label. This product also may be applied postemergence to the crop and preemergence to weeds in crops where postemergence applications are allowed on this label. Follow all restrictions (height, timing, rate, etc.) to avoid illegal residues. **Apply this product only through a center pivot irrigation system. Do not apply this product through any other type of irrigation system.** Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension specialists, equipment manufacturers, or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being with-drawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump or piston pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Prepare a mixture with a minimum of 1 part of water to 1 part herbicide(s) and inject this mixture into the center pivot system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the herbicide in suspension.
9. Meter into irrigation water during entire period of water application.
10. Apply in $\frac{1}{2}$ - 1 inch of water. Use the lower water volume ($\frac{1}{2}$ inch) on *coarse-textured soils* and the higher volume (1 inch) on *fine-textured soils*. More than 1 inch of water at application may reduce weed control by moving the herbicide below the effective zone in the soil.

PRECAUTIONS FOR CENTER PIVOT APPLICATIONS

1. Where sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result.
2. Where sprinkler distribution patterns overlap excessively, crop injury may result.

DRY BULK GRANULAR FERTILIZERS

Many dry bulk granular fertilizers may be impregnated or coated with this product alone or selected Stalwart C tank mixtures which are registered for preplant incorporated or preplant surface applications which are used to control weeds on labeled crops on the Stalwart C label and are not prohibited from use on dry bulk granular fertilizers.

When applying Stalwart C or Stalwart C mixtures with dry bulk granular fertilizers, follow all directions for use and precautions on the respective product labels, regarding target crops, corn, rates per acre, soil texture, application methods (including timing of application), and rotational crops.

All individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the herbicide/fertilizer mixture.

Prepare the herbicide/fertilizer mixtures by using any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender. Nozzles used to spray Stalwart C and Stalwart C mixtures onto the fertilizer must be placed to provide uniform spray coverage. Care should be taken to aim the spray directly onto the fertilizer only and to avoid spraying the walls of the blender.

If the herbicide/fertilizer mixture is too wet, add a highly absorptive material, such as Agsorb® or Celatom MP-79®, or similar granular clay or diatomaceous earth materials, to obtain a dry, free-flowing mixture. Absorptive materials should be added only after the herbicide has been thoroughly blended into the fertilizer mixture. Best application results will be obtained by using a granule of 6/30 particle size or of a size similar to that of the fertilizer material being used. Generally, less than 2% by weight of absorptive material will be needed. Avoid using more than 5% absorptive material by weight.

Calculate amounts of Stalwart C, atrazine, atrazine plus simazine, metribuzin, isoxaflutole or ethafluralin by the following formula:

$$\frac{2,000}{\text{lbs. of fertilizer per acre}} \times \text{pts./A of liquid or flowable product} = \text{pts. of liquid or flowable product per ton of fertilizer}$$
$$\frac{2,000}{\text{lbs. of fertilizer per acre}} \times \text{lbs./A of dry product} = \text{lbs. of dry product per ton of fertilizer}$$

Pneumatic (Compressed Air) Application (Stalwart C Alone): High humidity, high urea concentrations, low fertilizer use rates, and dusty fertilizer may cause fertilizer mixture to build up or plug the distributor head, air tubes, or nozzle deflector plates. To minimize buildup, premix Stalwart C with Exxon Aromatic 200 at a rate of 1.0 to 4.0 pts./gal. of Stalwart C. Aromatic 200 is a noncombustible / nonflammable petroleum product. Aromatic 200 may be used in either a fertilizer blender or through direct injection systems. Drying agents should not be used when using Aromatic 200.

PRECAUTIONS

1. Mixtures of Stalwart C and Aromatic 200 must be used on dry fertilizer only. Poor results or crop injury may result if these mixtures are used in water or liquid fertilizer solutions for spraying applications.
2. When impregnating Stalwart C in a blender before application, a drier mixture can be attained by substituting a drying agent for Aromatic 200. The use of Agsorb FG or drying agents of 6/30 particle size are recommended.
3. Drying agents are not recommended for use with On-The-Go impregnation equipment.

RESTRICTIONS

To avoid potential for explosion,

1. Do not impregnate Stalwart C or Stalwart C mixtures on ammonium nitrate, potassium nitrate, or sodium nitrate, either alone or in blends with other fertilizers.
2. Do not use Stalwart C or Stalwart C mixtures on straight limestone, since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated.

Application

Apply 200-700 lbs. of the herbicide/fertilizer mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury. Non-uniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil may improve weed control. On *fine-* or *medium-textured soils* in areas where soil incorporation is not planned, i.e., reduced tillage situations or in some conventional till situations, make applications approximately 30 days before planting to allow moisture to move the herbicide/fertilizer mixture into the soil. On *coarse-textured soils*, make applications approximately 14 days prior to planting.

USE PRECAUTION:

To avoid crop injury, do not use the herbicide/fertilizer mixture on crops where bedding occurs.

COMPATIBILITY TEST

Because liquid fertilizers can vary, even within the same analysis, always **check compatibility with herbicide(s) each time before use**. Be especially careful when using **complete** suspension or fluid fertilizers, as serious compatibility problems are more likely to occur. Commercial application equipment may improve compatibility in some instances. The following test assumes a spray volume of 25 gals./A. For other spray volumes, make appropriate changes in the ingredients. Check compatibility using this procedure:

1. Add 1.0 pt. of fertilizer to each of 2 one-qt. jars with tight lids.
2. To **one** of the jars, add 1 /4 tsp. or 1.2 milliliters of a compatibility agent approved for this use (1 /4 tsp. is equivalent to 2.0 pts./100 gals. spray). Shake or stir gently to mix. When an adjuvant is to be used with this product, Sipcam Agro USA recommends the use of Compex[®], Unite[®] or a Chemical Producers and Distributors Association (CPDA) certified adjuvant.
3. To **both** jars, add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:
Dry herbicides: For each pound to be applied per acre, add 1.5 level teaspoons to each jar.
Liquid herbicides: For each pint to be applied per acre, add 0.5 teaspoon or 2.5 milliliters to each jar.
4. After adding all ingredients, put lids on and tighten, and invert each jar 10 times to mix. Let the mixtures stand 15 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the

spray mixture by comparing the 2 jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) slurry the dry herbicide(s) in water before addition, or (B) add 1/2 of the compatibility agent to the fertilizer and the other 1/2 to the emulsifiable concentrate or flowable herbicide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

CROPS

CORN (FIELD, POP, SWEET, GROWN FOR SEED) – STALWART C ALONE

Apply this product, either preplant surface, preplant incorporated, or preemergence, using the appropriate rate specified below.

1. PREPLANT SURFACE-APPLIED

Refer to instructions for use of Stalwart C alone under **Application Procedures**.

a. Fall Application

- Apply after September 30 in MN, ND, SD, WI, and north of Route 30 in IA;
- Apply after October 15 north of Route 91 in NE and south of Route 30 in IA;
- Apply after October 31 north of Route 136 in IL

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67-2.0 pts./A on *medium-textured* and 2.0 pts./A on *fine-textured soils*. A tillage operation may precede the application. Minimize furrow and ridge formation in the tillage operations.

RESTRICTIONS:

1. If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for corn, or illegal residues may result.
2. DO NOT apply to frozen ground.
3. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2-3 inches.

- b. Use on *medium-* and *fine-textured soils* with minimum-tillage or no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY. Apply 2/3 the specified rate of Stalwart C (1.67 pts./A on *medium soils* and 2.0 pts./A on *fine soils*) as a split treatment 30-45 days before planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. Apply 1.33 pts./A on *coarse soils* not more than 2 weeks prior to planting.

RESTRICTION: If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for corn, or illegal residues may result.

- c. On *medium-* and *fine-textured soils* with minimum- or no-tillage systems in CT, DE, MA, MD, ME, MI, NH, NY, OH, PA, RI, VA, VT, and WV, preplant surface applications may be applied following the directions for use above. If the amount of rainfall results in unsatisfactory length of weed control following the earlier treatment, a post-emergence application of an appropriately labeled broadleaf and/or grass weed herbicide may be used, i.e., metolachlor, primisulfuron-methyl, prosulfuron, nicosulfuron, dicamba, bromoxynil, or 2,4-D. If the postemergence treatment includes the herbicide used preplant surface-applied, do not exceed the total labeled rate for corn on a given soil texture.

Observe all directions for use, precautions, and restrictions on the label of the postemergent herbicide.

2. PREPLANT INCORPORATED OR PREEMERGENCE

Follow instructions for use of Stalwart C alone under **Application Procedures**.

On *coarse soils*, apply 1.0-1.33 pts./A of Stalwart C if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On *medium soils*, apply 1.33-1.67 pts./A of Stalwart C. On *fine soils*, apply 1.33-1.67 pts./A of Stalwart C if organic matter content is less than 3%, or 1.67-2.0 pts./A if organic matter content is 3% or greater.

CALIFORNIA ONLY

Preplant Incorporated: See the **SPECIAL APPLICATION PROCEDURES** Section of this label.

3. LAY-BY OR POSTEMERGENCE: To extend the duration of weed control in corn, a maximum rate of 2.0 pts./A of Stalwart C may be applied after crop emergence until the corn plants reach 40 inches in height, following any preplant surface-applied, preplant incorporated, or preemergence herbicide application, including Stalwart C. For best results, applications should be made to soil free of emerged weeds and directed towards the base of the corn plants in excess of 5 inches tall. The total Stalwart C rate applied on corn during any one crop year must not exceed 4.0 pts./A, depending on soil texture.

Shattercane, Wild Proso Millet, Woolly Cupgrass, and Eclipta – Partial Control: For more consistent partial control of shattercane, wild proso millet, woolly cupgrass, or eclipta, apply 2.0-2.55 pts./A as a single application; or apply 1.0-1.33 pts./A of Stalwart C preplant incorporated followed by 1.0-1.33 pts./A of Stalwart C preemergence; however, do not apply more than a total of 2.55 pts./A. Make the preemergence application during or after planting, but before weeds and corn plants emerge. Apply the 1.33 pts./A rate of Stalwart C when a heavy infestation of shattercane, wild proso millet, woolly cupgrass, or eclipta is expected. A shallow cultivation may be needed to control any late emerging weeds.

Woolly Cupgrass and Wild Proso Millet Control Program: For control of these species, use the following 3-step program:

1. Apply Stalwart C early preplant, preplant incorporated, or preemergence at 1.67 pts./A on *medium soils* and 2.0 pts./A on *fine-textured soils*, up to the maximum label rate. Lightly incorporate with a rotary hoe if rainfall does not occur within 5-7 days;
2. Apply a postemergence tank mix of primisulfuron-methyl at 0.38 oz./A or Exceed primisulfuron-methyl and prosulfuron at 1 packet per 4 acres plus nicosulfuron at 0.33 oz./A plus 1.0 qt. of crop oil concentrate plus 1.0 gal./A of 28% nitrogen, or the equivalent amount of ammonium sulfate, when grasses are 2-3 inches tall and the corn plant is at least 4 inches tall; and
3. Cultivate 14-21 days after the postemergence application.

RESTRICTIONS (for all application methods):

1. To avoid possible illegal residues, DO NOT graze or feed forage from treated areas for 30 days following application.
2. DO NOT apply more than the labeled application rate for a given soil texture per year, either as a single or split treatment, or illegal residues may result.
3. In corn, Stalwart C may be used up to 2.75 pts./A as either a preplant surface, preplant incorporated, or preemergence treatment on soils having an organic matter content between 6% and 20% or up to 2.0 pts./A on any soil for extended residual control and where severe stands of problem weeds are expected.
4. In the event of escape of annual weeds following a preplant surface, preplant incorporated, or pre-emergence treatment of Stalwart C, follow with a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide, i.e., atrazine, primisulfuron-

methyl, metolachlor, primisulfuron-methyl, prosulfuron, nicosulfuron, dicamba, bromoxynil, or 2,4-D. If the postemergence treatment includes the herbicide used in the earlier treatment, i.e., atrazine, do not exceed the total labeled rate for corn on a given soil texture.

5. Bromoxynil may be applied postemergence alone or in tank mix combination with atrazine. DO NOT exceed 1.2 lbs. a.i./A of atrazine in tank mix combination with bromoxynil postemergence. Refer to the atrazine, bromoxynil labels for specific rates and precautions.
6. Do not use Stalwart C on peat or muck soils.
7. Do not apply to frozen ground.

CORN (FIELD, POP, SWEET, GROWN FOR SEED) STALWART C COMBINATIONS

Stalwart C in any tank mixture for corn (except Stalwart C + atrazine postemergence and Stalwart C + dicamba postemergence) may be applied in water or fluid fertilizer. Use only water in the Stalwart C + atrazine or the Stalwart C + dicamba postemergence tank mixes.

IMPORTANT: FOR ALL TANK MIXTURES – If applying Stalwart C in tank mixtures, all the restrictions and rate limitations on the mix partner label(s) must be followed if more restrictive than those on this label.

Stalwart C Tank Mixtures for Corn – Additional Weeds Controlled and Special Instructions

	Stalwart C + Atrazine and/or Simazine (1)	Stalwart C + Atrazine	Stalwart C + Dicamba (Field Corn)	Stalwart C + Atrazine + Linuron	Stalwart C + Atrazine or Simazine + Pendimethalin	Stalwart C + Atrazine / dicamba
Comments	2,3,4,6,7	2,3,4		2,3,4,5	1, 2,3,4	6
Browntop panicum	a			a	A	
Cocklebur	a	b	b	a	A	a
Common purslane	a			a	A	a
Hairy nightshade	a			a	A	a
Jimsonweed		a	b			a
Kochia		a				a
Lambsquarters	a	a	a	a	a	a
Morningglory	a	b	b	a	a	a
Mustard		a				a
Pigweed				a	a	a
Prickly sida		a				a
Ragweed	a	a	a	a	a	a
Smartweed	a	a	a	a	a	a
Velvetleaf	a	a	b	a	a	a

a = control; b = partial control.

(1) Preplant Surface, PPI, PRE

INSTRUCTIONS

1. Special Mixing Instructions for Stalwart C + atrazine or simazine and pendimethalin

- a) Fill the spray tank 1/4 full with water or fluid fertilizer and start agitation.
- b) To aid compatibility, add a compatibility agent, such as Unite® or X-77®, at 4.0 pts./100 gals. of spray mixture.
- c) Then add the atrazine or simazine and allow it to become dispersed.

- d) Then add Stalwart C and pendimethalin.
- e) Add the rest of the water.
2. Follow the directions for use, rates, and restrictions on the tank mix partner label(s).
3. In Minimum-Tillage and No-Tillage systems, mix with paraquat for control of most emerged annual weeds and suppression of perennial weeds; or with glyphosate/2, 4-D for suppression of emerged field bindweed and control or suppression of annual weeds; or with glyphosate for control of most emerged annual and perennial weeds.
4. Refer to **TANK MIXTURE WITH ATRAZINE; OR ATRAZINE + 2,4-D; OR ATRAZINE + 2,4-D + DICAMBA FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS** in the corn section of this label for specific directions for 2,4-D or dicamba burndown combinations in Minimum-Tillage and No-Tillage systems.
5. Stalwart C in any tank mixture for corn may be applied in water or fluid fertilizer, except as noted.
6. Refer to **Corn (Field, Pop, Sweet, Grown For Seed) – Stalwart C Alone**, for sequential postemergence treatments if escape weeds develop.

RESTRICTIONS FOR ALL TANK MIXES USED ON CORN

1. For all applications to corn, do not graze or feed forage from treated areas for 30 days following application, or possible illegal residues may result.
2. When applying Stalwart C in tank mixture with atrazine, do not exceed the specified amount of atrazine per acre per year. Certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.
3. In corn, Stalwart C may be used up to 2.0 pts./A in combinations on any soil for extended residual control and where severe stands of problem weeds are expected.
4. Do not apply more than the labeled rate for a given soil texture per year, either as a split or single treatment, or illegal residues may result.

1. TANK MIXTURE WITH ATRAZINE OR SIMAZINE, OR ATRAZINE + SIMAZINE– PREPLANT SURFACE, PREPLANT INCORPORATED, OR PREEMERGENCE

In addition to the weeds controlled by Stalwart C alone, Stalwart C+atrazine or simazine, or Stalwart C+=atrazine + simazine, applied preplant surface, preplant incorporated, or preemergence, also controls the following weeds: browntop panicum, cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

Apply Stalwart C + atrazine or simazine, or Stalwart C + atrazine + simazine either preplant surface, preplant incorporated, or preemergence.

Preplant Surface-Applied: Follow instructions for use of Stalwart C alone under **Application Procedures** and under application instructions for Stalwart C alone on corn. Apply Stalwart C + atrazine or simazine, or Stalwart C + atrazine + simazine on *medium soils* (1.67 pts./A of Stalwart C + the labeled rate of atrazine or simazine, or atrazine+ simazine combined) and on *fine soils* (1.67-2.0 pts./A of Stalwart C + the labeled rate of atrazine or simazine, or atrazine+simazine combined) in minimum-tillage and no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY. Apply the tank mixtures as a split or single treatment in those states and as indicated in the **Stalwart C Alone – Preplant Surface-Applied** section of the label. On *coarse soils*, apply 1.33 pts./A of Stalwart C and the labeled rate of atrazine or simazine, or atrazine+simazine combined.

Preplant Incorporated or Preemergence: Follow instructions for use of Stalwart C alone under **Application Procedures**. Apply Stalwart C + atrazine or simazine, or Stalwart C + atrazine + simazine, using the appropriate rates from Table 1.

CALIFORNIA ONLY

Preplant Incorporated: See the **SPECIAL APPLICATION PROCEDURES** Section of this label.

Shattercane, Wild Proso Millet, Woolly Cupgrass, and Eclipta – Partial Control

For more consistent partial control of shattercane, wild proso millet, woolly cupgrass, or eclipta, where Stalwart C is applied in tank mixture or sequentially with other registered corn herbicides, apply 2.0-2.33 pts. as a single application, or the following applications may be made:

1. Apply 1.0-1.33 pts./A of Stalwart C + the labeled rate of atrazine or simazine preplant incorporated, followed by 1.0-1.33 pts./A of Stalwart C preemergence. Make the preemergence application during or after planting, but before weeds and corn plants emerge.
2. Apply Stalwart C at 1.33 pts./A alone or in tank mix combination with up to the labeled rate of atrazine or simazine, preplant incorporated. Do not exceed the total rate of triazine herbicide specified for corn grown on a given soil texture. Follow with a post-directed application of ametryn at the labeled rate. Refer to the ametryn label for specific directions for the post-directed application.
3. Apply EPTC or butylate at labeled rates preplant incorporated, followed by a preemergence application of Stalwart C at 1.0-1.33 pts./A. Do not use EPTC or butylate on soils where rapid degradation has been shown to occur. Make the preemergence application during or after planting, but before weeds and corn plants emerge.

PRECAUTION: When following the application regimes in numbers 1-3 above, a shallow cultivation may be needed after the preemergence or postemergence application to help control any late emerging shattercane or wild proso millet plants.

Table 1: Stalwart C + atrazine or simazine, or Stalwart C + atrazine + simazine, Preplant Incorporated, or Preemergence – Corn (Field, Pop, Sweet, Grown For Seed)

Soil Texture	Broadcast Rates of Stalwart C Per Acre			
	Less than 3% Organic Matter		3% Organic Matter or Greater	
COARSE	0.85-1.0 pts. + the label rate of either atrazine or simazine	0.85-1.0 pts. + the label rates of atrazine and simazine	1.0 pts. + the label rate of either atrazine or simazine	1.0 pts. + the label rates of atrazine and simazine
MEDIUM	1.0-1.33 pts. + the label rate of either atrazine or simazine	1.0-1.33 pts. + the label rates of atrazine and simazine	1.33 pts. + the label rate of either atrazine or simazine	1.33 pts. + the label rates of atrazine and simazine
FINE	1.33 pts. + the label rate of either atrazine or simazine	1.33 pts. + the label rates of atrazine and simazine	1.33-1.67 pts. + the label rate of either atrazine or simazine	1.33-1.67 pts. + the label rates of atrazine and simazine
DO NOT USE ON MUCK OR PEAT (SOILS WITH MORE THAN 20% ORGANIC MATTER)				

* Use simazine in preference to atrazine when heavy infestations of crabgrass or fall panicum are expected. On soils having between 6% and 20% organic matter, Stalwart C may be used up to 2.33 pts./A in tank mix combination with the label rate of atrazine. Refer to the atrazine label for weeds controlled at this rate.

2. TANK MIXTURE WITH ATRAZINE– POSTEMERGENCE

Weeds Controlled			
Barnyardgrass (watergrass) crabgrass crowfootgrass fall panicum	giant foxtail green foxtail yellow foxtail ----- jimsonweed	Kochia Lambsquarters Mustard pigweed prickly sida	Purslane Ragweed smartweed velvetleaf
Weeds Partially Controlled			
cocklebur	morning glory	yellow nutsedge	

Apply 1.0 pt./A of Stalwart C + the label rate of atrazine on *coarse soils*, 1.33 pts./A of Stalwart C + the label rate of atrazine on *medium soils*, or 1.33-1.67 pts./A of Stalwart C + the label rate of atrazine on *fine soils*. Apply this tank mixture before grass and broadleaf weeds pass the 2-leaf stage and before corn exceeds 5 inches in height. Application to weeds larger than the 2-leaf stage will generally result in unsatisfactory control.

Lay-by: Apply to corn plants not more than 12 inches tall. Applications to corn plants in excess of 5 inches should be directed to the base of the plants; whereas, applications to plants less than 5 inches tall may be made over the top. Occasionally, some corn leaf burn may result, but this should not affect later growth or yield. Do not apply this postemergence tank mixture in fluid fertilizer, or severe crop injury may occur.

For better control of cocklebur, morningglory, velvetleaf, and yellow nutsedge on *fine-textured soils* above 3% organic matter, apply the label rate of atrazine with 1.33-1.67 pts./A of Stalwart C.

Tank mixtures of Stalwart C + atrazine may be applied following use of any registered preplant surface-applied, preplant incorporated, or preemergence corn herbicide, including Stalwart C + atrazine.

RESTRICTIONS

1. The total Stalwart C rate must not exceed 4.0 pts., and the atrazine rate must not exceed the amount allowed on the label during any one crop year, or illegal residues may result.
2. Refer to the atrazine label for geographic, soil-texture, and rotational restrictions.

3. TANK MIXTURE WITH DICAMBA

Preemergence: Use this tank mixture only on field corn, which is flat-planted (no furrows) in CO, IA, IL, IN, KS, MN, NE, OH, SD, and WI if the tank mix partner is registered in the state.

In addition to the weeds controlled by Stalwart C alone, Stalwart C + dicamba, applied preemergence, also controls lambsquarters, ragweed, smartweed, cocklebur*, jimsonweed*, morningglory*, and velvetleaf*.

*Partially controlled.

Apply Stalwart C + dicamba preemergence. Broadcast the label rate of dicamba with 1.33 pts./A of Stalwart C on *medium soils*, or with 1.33-1.67 pts./A of Stalwart C on *fine soils*. Do not apply on *coarse soils* or on soils with less than 2.5% organic matter. Apply this tank mixture to the soil surface at planting or after planting, but before field corn emerges. Plant seed at least 1.5 inches deep and apply behind planting equipment, avoiding incorporation by the planter wheel or other seed-covering device. Do not incorporate before corn plants emergence. If it is necessary to rotary hoe to break the soil crust, do not disturb the soil more than 1/2 inch deep.

Postemergence for Control of Pigweed (Mid-Atlantic states, including DE, MD, PA, VA, and WV): Apply 1.0-1.5 pts. of Stalwart C + the label rate of dicamba by ground equipment when pigweed

plants are less than 3 inches tall and before field corn exceeds 5 inches in height in a minimum of 20 gals. of spray per acre. Use the lower specified rate on *coarse-textured* and low organic matter soils. Use the higher specified rate on *fine-textured* and high organic matter soils.

RESTRICTIONS FOR ALL STALWART C AND DICAMBA TANK MIXES ON CORN

1. Avoid drift to sensitive nontarget plants, such as soybeans, during application, or injury may occur.
2. Do not apply with aircraft.

4. TANK MIXTURE WITH ATRAZINE AND LINURON FOR CONTROL OF LAMBSQUARTERS AND PIGWEED

For prolonged control of lambsquarters and pigweed in DE, MD, NJ, NY, PA, VA, and WV, Stalwart C may be applied preemergence in tank mix combination with atrazine + the label rate of linuron. Apply Stalwart C and atrazine according to the rates in Table 1 and linuron according to the labeled rates.

Observe all directions for use, precautions, and restrictions on the Stalwart C, atrazine, and linuron labels when applying these products in tank mix combinations.

5. TANK MIXTURE WITH ATRAZINE OR SIMAZINE +PENDIMETHALIN FOR PROLONGED CONTROL OF LAMBSQUARTERS AND PIGWEED IN FIELD CORN ONLY (NORTHEAST U.S., INCLUDING MI, IN, KY, AND STATES EAST OF THESE)

For prolonged control of lambsquarters and pigweed, in addition to a broad spectrum of annual broadleaf and grass weeds, Stalwart C in tank mix combination with atrazine* or simazine + pendimethalin may be applied after planting, but before field corn or weeds emerge. Apply by ground equipment in a minimum of 10 gals. of water or 20 gals. of liquid fertilizer. Apply by air in a minimum of 5.0 gals. of water. Refer to Table 1 of this label for rates of Stalwart C to be applied. Apply pendimethalin, atrazine, or simazine at the label rate.

Some formulations of atrazine and pendimethalin are not compatible. Before using this tank mixture, a compatibility test must be conducted. See the **COMPATIBILITY TEST** section of this label.

Mixing Instructions:

1. Fill the spray tank 1/4 full with water or fluid fertilizer and start agitation.
2. To aid compatibility, add a compatibility agent, such as Unite[®] or X-77[®], at 4.0 pts./100 gals. of spray mixture.
3. Then add the atrazine or simazine and allow it to become dispersed.
4. Then add Stalwart C and pendimethalin.
5. Add the rest of the water.

Observe all directions for use, precautions, and restrictions on the respective product labels when applying these products in tank mix combination. Refer to the pendimethalin label for replanting instructions in the event of crop loss.

6. TANK MIXTURE WITH ATRAZINE OR SIMAZINE, OR ATRAZINE + SIMAZINE, WITH PARAQUAT, GLYPHOSATE +2,4-D, OR GLYPHOSATE FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides paraquat, glyphosate or glyphosate+2,4-D, may be added to a tank mix of Stalwart C + atrazine and/or simazine, mixed with paraquat for control of most emerged annual weeds and suppression of perennial weeds; or with glyphosate/2, 4-D for suppression of emerged field bindweed and control or suppression of annual weeds; or with glyphosate for control of most emerged annual and perennial weeds. The Stalwart C + atrazine or simazine, or Stalwart C + atrazine + simazine portion of the tank mixture provides

preemergence control of the weeds listed on this label in the tank mixture section for **STALWART C + ATRAZINE OR SIMAZINE, OR STALWART C + ATRAZINE + SIMAZINE PREPLANT SURFACE, PREPLANT INCORPORATED, OR PREEMERGENCE.**

Application: Apply before, during, or after planting, but before the corn emerges, at the rates specified below. Add paraquat, glyphosate or glyphosate/2, 4-D at the labeled broadcast rate. See the paraquat, glyphosate or glyphosate/2, 4-D labels for weeds controlled, labeled rates for specific weeds, and other use directions.

- Apply surfactant at 1.0 or 2.0 pts./100 gals. of spray mixture with 75% or greater or 50-74% nonionic active ingredient, respectively. This treatment will not control weeds taller than 6 inches.
- Apply in 20-60 gals. of water or fluid fertilizer per acre with ground equipment.
- On *coarse soils*, apply 1.0 pt./A of Stalwart C with the labeled rate of all tank mix partners,
- On *medium soils*, apply 1.33 pts./A of Stalwart C with the labeled rate of all tank mix partners,
- On *fine soils****, apply 1.33-1.67 pts./A of Stalwart C with the labeled rate of all tank mix partners,

* Use simazine in preference to atrazine when heavy infestations of crabgrass or fall panicum are expected.

** When using the tank mixture of Stalwart C + atrazine + simazine, use equal rates of atrazine and simazine as shown when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of atrazine + simazine instead of the 1:1 ratio.

*** For cocklebur, yellow nutsedge, and velvetleaf control on *fine-textured soils* above 3% organic matter, apply the specified labeled rate of atrazine and/or simazine, with 1.33-1.67 pts./A of Stalwart C.

RESTRICTION: Do not apply combinations containing paraquat in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

7. TANK MIXTURE WITH ATRAZINE; OR ATRAZINE + 2,4-D; OR ATRAZINE + 2,4-D + DICAMBA FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, Stalwart C applied in combination with atrazine will kill most emerged small annual weeds. Apply Stalwart C + atrazine before, during, or after planting, but before corn emerges, according to the rates in Table 1.

Where heavy crop debris exists, add the label rate of an appropriately labeled 3.8 lbs. a.i./gal. of 2,4-D amine (such as Weedar® 64 or Formula 40®) to the spray tank last and apply in a minimum of 25 gals. of carrier per acre.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burndown of existing weeds, and therefore, are recommended instead of water. Add X-77 surfactant at 1.0-2.0 qts./100 gals. of diluted spray, or another appropriate surfactant at its recommended rate, or add crop oil concentrate plus 28% liquid nitrogen (or equivalent). Apply before weeds exceed 3 inches in height. If alfalfa is present, add dicamba to the spray mixture at the labeled rate and apply before alfalfa exceeds 6 inches in height.

For fields with existing sod grasses (e.g., brome grass, orchardgrass, rye, or timothy), when existing weeds exceed 3 inches in height or when very dry conditions exist, add paraquat at the label rate in place of or in addition to 2,4-D, as indicated above. Do not apply paraquat in suspension-type liquid fertilizer. Observe all directions for use, precautions, and restrictions on the respective product labels when applying these products in tank mix combination.

8. TANK MIXTURE WITH DICAMBA / ATRAZINE IN CONSERVATION TILLAGE – FIELD CORN

In conservation tillage systems where field corn is planted directly into a cover crop or previous crop residue, Stalwart C + dicamba / atrazine will kill most emerged small annual weeds. Apply Stalwart C + dicamba / atrazine before, during, or after planting, but before field corn emergence on *medium* and *fine soils* with greater than 2.5% organic matter. For fields with existing vegetation exceeding 3 inches in height or when very dry conditions exist, add paraquat at its labeled rate. Stalwart C + dicamba / atrazine may be applied postemergence to field corn plants less than 3 inches tall and before weedy grasses exceed the 2-leaf stage. As carriers, nitrogen solutions and complete liquid fertilizers, applied before crop emergence enhance burndown of existing weeds. Do not apply paraquat in suspension-type liquid fertilizer or use on emerged crop.

Refer to the dicamba / atrazine label and follow all directions, restrictions, and use precautions regarding application and use in field corn.

COTTON – STALWART C ALONE

Application: Apply Stalwart C preemergence only in Area 1* at the rate of 0.75 to 1 pt./A on sandy loams, 1 to 1.33 pts./A on *medium soils*, or 1 to 1.33 pts./A on *fine soils*. Apply this product preplant incorporated or preemergence in Area 2** at 1 pt./A on sandy loams, 1 to 1.33 pts./A on *medium soils*, or 1.33 pts./A on *fine soils*. Apply this product postemergence to cotton and preemergence to weeds at 0.75 to 1.33 pts./A according to the state rate restrictions in the **Postemergence** section below.

DO NOT use on sands and loamy sand.

* Area 1 = AR, LA, MS, TN, and Bootheel of MO

** Area 2 = NM, OK, and TX

Preplant Incorporated (NM, OK, and TX Only): Apply Stalwart C to the soil and incorporate into the top inch of soil immediately before planting, at planting, or after planting, but before crop or weeds emerge. Use a rolling cultivator or similar implement to uniformly incorporate not more than 1 inch deep. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. Where furrow irrigation is used, wet the top of the bed for best results. If the crop is to be planted on beds, apply and incorporate after bed formation. Cotton should be planted below the zone of incorporation; i.e., at least 1 inch on *fine soils* and 1.5 inches on *coarse* and *medium soils*. If incorporated prior to planting, use a planter that will result in a minimum of soil disturbance.

RESTRICTIONS:

1. For best control of yellow nutsedge and suppression of seedling johnsongrass, apply this product preplant incorporated at the maximum rate for the soil texture, whether applied alone or mixed with prometryn.

Preemergence: Apply Stalwart C to the soil surface at planting or after planting, but before weeds or crop emerge.

Postemergence: Apply Stalwart C broadcast over-the-top or directed to the soil surface, according to the rate and cotton height restrictions listed below by state. Application before weeds emerge or after clean cultivation to remove existing weeds is necessary as this product will not control emerged weeds. This product postemergence may be applied over any previous registered herbicide treatment. In sprinkler-irrigated areas, sprinkler irrigate after application with one-half to 1 inch of water (one-half inch on *coarse-textured soils* to 1 inch on *fine-textured soils*). To incorporate this product in furrow-irrigated areas, apply this product, incorporate with a rolling cultivator or similar implement that provides uniform shallow incorporation (2 inches or less), and then irrigate. In non-irrigated areas, if at least one-half inch of rainfall does not occur within 10 days after application, cultivate with a rolling cultivator or similar implement that provides uniform shallow incorporation of this product.

VA, NC, SC, GA, FL, and AL	Apply Stalwart C at 1 to 1.33 pts./A when cotton is 3 to 6 inches tall.
TN, AR, MS, MO, and LA	Apply Stalwart C at 0.75 to 1.33 pts./A when cotton is 3 to 12 inches tall.
TX, OK, NM, AZ, CA, and Clay Soils in AR	Apply Stalwart C at 1 to 1.33 pts./A when cotton is 3 to 12 inches tall, but before August 1.

Multiple Applications: Where weed pressure is heavy, difficult-to-control species are expected, or reinfestation may occur, and a weed control program is used, multiple applications of this product are effective when used as part of the weed control program. Apply as a preplant incorporated or preemergence treatment and follow with an application postemergence to cotton before weeds emerge or after clean cultivation to remove existing weeds since this product will not control emerged weeds. Cotton should be at least 3 inches tall at the postemergence timing. Apply this product postemergence over a previous preplant or preemergence application of this product as shown in the following table.

Stalwart C Multiple Applications to Cotton			
State	Preplant incorporated or Preemergence Pts./A		Postemergence and Height Pts./A
MS, LA, TN, AR, MO	0.75 – 1.33 (Preemergence Only)	+	0.75 – 1.33 to 3-12" cotton
TX, OK, NM	1.0 – 1.33	+	1.0– 1.33 to 3-12" cotton before August 1
NC, VA	1.0 – 1.33 (Preemergence Only)	+	1.0– 1.33 to 3-12" cotton

In sprinkler-irrigated areas, sprinkler irrigate after application with one-half to 1 inch of water (one-half inch on *coarse-textured soils* to 1 inch on *fine-textured soils*) to incorporate this product. In furrow-irrigated areas, apply this product, incorporate with a rolling cultivator or similar implement that provides uniform shallow incorporation (2 inches or less), and then irrigate. In non-irrigated areas, if at least one-half inch of rainfall does not occur within 10 days after application, cultivate with a rolling cultivator or similar implement that provides uniform shallow incorporation of this product.

For best control of yellow nutsedge and suppression of seedling johnsongrass, apply this product preplant incorporated, preemergence, or postemergence to cotton and preemergence to weeds at the maximum rate for the soil texture, whether applied alone or in combinations. Do not apply more than a total of 2 pts./A on *coarse soils* or 4 pts./A of this product on *medium* and *fine soils* during a growing season (do not make tandem applications of metolachlor and s-metolachlor herbicides). These treatments may be applied over previous application of herbicides.

RESTRICTIONS:

1. Do not graze or feed forage or fodder from cotton to livestock, or illegal residues may result.
2. To avoid crop injury, do not apply this product on sand or loamy sand soils, or in areas where water is likely to "pond" over the bed;
3. To avoid concentration in the seed furrow, do not make broadcast applications of this product to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow;
4. In furrow-planted cotton, to avoid concentration in the furrow and potential injury, do not apply this product postemergence until after first "knifing" or cultivation to level soil surface.
5. Do not apply over-the-top in fluid fertilizer or any other adjuvant, surfactant, oil, or other pesticide not specified in the cotton section of this label, or injury may occur;
6. Do not apply on Taloka silt loam.
7. Do not use in Gaines County, TX.

COTTON – STALWART C COMBINATIONS

TANK MIXTURE WITH PROMETRYN

Stalwart C tank mixed with prometryn may be applied preplant incorporated or preemergence in water or fluid fertilizer. When fluid fertilizer is used as a carrier for this product, either alone or in combination with prometryn, mix only the amount that will be sprayed in one operation. Do not allow these to stand without agitation. Only water may be used as a carrier for postemergence-directed application.

In addition to those weeds controlled by Stalwart C alone, Stalwart C + prometryn, applied preplant incorporated or preemergence, also controls the following weeds: junglerice, wild oats, annual morningglory, groundcherry, hairy nightshade, lambsquarters, malva, mustard, prickly sida (teaweed), purslane, ragweed, and shallow-germinating seedlings of cocklebur and coffeeweed. As a postemergence-directed application, prometryn provides postemergence control and residual control of weeds on its label, while this product provides residual control of weed species on its label. This product will not control emerged weeds.

Preplant Incorporated or Preemergence: Apply Stalwart C + prometryn, either preplant incorporated or preemergence, using the appropriate rate from the table below. Cotton should be planted below the zone of incorporation; i.e., at least 1 inch on *fine soils* and 1.5 inches on *coarse* and *medium soils*. If incorporated before planting, use a planter that will result in a minimum of soil disturbance.

TABLE 2: STALWART C + PROMETRYN – COTTON (NM, OK, TX)

Use Areas	Soil Texture	Broadcast Rates Per Acre	
		Stalwart C	Prometryn
ALL	Sand, Loamy sand	DO NOT USE	
OK, and Blacklands, Gulf Coast, and Rio Grande Valley of TX	Loams	0.85-1.33 pts.	Label rate
	Clays	1.33 pts.	Label rate
NM; High Plains, Rolling Plains, Edwards Plateau of TX; and Southwest TX	Sandy loam	0.85-1.0 pt.	Label rate
	Loams	0.85-1.33 pts.	Label rate
	Sandy clay loams	1.33 pts.	Label rate
	Other clay soils	1.33 pts.	Label rate

Postemergence-Directed (AR, AZ, CA, LA, MO, MS, NM, OK, TN, and TX): Stalwart C may be tank mixed with prometryn in water and applied postemergence directed in cotton for control of emerged weeds listed on the prometryn label and residual preemergence control of weeds controlled by this product and prometryn, or application may be made after cultivation for residual preemergence control. These treatments may be applied over previous application of herbicides, including this product, provided the maximum label rate of any product is not exceeded (do not make tandem applications of metolachlor and s-metolachlor herbicides). Do not apply over-the-top of cotton or injury may occur.

Apply Stalwart C + prometryn in a minimum of 20 gals. of spray volume per acre. Follow the directions, restrictions, and use precautions on the prometryn label when prometryn is applied as a postemergence-directed application. Refer to the directions, restrictions, and use precautions for this product under the **Cotton — Stalwart C Alone — Postemergence** section.

RESTRICTIONS:

1. To avoid concentration in the seed furrow, do not make broadcast applications of this product + prometryn to cotton planted in furrows more than 2 inches deep. Band applications may be

made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow.

2. To avoid crop injury,
 - a. Do not apply on sand or loamy sand soils, or in areas where water is likely to “pond” over the bed;
 - b. Do not apply in cut areas of newly leveled fields, or in areas of excess salt;
 - c. Do not apply to glandless cotton varieties; and
 - d. Do not apply on Taloka silt loam.
3. Do not use in Gaines County, TX.
4. Do not graze or feed forage or fodder from cotton to livestock, or illegal residues may result.
5. Refer to the prometryn label for further instructions and restrictions.

TANK MIXTURE WITH FLUMETURON

Stalwart C may be applied in tank mixture with flumeturon preemergence for control of those weeds controlled by this product alone and those as listed on the flumeturon label. This combination will also control spotted spurge, hyssop spurge, nodding spurge, and prostrate spurge. Apply to the soil surface at planting or after planting, but before weeds or crop emerges, using the appropriate rates from the table below. The tank mixture may be applied postemergence to cotton, but preemergence to weeds, or it may be applied postemergence to both cotton and broadleaf weeds for control of weeds on the flumeturon label. Apply as a directed, semi-directed, or over-the-top spray. This product will not control emerged weeds, but will provide preemergence control of species on its label.

Mixing Instructions: Incompatibility may occur when tank mixing Stalwart C and flumeturon. To help overcome this condition, fill the spray tank one-quarter full with water or fluid fertilizer and start agitation, add the flumeturon and allow it to become dispersed. Add X-77® at 0.5% volume/volume final spray (4 pts./100 gals.), then add this product and finally the rest of the water or fluid fertilizer. Agitate during mixing and application to maintain a uniform suspension. Do not use fluid fertilizer as a carrier for postemergence applications.

TABLE 3: STALWART C + FLUMETURON – COTTON

Soil Texture	Broadcast Rates Per Acre		
	Stalwart C (pts.)		Flumeturon
	Area 1*	Area 2**	
Sand, Loamy sand	DO NOT USE		
Sandy loam	0.75-1.0	0.85-1.0	Label rate
Loam, Silt loam, Silt	1.0-1.33	1.0-1.33	Label rate
Fine soil	1.0-1.33	1.33	Label rate

* Area 1 = AR, LA, MS, Bootheel of MO and TN

** Area 2 = Eastern OK, Gulf Coast, Rio Grande Valley, and Eastern TX

Postemergence: This tank mixture may be applied postemergence to cotton, but preemergence to weeds or postemergence to both cotton and weeds for control of weeds on the flumeturon label. Apply as a directed, semi-directed, or over-the-top spray. This product will not control emerged weeds, but will provide preemergence control of species on its label. Apply when cotton is in the 3- to 12-inch stage. Where rate ranges are given for flumeturon, use the higher rate when applying postemergence to weeds that are 2 inches or less. These treatments may be applied over previous application of herbicides, including this product, provided the maximum label rate of any product is not exceeded (do not make tandem applications of metolachlor and s-metolachlor herbicides).

RESTRICTIONS

1. To avoid possible illegal residues, DO NOT feed treated forage or gin trash to livestock, or graze treated areas.

2. DO NOT apply Stalwart C + flumeturon on sand or loamy sand soils, or in areas where water is likely to “pond” over the bed, or crop injury may occur.
3. To avoid concentration in the seed furrow, do not make broadcast applications of this product + flumeturon to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow.
4. The use of flumeturon following the use of a systemic insecticide at planting may result in crop injury.
5. DO NOT use on Taloka silt loam, or crop injury may occur.
6. DO NOT use in Gaines County, TX.

Refer to the flumeturon labels for further instructions, use precautions, and restrictions.

TANK MIXTURE OF STALWART C + FLUMETURON WITH PARAQUAT OR GLYPHOSATE FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where cotton is planted directly into a cover crop, stale seedbed, or previous crop residues, the contact herbicides paraquat or glyphosate may be added to a tank mix of either Stalwart C or Stalwart C + flumeturon. When used as directed, the paraquat portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Glyphosate combinations will control emerged annual and perennial weeds when applied as directed on the glyphosate label. Stalwart C and Stalwart C + flumeturon portion of the tank mixture provides preemergence control of the weeds listed on each label, respectively.

Refer to the label of each product used in combination and observe the planting details, information regarding application, geographical restrictions, and all other precautions and restrictions. Refer to **Mixing Instructions** under **Tank Mixture with Flumeturon** section.

Application: Apply before, during, or after planting, but before the cotton emerges, at the rates specified below. Apply this product at 0.85 to 1 pt./A on sandy loams, *medium-*, and *fine-textured soils*. Use flumeturon at the labeled rates.

Do not apply this product + flumeturon + glyphosate in tank mixture because of compatibility problems.

Add paraquat or glyphosate at the following broadcast rates:

Paraquat: Use the labeled rates. Apply surfactant at 1 or 2 pts./100 gals. of spray mixture with 75% or greater or 50 to 74% nonionic active ingredient, respectively. This treatment will not control weeds taller than 6 inches.

RESTRICTION: Do not apply combinations containing paraquat in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

Glyphosate: See the glyphosate label for weeds controlled, specified rates, and other use directions.

Apply in 20 to 60 gals. of water or fluid fertilizer per acre with ground equipment.

RESTRICTIONS:

1. If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed.
2. Refer to the flumeturon labels and the **Tank Mixture with Flumeturon** section of this label for further instructions, use precautions, and restrictions.
3. Do not use in Gaines County, TX.

TANK MIXTURE WITH MSMA, MSMA + PROMETRYN, OR MSMA + FLUMETURON

Stalwart C may be tank mixed with MSMA in water and applied postemergence-directed for control of emerged weeds listed on the MSMA product label and residual preemergence control of weeds controlled by this product. The addition of prometryn or flumeturon will add control of weed species on their respective labels.

Postemergence-Directed (AR, AZ, CA, LA, MS, NM, OK, TN, TX, and Bootheel of MO): Apply Stalwart C + MSMA postemergence-directed to 3 to 12-inch cotton according to the directions, restrictions, and use precautions on the MSMA product label, as well as the directions, restrictions, and use precautions for use of this product in the section for **Cotton — Stalwart C Alone – Postemergence**. Do not apply after first cotton bloom. These treatments may be applied over previous registered treatments, including this product, provided the maximum label rate of any product is not exceeded. Flumeturon or prometryn may be added to this product + MSMA tank mixture according to the respective label directions for application to 3- to 12-inch cotton. When these mixtures are used, follow the mixing instructions for Stalwart C + prometryn or flumeturon and then add the MSMA product.

Do not use this product in tank mix with premixes of MSMA plus herbicides other than those registered for use in tank mixture with this product on cotton.

TANK MIXTURE OF STALWART C WITH GLYPHOSATE FOR USE ON ROUNDUP READY® COTTON AND ROUNDUP READY FLEX COTTON ONLY

Stalwart C may be tank mixed with glyphosate in water and applied postemergence over-the-top or postemergence-directed spray only to RoundUp Ready or RoundUp Ready Flex cotton or other cotton varieties or cultivars warranted as tolerant to glyphosate. This tank mixture will control emerged weeds listed on the glyphosate label and residual preemergence control of weeds listed on this label. See the **Cotton – Stalwart C Alone – Postemergence** section for proper rates and timing of Stalwart C. Also follow the glyphosate label for appropriate use rate, method of application, and restrictions of application timing. For postemergence over-the-top application, do not add any adjuvants, surfactants, fertilizers, or other pesticides to this tank mixture as unacceptable injury may occur.

RESTRICTIONS:

1. Do not apply this tank mixture postemergence to any cotton variety unless it is designated glyphosate tolerant, and unless the glyphosate formulation being used is registered for postemergence use in Roundup Ready Cotton or glyphosate tolerant cotton.
2. Postemergence over-the-top applications of this tank mixture may cause temporary injury in the form of necrotic spotting to exposed cotton leaves, which will not affect normal plant development.
3. Do not apply glyphosate postemergence over-the-top to cotton past the growth stage limit specified on the label.
4. Do not use on sand or loamy sand soils in Gaines County, TX.

TANK MIXTURE OF STALWART C WITH GLUFOSINATE TOLERANT COTTON

Stalwart C may be tank mixed with glufosinate in water and applied as a postemergence, broadcast over-the-top spray or as a postemergence-directed spray only to LibertyLink cotton or other cotton varieties or cultivars warranted as tolerant to glufosinate. This tank mixture will control emerged weeds listed on the glufosinate label and provide residual preemergence control of weeds listed on this label. See the **Cotton – Stalwart C Alone – Postemergence** section for proper rates and timing of Stalwart C. Also follow the glufosinate label for appropriate use rate, method of application, and restrictions of application timing. For postemergence over-the-top application, do not add any adjuvants, surfactants, fertilizers, or other pesticides to this tank mixture as unacceptable injury may occur.

RESTRICTIONS:

1. Do not apply this tank mixture postemergence to any cotton variety unless it is designated glufosinate tolerant, and unless the glufosinate formulation being used is registered for postemergence use in glufosinate tolerant cotton.
2. Postemergence over-the-top applications of this tank mixture may cause temporary injury in the form of necrotic spotting to exposed cotton leaves, which will not affect normal plant development.
3. Do not apply glufosinate postemergence to cotton beyond early bloom stage.
4. Do not use on sand or loamy sand soils in Gaines County, TX.

PEANUTS – STALWART C ALONE

Apply Stalwart C, either preplant incorporated, postplant incorporated, preemergence, or lay-by, using the appropriate rate specified below. **Preplant Incorporated or Preemergence:** Follow instructions for use of this product alone under **Application Procedures**.

Postplant Incorporated: Apply and shallowly incorporate this product into the soil after planting, but before peanut germination. Incorporation depth and incorporating implements must be kept above the seed, or seed will be damaged.

Lay-by: Apply Stalwart C to the soil immediately after the last normal cultivation. Apply this product alone, preplant incorporated, postplant incorporated, or pre-emergence, or lay-by, at a broadcast rate of 1 to 1.33 pts./A in the Southeast* and 0.85 to 1.33 pts./A in NM, OK, and TX.

*In the Southeast, use 1.33 to 2 pts./A and apply preemergence for partial control of Florida beggarweed.

RESTRICTIONS:

1. Do not apply within 90 days of harvest, or illegal residues may result.
2. Do not graze or feed peanut forage or fodder to livestock for 30 days following application.
3. Stalwart C alone may be applied as directed after any of the following preplant incorporated herbicides when used according to their label: trifluralin at the labeled rate; ethafluralin at the labeled rate; imazethapyr at the labeled rate; or pendimethalin at the labeled rate.

PEANUTS – STALWART C COMBINATIONS

TANK MIXTURE OR SEQUENTIALLY WITH IMAZETHAPYR

The tank mixture or sequential treatment of Stalwart C and imazethapyr controls both all weeds controlled by this product alone and all weeds controlled by imazethapyr alone. Refer to the **Stalwart C Applied Alone** section for weeds controlled by this product and to the imazethapyr label for weeds controlled by imazethapyr.

Refer to the respective labels for application methods, timing, rates, restrictions, and use precautions; and use in accordance with the most restrictive label. Do not exceed the label rate of either product. This product will not control emerged weeds.

TANK MIXTURE WITH ETHAFLURALIN

The tank mixture controls all weeds controlled by Stalwart C alone and by ethafluralin alone. Refer to the **Stalwart C Applied Alone** section for weeds controlled by this product and to the ethafluralin label for weeds controlled by ethafluralin.

Apply Stalwart C + ethafluralin preplant incorporated, using the appropriate rate from the table below. Follow label recommended soil preparation and soil-incorporation procedures for ethafluralin.

Table 4: Stalwart C + Ethafluralin – Peanuts

Soil Texture	Broadcast Rates Per Acre (pts.)			
	Southeast		NM, OK, TX	
	Stalwart C	Ethafluralin	Stalwart C	Ethafluralin
COARSE	1.0-1.33	Label rate	0.85-1.33	Label rate
MEDIUM	1.0-1.33	Label rate	0.85-1.33	Label rate
FINE	1.0-1.33	Label rate	0.85-1.33	Label rate

Follow all use directions, restrictions, and use precautions regarding application to peanuts on this product and ethafluralin labels.

TANK MIXTURE WITH PENDIMETHALIN

Stalwart C + pendimethalin applied preplant incorporated controls all weeds controlled by this product alone plus Texas panicum, field sandbur, johnsongrass from seed, lambsquarters, kochia, annual spurge, and other species on the pendimethalin label. Apply Stalwart C + pendimethalin by ground or by aerial equipment within 14 days before planting. Incorporate into the top 1 to 2 inches of soil before planting and within 7 days of application, using a finishing disk or similar implement capable of providing uniform incorporation. If peanuts will be planted on beds, apply and incorporate after bed formation. Refer to the **Incorporation** instructions of the respective labels for additional directions.

Apply Stalwart C + pendimethalin preplant incorporated, using the appropriate rates from the table below.

Table 5: Stalwart C + pendimethalin – Peanuts

Soil Texture	Broadcast Rates of Stalwart C (pints per acre)			
	NM, OK, TX		Other Peanut Growing States	
	Stalwart C	Pendimethalin	Stalwart C	Pendimethalin
Sand, Loamy sand	0.85	Label rate	1.0-1.33	Label rate
Sandy loam	0.85-1.0	Label rate	1.0-1.33	Label rate
Fine soil	1.33	Label rate	1.33	Label rate

Follow all use directions, restrictions, and use precautions regarding application to peanuts on Stalwart C and pendimethalin labels.

TANK MIXTURE OR SEQUENTIALLY WITH BENTAZON/ACIFLUFEN

Stalwart C + bentazon/aciflufen applied at ground cracking through 2 expanded tetrafoliate leaves or Stalwart C applied according to the directions for **Stalwart C Alone** and followed with an at-cracking through postemergence treatment of bentazon/aciflufen as specified on its label will control species on the bentazon/aciflufen label and provide residual control of species listed in the **Stalwart C Applied Alone** section of this label. This product will not control emerged weeds. Refer to the **Peanuts – Stalwart C Alone** section and to the bentazon/aciflufen label and follow all directions, use precautions, and restrictions for each product.

**CROP GROUP 6 LEGUME VEGETABLES (succulent and dried) –
 STALWART C ALONE**

Crop Group 6 – Legume Vegetables (Succulent or Dried) Group – Beans, peas and lentils (includes grain lupin, sweet lupin, white lupin, and white sweet lupin, field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean, adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean, Broad bean (fava bean), Chickpea (garbanzo bean), Guar,

Jackbean, Lablab bean, Lentil, dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea, Pigeon pea, and Sword bean.)

*On English peas, use only preemergence applications. Do not use on English peas in Northeastern U.S., or injury may occur.

Spring Application: Apply Stalwart C either preplant incorporated or preemergence, using the appropriate rate specified below.

Preplant Incorporated or Preemergence: Follow instructions for use of this product alone under **Application Procedures**. On *coarse soils* with less than 3% organic matter, apply 1 to 1.33 pts./A of Stalwart C or 1.33 pts./A if organic matter is 3% or greater. On *medium soils*, apply 1.33 to 1.67 pts./A of this product. On *fine soils*, apply 1.33 to 1.67 pts./A of this product if organic matter content is less than 3%, or 1.67 to 2 pts./A if organic matter content is 3% or greater.

RESTRICTIONS: To avoid possible illegal residues,

1. Do not cut for hay within 120 days following application of this product.
2. Do not use for forage within 60 days following application of this product.
3. Do not apply more than 3 pts./A of this product during any one crop year.

CROP GROUP 6 LEGUME VEGETABLES (succulent and dried) – STALWART C COMBINATIONS

TANK MIXTURE AND SEQUENTIAL APPLICATIONS WITH EPTC – BEANS (GREEN OR DRY)

This mixture controls all weeds controlled by Stalwart C alone and by EPTC alone. Refer to the **Stalwart C Applied Alone** section of this label for weeds controlled by this product alone and to the EPTC label for weeds controlled by EPTC.

Preplant Incorporated: Follow instructions for use of this product alone under **Application Procedures**.

Sequential: Apply EPTC alone preplant incorporated, as specified on that label. Follow with a preemergence application of this product at rates specified for this product alone, during planting (behind the planter) or after planting, but before the weeds or crop emerge.

CALIFORNIA ONLY

Preplant Incorporated: See the **SPECIAL APPLICATION PROCEDURES** Section of this label.

Refer to the **Product Information** section of this label and to the EPTC label for weather, cultural practices, and all other use precautions and restrictions that affect performance of these products.

Apply the labeled rate of EPTC* with Stalwart C as specified. On *coarse soils*, apply 0.85 pts./A of this product if organic matter content is less than 3%, or 1 pt./A if organic matter content is 3% or greater. On *medium soils*, apply 1 pt./A of Stalwart C if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On *fine soils*, apply 1.33 pts./A of Stalwart C if organic matter is less than 3%, or 1.33 to 1.67 pts./A if organic matter is 3% or greater.

*Refer to the EPTC label for rate limitations depending on geographical area, and for species and varietal restrictions.

RESTRICTIONS:

1. DO NOT exceed the label rate of EPTC on small white beans or green beans grown on coarse-textured soils.
2. DO NOT cut for hay within 120 days following application, or illegal residues may result.

TANK MIXTURE WITH TRIFLURALIN – BEANS (DRY – KIDNEY, NAVY, PINTO, ETC.; LIMA; AND SNAP)

Stalwart C + trifluralin tank mix applied preplant incorporated controls those weeds listed under **Stalwart C Applied Alone** and those weeds listed for trifluralin alone on the trifluralin label. Stalwart C + trifluralin may be applied by ground or by aerial equipment and incorporated up to 14 days prior to planting. Follow the procedures on this label and on the respective trifluralin label, using equipment that provides uniform 2-inch incorporation.

Apply Stalwart C + trifluralin tank mix, using the appropriate label rate of this product, and the trifluralin specified label rate. Choose the product rate for the specific soil texture/organic matter classification and weed species expected.

Follow all restrictions and use precautions on the respective trifluralin label and in the Crop Group 6 Legume Vegetables (succulent and dried) – **Stalwart C Alone** section of this label.

POTATOES – STALWART C ALONE

Apply Stalwart C, either incorporated, preemergence, or after hilling/lay-by, according to directions specified below for control of weeds listed under the **Product Information** section. Within a rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil.

Incorporated: Apply Stalwart C at 1 to 2 pts./A to the soil and incorporate into the top 3 inches before planting, using a finishing disk, harrow, rolling cultivator, or similar implement. Planting and later cultural practices should not bring untreated soil to the surface. Postplant incorporated application may be made any time after planting to drag-off, but before potato emergence. Use an implement that evenly distributes this product in the top 2 inches of soil. Do not damage potato seed pieces or sprouts with incorporation equipment.

Preemergence: Apply Stalwart C at 1 to 2 pts./A, either after planting as a preemergence, delayed preemergence, after drag-off or hilling treatment, but before weeds emerge. Up to 2.75 pts./A of this product alone may be used where soil organic matter is between 6% and 20%.

After Hilling/Lay-by: Apply 1.67 pts./A of Stalwart C after hilling/at lay-by to control species sensitive to this product for remainder of the growing season. This application will not control emerged weeds. It may be applied over a previous application of this product, but do not apply more than 3.7 pts./A of this product in a single crop season.

RESTRICTIONS:

- (1) Potatoes treated with this product must not be harvested within 60 days after the at-planting to drag-off application, or within 40 days after a lay-by application, or illegal residues may result.
- (2) Do not use on muck or peat soils. If cool, wet soil conditions occur after application, this product may delay maturity and/or reduce yield of Superior and other early maturing potato varieties.
- (3) To avoid crop injury,
 - a. Do not use on sweet potatoes or yams;
 - b. Do not apply both as a preemergence and an incorporated treatment;
- (4) Do not use in Kern County, CA.

POTATOES – STALWART C COMBINATIONS

TANK MIXTURE WITH METRIBUZIN

In addition to those weeds controlled by Stalwart C alone, this product applied in tank mix combination with, or sequentially with, any of the registered metribuzin formulations, also controls the following broadleaf weeds: cocklebur*, hairy nightshade*, hemp sesbania, jimsonweed*, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, venice mallow, and wild mustard.

*Partially controlled.

Stalwart C at 1 to 2 pts./A plus metribuzin at the labeled use rate may be used preemergence through after last hilling. Apply 1 to 1.33 pts./A of Stalwart C on *coarse soils* and 1.33 to 2 pts./A on other soil textures. Within this rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. This product will not control emerged weeds.

Refer to the metribuzin label for precautionary statements, restrictions, application information, weeds controlled, and varietal limitations.

RESTRICTIONS:

- (1) Potatoes treated with this product in tank mixture with metribuzin cannot be harvested within 60 days after application, or illegal residues may result.
- (2) Potatoes must not be harvested within 40 days after a lay-by application of this product, or illegal residues may result.
- (3) DO NOT apply to sweet potatoes or yams.
- (4) DO NOT use this tank mixture on muck or peat soils.
- (5) DO NOT use this product + metribuzin on potatoes in Kern County, CA
- (6) Postemergence applications to potatoes should be made only as a directed or semi-directed spray to avoid chlorosis, minor necrosis, or leaf distortion.

STALWART C + LINURON TANK MIXTURE (EAST OF ROCKY MOUNTAINS)

Stalwart C may be applied in a tank-mix combination with any of the registered linuron formulations as a preemergence broadcast application to potatoes. Apply to the soil surface after planting and before emergence of the crop or after final drag-off, according to the rates specified in the table below.

TABLE 6: STALWART C + LINURON – POTATOES (EAST OF ROCKY MOUNTAINS)

Soil Texture	Broadcast Rates Per Acre			
	1% to Less Than 3% Organic Matter		3-5% Organic Matter	
	Stalwart C	Linuron	Stalwart C	Linuron
COARSE Sandy loam	1 pt.	Label rate	1.33 pts.	Label rate
MEDIUM Loam, Silt loam, Silt	1.33 pts.	Label rate	1.67-2 pts.	Label rate

RESTRICTIONS:

To avoid crop injury,

1. Do not use on sands or loamy sands, and
2. Do not incorporate or spray over the top of emerged potatoes.

Refer to the **Product Information** section of this label and to the linuron label for precautionary statements, restrictions, application information, and weeds controlled.

TANK MIXTURE WITH PENDIMETHALIN

In addition to the weeds controlled by Stalwart C alone, this tank mixture with pendimethalin controls such problem species as kochia, lambsquarters, purslane, annual spurge, stinging nettle, and others specified on the pendimethalin label. Apply Stalwart C + pendimethalin preemergence, preemergence incorporated or early postemergence according to the specific directions on the pendimethalin label, using the rates in the table below.

TABLE 7: STALWART C + PENDIMETHALIN – POTATOES

Soil Texture	Broadcast Rates Per Acre (pts.)	
	Less Than 3% Organic Matter	More Than 3% Organic Matter
	Stalwart C + pendimethalin	Stalwart C + pendimethalin
COARSE	1.0-1.33 + label rate	1.0-1.33 + label rate
MEDIUM	1.33 + label rate	1.33-1.67 + label rate
FINE	1.33-1.67 + label rate	1.67 + label rate

Refer to Stalwart C and pendimethalin labels and observe all directions, timings, use precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.

TANK MIXTURE WITH PENDIMETHALIN + EPTC

In addition to the weeds controlled by Stalwart C alone, this tank mixture will control those species on the pendimethalin and EPTC labels. Refer to the Stalwart C, pendimethalin and EPTC labels for rates of those products (depending on geographical area); and observe all directions, use precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.

SAFFLOWERS – STALWART C ALONE

Preplant Incorporated or Preemergence: Follow instructions for use of Stalwart C alone under **Application Procedures**.

On *coarse soils*, apply 1 to 1.33 pts./A of Stalwart C if organic matter content is less than 3%, or 1.33 pts./A if organic matter is 3% or greater. On *medium soils*, apply 1.33 to 1.67 pts./A of this product. On *fine soils*, apply 1.33 to 1.67 pts./A of this product if organic matter content is less than 3%, or 1.67 to 2 pts./A if organic matter content is 3% or greater.

CALIFORNIA ONLY

Preplant Incorporated: See the **SPECIAL APPLICATION PROCEDURES** Section of this label.

SORGHUM – STALWART C ALONE

USE ONLY ON SORGHUM (GRAIN OR FORAGE) SEED TREATED WITH CONCEP® OR SCREEN®

Apply Stalwart C, either preplant surface, preplant incorporated, or preemergence, using the appropriate rate specified below. Apply this product alone only when the sorghum seed has been properly treated by the seed company with Concep or Screen.

Pre-plant Surface Applied: Refer to instructions for this product under **Application Procedures**. For minimum-tillage or no-tillage systems only, Stalwart C may be applied up to 45 days before planting in CO, IA, IL, KS, MO, NE, and SD. Use only split applications for treatments made 30 to 45 days prior to planting, with two-thirds of the broadcast rate applied initially and the remaining one-third at planting. Apply 1.5 pts./A of this product on *medium soils* or 1.67 pts./A on *fine soils*. Treatments less than 30 days prior to planting may be either as a split or single application. Apply 1.33 pts./A of this

product on *coarse soils* not more than 2 weeks prior to planting. Under dry conditions, irrigation after application is recommended to move this product into the soil.

Preplant Incorporated or Preemergence: Refer to instructions for use of Stalwart C under **Application Procedures**. Broadcast 1 to 1.33 pts./A of this product on *coarse soils*, 1.33 to 1.5 pts./A on *medium soils*, or 1.33 to 1.67 pts./A on *fine soils*.

RESTRICTIONS:

1. If sorghum seed is not properly treated with Concep or Screen seed treatment, this product will severely injure the crop.
2. Under high soil moisture conditions prior to sorghum emergence, injury may occur following the use of this product. The crop will normally outgrow this effect.
3. DO NOT use this product on sorghum grown under dry mulch tillage, or injury may occur.
4. Except for the split preplant surface treatment, DO NOT make more than one application per year, or illegal residues may result.

SORGHUM – STALWART C COMBINATIONS

USE ONLY ON SORGHUM (GRAIN OR FORAGE) SEED TREATED WITH CONCEPT[®] OR SCREEN[®]

Stalwart C tank mixtures with atrazine may be applied in water or fluid fertilizer. Apply this product in tank mixtures only when the sorghum seed has been properly treated by the seed company with Concep or Screen.

IMPORTANT: FOR TANK MIXTURES WITH ATRAZINE – If applying this product in tank mixture with atrazine, all the restrictions and rate limitations on the atrazine label must be followed if more restrictive than those on this label. In addition, if atrazine is/must be applied at rates lower than the label rate, broadleaf weed control may be affected. Refer to the atrazine label for weeds controlled at the reduced rates.

RESTRICTIONS:

1. If sorghum seed is not properly treated with Concep or Screen seed treatment, this product + atrazine may severely injure the crop.
2. Applications of this product + atrazine on highly alkaline soils or on eroded areas where calcareous subsoils are exposed may cause sorghum injury.
3. Under high soil moisture conditions prior to sorghum emergence, injury may occur following the use of this product + atrazine. The crop will normally outgrow this effect.
4. DO NOT use this product + atrazine on sorghum grown under dry mulch tillage, or injury may occur.
5. Except for the split preplant surface treatment, DO NOT make more than one application per year, or illegal residues may result.

TANK MIXTURE WITH ATRAZINE

In addition to the weeds controlled by Stalwart C alone, Stalwart C + atrazine also controls the following broadleaf weeds when applied either preplant surface, preplant incorporated, or preemergence: cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

Preplant Surface-Applied: Refer to instructions for use of Stalwart C under **Application Procedures**. For minimum-tillage or no-tillage systems only, Stalwart C + atrazine may be applied up to 45 days prior to planting in IA, IL, Eastern KS, MO, NE, and SD. Use only split applications for

treatments made 30 to 45 days prior to planting, with two-thirds of the broadcast rate applied initially and the remaining one-third at planting. Apply 1.5 pts./A of Stalwart C + the label rate of atrazine on *medium soils* with 1.5% organic matter or greater. Apply 1.5 pts./A of Stalwart C + the label rate of atrazine on *fine soils* with less than 1.5% organic matter, or apply 1.67 pts./A of Stalwart C + the label rate of atrazine on *fine soils* with 1.5% organic matter or greater. Treatments less than 30 days prior to planting may be made either as a split or single application. Under dry conditions, irrigation after application is recommended to move Stalwart C + atrazine into the soil.

RESTRICTIONS:

To avoid crop injury,

1. Do not use on coarse soils,
2. Do not use on medium soils with less than 1.5% organic matter.

Preplant Incorporated or Preemergence: Refer to instructions for use of Stalwart C under **Application Procedures**. On *medium soils* with 1.5% organic matter or greater, apply 1 pt./A of Stalwart C + the label rate of atrazine. On *fine soils* with less than 1.5% organic matter, apply 1 pt./A of Stalwart C + the label rate of atrazine. On *fine soils* with 1.5% organic matter or greater, apply 1.2-1.33 pts./A of Stalwart C + the label rate of atrazine.

RESTRICTIONS:

To avoid crop injury,

1. DO NOT use on coarse soils;
2. DO NOT use on medium soils with less than 1.5% organic matter;
3. DO NOT use in NM, OK, or TX, except in Northeast OK and the TX Gulf Coast and Blacklands areas;
4. DO NOT apply preplant incorporated in AZ or the Imperial Valley of CA.

MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

TANK MIXTURE OF STALWART C OR STALWART C + ATRAZINE WITH PARAQUAT, GLYPHOSATE/2, 4-D OR GLYPHOSATE

In minimum-tillage or no-tillage systems where sorghum (seed treated with Concep or Screen) is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides paraquat, glyphosate/2, 4-D or glyphosate may be tank mixed with Stalwart C or Stalwart C + atrazine. Mix with paraquat for control of most emerged annual weeds and suppression of perennial weeds; or with glyphosate/2, 4-D for suppression of emerged field bindweed and control or suppression of annual weeds; or with glyphosate for control of most emerged annual and perennial weeds. Stalwart C or Stalwart C plus atrazine portion of the tank mixture provides preemergence control of the weeds listed on this label under the respective sections.

Refer to the label of each product used in combination and observe the planting details, restrictions, and all other use precautions.

Application: Apply before, during, or after planting, but before sorghum emerges, at the appropriate rates listed under **Sorghum (Grain or Forage) – Stalwart C Alone** or **– Stalwart C in Combinations – Tank Mixture with atrazine**, respectively. Add paraquat, glyphosate/2, 4-D, or glyphosate at the following broadcast rates:

Paraquat: Use the labeled rate. Apply surfactant at 1 or 2 pts./100 gals. of spray mixture with 75% or greater or 50 to 74% nonionic active ingredient, respectively. This treatment will not control weeds taller than 6 inches.

Glyphosate/2, 4-D: Use the labeled rate depending on weed species and size. See the glyphosate/2, 4-D label for weeds controlled, rates for specific weeds, and other information concerning use.

Glyphosate: See the glyphosate label for weeds controlled, rates, and other use directions. Apply in a minimum of 20 gals. of water per acre with conventional spray equipment.

SOYBEANS – STALWART C ALONE

Apply Stalwart C, either preplant surface-applied, preplant incorporated, preemergence, or post emergence using the appropriate rate specified below.

Preplant Surface-Applied, Preplant Incorporated, Preemergence, or Postemergence: Follow instructions for use of this product alone under **Application Procedures**.

Preplant Surface-Applied

Fall Application

- **Apply after September 30 in MN, ND, SD, WI, and North of Route 30 in IA**
- **Apply after October 15 North of Route 91 in NE and South of Route 30 in IA**
- **Apply after October 31 North of Route 136 in IL**

In all locations, apply Stalwart C to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 to 2 pts./A on *medium-textured* and 2 pts./A on *fine-textured soils*. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2 to 3 inches. Minimize furrow and ridge formation in the tillage operations.

Preplant Surface – Spring Application

Use on *medium* and *fine* soils with minimum-tillage or no-tillage systems in CO, CT, DE, IA, IL, IN, KS, KY, MA, MD, ME, MI, MN, MO, MT, ND, NE, NH, NY, OH, PA, RI, SD, TN, VA, VT, WI, WV, and WY. Apply two-thirds of the specified rate of Stalwart C (1.67 pts./A on *medium soils* and 2 pts./A on *fine soils*) as a split treatment 30 to 45 days prior to planting and the remainder at planting. Applications made less than 30 days before planting may be as either a split or single treatment. Apply 1.33 pts./A on *coarse soils* not more than 2 weeks before planting.

Preplant Incorporated or Preemergence: On *coarse soils*, apply 1 to 1.33 pts./A of Stalwart C if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On *medium soils*, apply 1.33 to 1.67 pts./A of this product. On *fine soils*, apply 1.33 to 1.67 pts./A of this product if organic matter content is less than 3%, or 1.67 to 2 pts./A if organic matter content is 3% or greater.

On soybeans, Stalwart C may be used up to 2.75 pts./A as a preplant surface-applied, preplant incorporated, or preemergence treatment on soils having an organic matter content between 6% and 20%. The total rate of this product applied to soybeans during any one crop season must not exceed 2.75 pts./A.

RESTRICTIONS: (Preplant surface applied, preplant incorporated, preemergence):

1. On soybeans, use up to 2.5 pints / acre of Stalwart C preplant incorporated or preemergence treatment on soils having an organic matter content between 6% and 20%.
2. The total Stalwart C rate applied to must not exceed 2.75 pints / acre per year.
3. If a pre-plant surface, pre-plant incorporated or pre-emergence application of metolachlor products has already been applied, a postemergence application of this product cannot be used.
4. DO NOT graze or feed treated soybean forage, hay, or straw to livestock 30 days following treatment, or illegal residues may result.
5. If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate on soybeans, or illegal residues may result.
6. DO NOT apply to frozen ground.

POSTEMERGENCE APPLICATIONS –

Postemergence (From emergence up through the 5th trifoliolate leaf stage)

For postemergence treatments, apply 1.0 to 1.33 pints per acre to soybeans from emergence through the 5th trifoliolate leaf stage. This product will not control emerged weeds, it must be applied to a weed free surface or in tank mixture with products that provide postemergence control of weeds present at the time of application.

Make postemergence applications at least 90 days before harvest.

RESTRICTIONS (Postemergence)

1. To prevent illegal residues, DO NOT apply more than 1.33 pints per acre postemergence.
2. DO NOT graze or feed treated soybean forage or soybean hay to livestock following a postemergence application of this product;
3. DO NOT apply a postemergence application of this product if a pre-plant surface, pre-plant incorporated or pre-emergence application of metolachlor products has already been applied.
4. **NOT FOR USE IN CALIFORNIA**

SOYBEANS – STALWART C COMBINATIONS

Preplant Incorporated or Pre-emergence

Water or fluid fertilizer may be used as carrier for Stalwart C in combination with metribuzin, linuron, metribuzin / chlorimuron-ethyl, imazethapyr, imazaquin, ethafluralin, or clomazone.

RESTRICTIONS:

1. For all of the following combinations, this product may be used up to 2.5 pts./A on soils having an organic matter content between 6% and 20%.
2. The total rate of this product applied to soybeans during any one crop year must not exceed 2.75 pts./A.
3. DO NOT apply to frozen ground.

TANK MIXTURE WITH METRIBUZIN

In addition to those weeds controlled by Stalwart C alone, Stalwart C + metribuzin, when applied as directed, also controls the following broadleaf weeds: cocklebur*, hairy nightshade, hemp sesbania, jimsonweed*, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, venice mallow, and wild mustard.

*Partially controlled.

Preplant Incorporated or Preemergence: Follow instructions for use of this product alone under **Application Procedures**.

Sequential: Apply Stalwart C alone **Preplant Incorporated**, as specified in the table below. Follow with a preemergence application of metribuzin during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

Refer to the metribuzin label for planting details and soybean variety restrictions.

TABLE 8: STALWART C + METRIBUZIN – SOYBEANS

Soil Texture*	Broadcast Rates Per Acre			
	0.5 to Less Than 3% Organic Matter		3% Organic Matter or Greater	
	Stalwart C	Metribuzin	Stalwart C	Metribuzin
COARSE	0.85-1.0 pt.	Label rate	1.0 pt.	Label rate
MEDIUM	1.0-1.33 pts.	Label rate	1.33 pts.	Label rate
FINE	1.33 pts.	Label rate	1.33-1.67 pts.	Label rate
Mississippi Delta only Silty clay, Clay	1.33 pts.	Label rate	1.33-1.67 pts.	Label rate
DO NOT USE ON MUCK OR PEAT (SOILS WITH MORE THAN 20% ORGANIC MATTER)				

* On all sand and on loamy sand with less than 2% organic matter, do not use this tank mixture preemergence or the sequential treatment. Do not use the tank mixture preplant incorporated on any sand, loamy sand, or sandy loam, or crop injury may occur.

RESTRICTION:

1. Do not use the tank mix or sequential application on soil with less than 0.5% organic matter or on alkaline soil with a pH over 7.4, or crop injury may occur.

PRECAUTION:

1. If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days.

TANK MIXTURE WITH LINURON

In addition to those weeds controlled by Stalwart C alone, Stalwart C + linuron, applied preemergence, also controls the following broadleaf weeds: cocklebur*, jimsonweed*, lambsquarters, morningglory*, prickly sida, ragweed, smartweed, velvetleaf*, venice mallow, and wild mustard.

*Partially controlled.

Preemergence: Apply during planting (behind planter) or after planting, but before weeds or soybeans emerge. Refer to the linuron label for planting details. Apply the appropriate rates from the table below.

RESTRICTIONS:

Do not use on soil with less than 0.5% organic matter, or crop injury may occur.

TABLE 9: STALWART C + LINURON – SOYBEANS

Soil Texture*	Broadcast Rates Per Acre			
	0.5 to Less Than 3% Organic Matter		3% Organic Matter or Greater	
	Stalwart C	Linuron	Stalwart C	Linuron
COARSE**	0.85 pt.	Label rate	1.0 pt.	Label rate
MEDIUM	1.0 pt.	Label rate	1.33 pts.	Label rate
FINE	1.33 pts.	Label rate	1.33-1.67 pts.	Label rate
DO NOT USE ON MUCK OR PEAT (SOILS WITH MORE THAN 20% ORGANIC MATTER)				

* Do not use on sand, gravelly soils, or exposed subsoils.

** Do not use on loamy sand, except in the Northeastern U.S. on loamy sand with over 1% organic matter.

TANK MIXTURE WITH TRIFLURALIN

Stalwart C + trifluralin tank mix applied preplant incorporated controls those weeds listed under the **Stalwart C Applied Alone** section and those weeds listed for trifluralin alone on the respective trifluralin label. Stalwart C + trifluralin may be applied by ground or by aerial equipment and incorporated up to 14 days before planting. Follow the procedures on the trifluralin and Stalwart C labels, using equipment that provides uniform 2-inch incorporation.

Apply Stalwart C + trifluralin tank mix, using the appropriate rate from the **Soybeans – Stalwart C Alone** section of this label and the appropriate section of the trifluralin label for the specific soil texture/organic matter classification and weed species expected.

To control DNA-resistant goosegrass* and other species on the respective labels where the soil organic matter is 3% or less, apply the rate in the table below.

TABLE 10: STALWART C + TRIFLURALIN – SOYBEANS (Organic Matter Content Less Than 3%)

Soil Texture	Broadcast Rates Per Acre (pts.)		
	Stalwart C	Trifluralin	
	Organic Matter Less Than 3%	Organic Matter	
		Less Than 2%	2-3%
COARSE*	0.85-1.0	Lower label rate	Higher label rate
MEDIUM	1.0	Lower label rate	Higher label rate
FINE	1.33	Lower label rate	Higher label rate

* Where a range of rates is given for this product use the minimum rate where DNA-resistant goosegrass is the predominant species.

Follow all restrictions and use precautions on the respective trifluralin label and in the **Soybeans – Stalwart C Alone** section of this label.

TANK MIXTURE WITH IMAZAQUIN

This tank mixture controls all weeds controlled by Stalwart C alone and by imazaquin alone. Refer to the Stalwart C Applied Alone section for weeds controlled by this product and to the imazaquin label for weeds controlled by imazaquin. Refer to the imazaquin label for geographical locations where this tank mixture may be applied.

Apply Stalwart C + imazaquin preplant incorporated or preemergence, using rates in the table below. Follow use directions under **Application Instructions** on the imazaquin label. For preplant incorporated applications, apply and incorporate within 30 days before planting. Observe all other use precautions and restrictions on the imazaquin labels.

TABLE 11: STALWART C + IMAZAQUIN – SOYBEANS

Soil Texture	Broadcast Rates Per Acre (pts.)			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	Stalwart C	Imazaquin	Stalwart C	Imazaquin
COARSE	0.85	Label rate	1.0	Label rate
MEDIUM	1.0	Label rate	1.33	Label rate
FINE	1.33	Label rate	1.33-1.67*	Label rate
DO NOT USE ON MUCK OR PEAT (SOILS WITH MORE THAN 20% ORGANIC MATTER)				

* Use the higher rate of this product if heavy weed infestations are expected.

RESTRICTIONS:

1. DO NOT apply within 90 days of harvest, and
2. DO NOT graze or feed treated soybean forage, hay, or straw to livestock, or illegal residues may result.

TANK MIXTURE WITH LINURON/CHLORIMURON-ETHYL

This tank mixture controls all weeds controlled by Stalwart C alone and by linuron / chlorimuron-ethyl alone. Refer to the Stalwart C Applied Alone section for weeds controlled by Stalwart C and to the linuron / chlorimuron-ethyl label for weeds controlled.

Apply Stalwart C + linuron / chlorimuron-ethyl preemergence after planting, but before soybeans emerge, using rates in the table below

Follow all use directions, restrictions, and use precautions regarding application to soybeans, and rotational restrictions on Stalwart C and linuron / chlorimuron-ethyl labels.

TABLE 12: STALWART C + LINURON / CHLORIMURON-ETHYL — SOYBEANS

Soil Texture	Broadcast Rates Per Acre	
	0.5 to 3% Organic Matter	
	Stalwart C	Linuron / Chlorimuron-ethyl
COARSE	0.85 pt.	Label rate
MEDIUM	1.0 pt.	Label rate
FINE	1.33 pts.	Label rate

RESTRICTION:

1. Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 6.8.

TANK MIXTURE WITH METRIBUZIN / CHLORIMURON-ETHYL

This tank mixture controls all weeds controlled by Stalwart C alone and by metribuzin / chlorimuron-ethyl alone. Refer to the **Stalwart C Applied Alone** section for weeds controlled by this product and to the metribuzin / chlorimuron-ethyl label for weeds controlled by metribuzin / chlorimuron-ethyl.

Apply preplant incorporated or preemergence, using the appropriate rates from the table below.

Preplant Incorporated: Apply within 2 weeks of planting. Uniformly incorporate into the top 1 to 2 inches of soil before planting soybeans.

Preemergence: Apply after planting, but before soybeans emerge.

Follow all use directions, restrictions, and use precautions regarding application to soybeans, and rotational restrictions on Stalwart C and metribuzin / chlorimuron-ethyl labels.

TABLE 13: STALWART C + METRIBUZIN / CHLORIMURON-ETHYL – SOYBEANS

Soil Texture	Broadcast Rates Per Acre (pts.)			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	Stalwart C	Metribuzin / Chlorimuron-ethyl	Stalwart C	Metribuzin / Chlorimuron-ethyl
COARSE	0.85 pt.	Label rate	1.0 pt.	Label rate
MEDIUM*	1.0 pt.	Label rate	1.33 pts.	Label rate
FINE*	1.33 pts.	Label rate	1.33-1.67 pts.	Label rate

*Refer to the metribuzin / chlorimuron-ethyl label for appropriate rate according to geographical location, soil and organic matter classification, and pH limitations.

RESTRICTION:

- Do not apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7, except as noted on the metribuzin / chlorimuron-ethyl label.

TANK MIXTURE WITH CLOMAZONE

This tank mixture controls all weeds controlled by Stalwart C alone and by clomazone alone. Refer to the **Stalwart C Applied Alone** section for weeds controlled by Stalwart C and to the clomazone label for weeds controlled by clomazone.

Apply Stalwart C + clomazone preplant incorporated, using rates in the table below. Follow all clomazone application instructions as to incorporation interval, geographical location, equipment operation, soil moisture conditions, etc.

Before making applications, read and strictly follow all use directions, restrictions, and use precautions regarding application to soybeans, and rotational restrictions on Stalwart C and clomazone labels.

TABLE 14: STALWART C + CLOMAZONE – SOYBEANS

Soil Texture	Broadcast Rates Per Acre (pts.)			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	Stalwart C	Clomazone	Stalwart C	Clomazone
COARSE	0.85 pt.	Label rate	1.0 pt.	Label rate
MEDIUM*	1.0 pt.	Label rate	1.33 pts.	Label rate
FINE*	1.33 pts.	Label rate	1.33-1.67 pts.	Label rate

TANK MIXTURE WITH ETHAFLURALIN

This tank mixture controls all weeds controlled by Stalwart C alone and by ethafluralin alone. Refer to the **Stalwart C Applied Alone** section for weeds controlled by Stalwart C and to the ethafluralin label for weeds controlled by ethafluralin.

Apply Stalwart C and ethafluralin preplant incorporated, using the appropriate rates from the table below.

Preplant Incorporated: Follow label recommended soil preparation and soil-incorporation procedures for ethafluralin.

Sequential: Apply ethafluralin alone preplant incorporated as specified on the ethafluralin label. Follow with a preemergence application of this product during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

TABLE 15: STALWART C + ETHAFLURALIN – SOYBEANS

Soil Texture	Broadcast Rates Per Acre (pts.)			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	Stalwart C	Ethafluralin	Stalwart C	Ethafluralin
COARSE	1.0-1.33	Label rate	1.33	Label rate
MEDIUM*	1.33-1.67	Label rate	1.33-1.67	Label rate
FINE*	1.33-1.67	Label rate	1.67-2.0	Label rate
DO NOT USE ON MUCK OR PEAT (SOILS WITH MORE THAN 20% ORGANIC MATTER)				

*For Eastern black nightshade on these soils, apply ethafluralin at 3.0 pts./A on *medium* and 3.5 pts./A on *fine-textured soils*, and follow with 2 incorporation passes.

Follow all use directions, restrictions, and use precautions regarding application to soybeans on the Stalwart C and ethafluralin labels.

TANK MIXTURE WITH IMAZETHAPYR

This tank mixture controls all weeds controlled by Stalwart C alone and by imazethapyr alone. Refer to the **Stalwart C Applied Alone** section for weeds controlled by Stalwart C and to the imazethapyr label for weeds controlled by imazethapyr. Refer to the imazethapyr label for geographical locations where this tank mixture may be applied.

Apply Stalwart C + imazethapyr early preplant, preplant incorporated, or preemergence after planting, using rates in the table below. Application can be made in water or liquid fertilizer.. For early preplant and preplant incorporated applications, apply within 30 days before planting.

Follow all use directions, restrictions, and, use precautions regarding application to soybeans, and rotational restrictions on the Stalwart C and imazethapyr labels.

TABLE 16: STALWART C + IMAZETHAPYR – SOYBEANS

Soil Texture	Broadcast Rates Per Acre (pts.)			
	Less Than 3% Organic Matter		3% or More Organic Matter	
	Stalwart C	Imazethapyr	Stalwart C	Imazethapyr
COARSE	0.85	Label rate	1.0	Label rate
MEDIUM	1.0	Label rate	1.33	Label rate
FINE	1.33	Label rate	1.33-1.67	Label rate

Sequential: Apply Stalwart C early preplant, preplant incorporated, or preemergence after planting at 0.85 pt./A on *coarse soils* and 1 pt./A on *medium- and fine-textured soils*. Follow with a sequential postemergence application of imazethapyr to control emerged weeds according to the imazethapyr label. Stalwart C will improve the consistency and level of control from imazethapyr on most grass species. Refer to the imazethapyr postemergence label for a listing of weeds controlled, application rate, and growth stage limitations.

MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

TANK MIXTURES WITH METRIBUZIN, IMAZAQUIN, LINURON, LINURON / CHLORIMURON-ETHYL, PRODIAMINE / ISOXABEN, METRIBUZIN / CHLORIMURON-ETHYL, OR IMAZETHAPYR, PLUS PARAQUAT OR GLYPHOSATE

In minimum-tillage or no-tillage systems where soybeans are planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides paraquat or glyphosate may be added to a tank mix of either Stalwart C + metribuzin, Stalwart C + imazaquin, Stalwart C + linuron, Stalwart C + linuron / chlorimuron-ethyl, Stalwart C + prodiamine / isoxaben, Stalwart C + metribuzin / chlorimuron-ethyl, or Stalwart C + imazethapyr.

When used as directed, the paraquat portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds.

Glyphosate combinations will control emerged annual and perennial weeds when applied as directed on the glyphosate label.

Stalwart C + metribuzin , imazaquin, linuron, linuron / chlorimuron-ethyl, prodiamine / isoxaben, metribuzin / chlorimuron-ethyl, or imazethapyr portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for Stalwart C + metribuzin , Stalwart C + imazaquin, Stalwart C + linuron, Stalwart C + linuron / chlorimuron-ethyl, Stalwart C + prodiamine / isoxaben, Stalwart C + metribuzin / chlorimuron-ethyl, and Stalwart C + imazethapyr, respectively.

Application: Apply before, during, or after planting, but before the soybeans emerge, at the rates specified in the chart below.

Apply in 20-60 gals. of water or fluid fertilizer per acre with ground equipment.

For all tank mix partners - Refer to the label of each product used in combination and observe the planting details, soybean variety restrictions, information regarding application to soybeans, geographical restrictions, soil and organic matter classification, pH limitations and all other use precautions and restrictions.

Paraquat: Use the labeled rate. Apply surfactant at 1 or 2 pts./100 gals. of spray mixture with 75% or greater or 50 to 74% nonionic active ingredient, respectively. This treatment will not control weeds taller than 6 inches.

RESTRICTION: Do not apply combinations containing paraquat in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

Tank Mix Partners	Stalwart C Rates Per Acre (pts.)			Tank Mix Partner	RESTRICTIONS
	Coarse Soil	Medium Soil	Fine Soil	Pints Per Acre	
Metribuzin + Paraquat Or Glyphosate	1.0	1.33	1.33 - 1.67	Labeled Rate	To avoid crop injury, Do not use this tank mixture on soil with less than 0.5% organic matter, on alkaline soil with a pH over 7.4, or on all sand and on loamy sand with less than 2% organic matter. If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed.
Imazaquin + Paraquat Or Glyphosate	1.0	1.33	1.67	Labeled Rate	Do not apply within 90 days of harvest, Do not graze or feed treated soybean forage, hay, or straw to livestock, or illegal residues may result.
Linuron + Paraquat Or Glyphosate	1.0	1.33	1.33 - 1.67	Labeled Rate	Do not use on loamy sand, except in the Northeastern U.S. on loamy sand with over 1% organic matter, or injury may occur. Do not use on sand, gravelly soils, or exposed subsoils, or injury may occur. Do not use on soil with less than 0.5% organic matter, or crop injury may occur.
Linuron / Chlorimuron-Ethyl + Paraquat Or Glyphosate	1.0	1.33	1.33 - 1.67	Labeled Rate	Use only where soils have 0.5 to 3% organic matter. Do not apply to sand or to any soil with pH greater than 6.8
Metribuzin / Chlorimuron-Ethyl + Paraquat Or Glyphosate	1.0 Do not use on sand.	1.33	1.33 - 1.67	Labeled Rate	Use only where soils have 0.5 to 5% organic matter. DO NOT apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 6.8, except as noted on the metribuzin / chlorimuron-ethyl label.
Imazethapyr + Paraquat Or Glyphosate	1.0	1.33	1.67		None

SOYBEANS – STALWART C COMBINATIONS POSTEMERGENCE

TANK MIXTURE OF STALWART C WITH GLYPHOSATE FOR USE ON GLYPHOSATE TOLERANT SOYBEANS

Stalwart C may be tank mixed with glyphosate in water and applied postemergence over-the top or postemergence-directed spray only up through the 5th trifoliolate leaf stage of soybean varieties or cultivars warranted as tolerant to glyphosate. This tank mixture will control emerged weeds listed on the glyphosate label and residual preemergence control of weeds listed on this label.

See the **Soybean – Stalwart C Alone – Postemergence** section for proper rates and timing of **Stalwart C**. Also follow the glyphosate label for appropriate use rate, method of application, and restrictions of application timing. For postemergence over-the-top application, do not add any adjuvants, surfactants, fertilizers, or other pesticides to this tank mixture as unacceptable injury may occur.

RESTRICTIONS:

- DO NOT apply this tank mixture postemergence to any soybean variety unless it is designated glyphosate tolerant and unless the glyphosate formulation being used is registered for postemergence use in Roundup Ready Soybeans or glyphosate tolerant soybeans.
- DO NOT apply more than 1.33 pints per acre postemergence.
- Do not apply a postemergence application of this product if a preplant surface, preplant incorporated, or preemergence application of metolachlor products has already been applied
- Following a postemergence application of this product, DO NOT graze or feed treated soybean forage or soybean hay to livestock.
- Postemergence applications must be made at least 90 days before harvest.
- **NOT FOR USE IN CALIFORNIA**

TANK MIXTURE OF STALWART C WITH GLUFOSINATE FOR USE ON GLUFOSINATE TOLERANT SOYBEANS

Stalwart C may be tank mixed with ~~glyphosate~~ glufosinate in water and applied postemergence over-the top or postemergence-directed spray only up through the 5th ~~3rd~~ trifoliolate leaf stage of soybean varieties or cultivars warranted as tolerant to glufosinate. This tank mixture will control emerged weeds listed on the glufosinate label and provide residual preemergence control of weeds listed on this label.

See the **Soybean – Stalwart C Alone – Postemergence** section for proper rates and timing of **Stalwart C**. Also follow the glufosinate label for appropriate use rate, method of application, and restrictions of application timing. For postemergence over-the-top application, do not add any adjuvants, surfactants, fertilizers, or other pesticides to this tank mixture as unacceptable injury may occur.

RESTRICTIONS:

- DO NOT apply this tank mixture postemergence to any soybean variety unless it is designated glufosinate tolerant and unless the glufosinate formulation being used is registered for postemergence use in Liberty Link Soybeans or glufosinate tolerant soybeans.
- DO NOT apply more than 1.33 pints per acre postemergence.
- Following a postemergence application of this product, DO NOT graze or feed treated soybean forage or soybean hay to livestock.
- Postemergence applications must be made at least 90 days before harvest.

TOMATOES – STALWART C ALONE

TRANSPLANTED TOMATOES

Stalwart C may be applied preplant incorporated or preplant before transplanting. If the latter method is used, keep soil disturbance to a minimum during transplanting. Application may also be made post-directed to transplants after the first settling rain or irrigation. When an application is made post-directed, apply in a minimum of 20 gallons of water per acre and minimum contact with tomato plants. This product will not control emerged weeds. In bedded transplanted tomatoes, apply this product preplant non-incorporated to the top of the pressed bed, as the last step, prior to laying plastic. **Stalwart C** may also be used to treat row-middles in bedded tomatoes, as long as the total amount of this product does not exceed the maximum allowed per crop.

SEEDED TOMATOES

Stalwart C may be applied post-directed to direct seeded tomatoes. Tomato plants must be at least 4 inches tall at the time of application and the product must be applied in a minimum of 20 gallons of water per acre. Minimize spray contact with tomato plants. This product will not control emerged weeds.

TOMATO USE RATES

Soil Texture	Broadcast Rates Per Acre (pts.)	
	Less Than 3% Organic Matter	3% or More Organic Matter
	Stalwart C	Stalwart C
COARSE	1.0 – 1.33 pints/acre	1.33 pints/acre
MEDIUM	1.33 pints/acre	1.33 pints/acre
FINE	1.33 – 1.67 pints/acre	1.67 – 2.0 pints/acre

RESTRICTIONS

To avoid possible illegal residues:

1. DO NOT apply this product within 90 days of tomato harvest.
2. DO NOT exceed the maximum label rate for the soil texture per year.
3. Apply only by ground application.

PRECAUTIONS

1. DO NOT apply to varieties or cultivars with unknown tolerance to Stalwart C.
2. This product may damage transplants that have been weakened by any cause. To prevent damage, plant only healthy transplants. Do not plant when wet, cool, or unfavorable growing conditions exist.
3. In transplanted tomatoes, if this product is applied preplant incorporated, incorporate to a depth less than the depth of transplanting, and use the lower end of the rate range for the given soil type, or damage may occur.
4. For row middle applications where tomatoes are grown on sandy soils and where high soil moisture conditions can exist (i.e. low binding and high evaporation conditions), as may be found in the States of Florida, Georgia, Maryland, and Virginia, there is potential for crop injury in the form of leaf epinasty. The risk of this type of injury can be reduced by:
 - a) incorporating this product immediately following application,
 - b) applying this product seven or more days before transplanting (but only after the beds have been formed),
 - c) minimizing the application of this product onto the plastic of the bed, or
 - d) any combination of the above.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: This product may be stored at temperatures down to -30°F. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

Pesticide Disposal: Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Open dumping is prohibited. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Container Handling:

[For Containers < 5 gallons:] Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, by incineration.

For Minibulk Containers [for nonrefillable containers > 5 gallons]: Nonrefillable container. Do not reuse or refill this container. Triple rinse container or pressure rinse promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration.

For Bulk Containers [for refillable containers > 5 gallons]: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When the container is empty, replace the cap and seal all openings that have been opened during use; and return to the point of purchase, or to a designated location named at the time of purchase of this product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged or leaking, call CHEMTREC. If the container is damaged and leaking or material has been spilled, follow these procedures:

- Cover spill with absorbent material.
- Sweep into disposal container.
- Wash area with detergent and water and follow with clean water rinse.
- Do not allow to contaminate water supplies.
- Dispose of according to instructions.

If not returned to the point of purchase or to a designated location, clean empty container as instructed above and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

WARRANTY AND LIMITATION OF DAMAGES

Conditions of Sale: To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. Sipcam Agro USA, Inc. disclaims all other warranties, express or implied. To the extent consistent with applicable law, Sipcam Agro USA, Inc. shall not be liable for consequential, special, or indirect damages resulting from the use or handling of this product, and Sipcam Agro USA, Inc.'s sole liability and buyer's and user's exclusive remedy shall be limited to the refund of the purchase price. Buyer and user acknowledge and assume all risks and liability resulting from handling, storage and use of this product. Sipcam Agro USA, Inc. does not authorize any agent or representative to make any other warranty, guarantee or representation concerning this product.

Stalwart is a registered trademark of Sipcam Agro USA, Inc.
LibertyLink® is a trademark of Bayer CropScience
Celatom MP-79 is a trademark of EP Minerals.
Unite is a registered trademark of HACO, Inc.
Compex is a registered trademark of KALO Agricultural Chemicals, Inc.
RoundupReady is a trademark of Monsanto Company
Formula 40 and Weedar are registered trademarks of Nufarm, Inc.
Agsorb is a registered trademark of Oil-Dri Corporation
X-77 is a registered trademark of Loveland Industries, Inc.