



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. Jon M. Gehring
Product Registration
Sipcam Agro USA, Inc.
300 Colonial Center Pkwy., Suite 230
Roswell, GA 30076

MAR 28 2008

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request General Label Change – Package Size
EPA Reg. No. 60063-28
Application Dated March 4, 2008

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 03/04/08 for the above product. The Registration Division 07(RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



SIPCAM AGRO USA, INC.

March 4, 2008

Ms. Sherada D. Hobgood
Document Processing Desk (NOTIF)
U.S. EPA, Registration Division (7505P)
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Subject: **Turf Fertilizer with Prodiamine 20/75**
EPA Reg. No. 60063-28
Notification of Minor Label Modifications

Dear Ms. Hobgood:


This notification, as per PR Notice 98-10, is to advise the Agency that we wish to add, in addition to use rates in pounds per acre, an option of listing rates in pounds of product per 1,000 square feet. Enclosed are the following documents to support this notification:

- Application for Pesticide (8570-1) with notification box checked
- Revised Draft Labeling

This product is primarily produced for use by homeowners in package sizes ranging from 15 to 25 pounds; however the current use rates in pounds per acre is not practical. Adding the option of use rates listed in pounds per 1,000 square feet for these smaller packages will assist the user in making applications for areas less than an acre. The option to use the rates in pounds per acre is, however, applicable to commercial markets. No other label changes have been made.

If you have any questions, please call me at 770-587-1032.

Sincerely,


Jon M. Gehring
Technical and Regulatory Manager



United States
Environmental Protection Agency
 Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number
NOTIFICATION

Application for Pesticide - Section I

1. Company/Product Number 60063-28	2. EPA Product Manager Miller (23)	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product Name Turf Fertilizer with Prodiamine 20/75	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Sipcam Agro USA, Inc. 300 Colonial Center Pkwy, Suite 230 Roswell, GA 30076 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of Minor Label Changes per PR Notice 98-10: Addition of use rates in lbs./1,000 sq.ft. converting from lbs./Acre
 This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Unit Packaging wgt. No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Jon M. Gehring	Title Technical and Regulatory Manager	Telephone No. (Include Area Code) (770) 587-1032
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Technical and Regulatory Manager	
4. Typed Name Jon M. Gehring	5. Date 3/4/08	



SIPCAM AGRO USA, INC.

[For lbs. of product for specific a.i. %, see chart on page 11.]

Turf Fertilizer With Prodiamine 20/75

Contains (0.20%-0.75%) Cavalcade®

For selective preemergence control of grass and broadleaf weeds in:

- Established Turf Grasses (excluding golf course putting greens) and lawns
- Landscape Ornamentals (including established perennials and wildflower plantings)

20 pounds treats (up to 2,680 sq.ft.)(up to 10,000 sq.ft.)

Active Ingredient:

Prodiamine, [N³,N³-Di-*n*-propyl-2,4-dinitro-6-(trifluoromethyl)-*m*-phenylenediamine 0.20% (0.75%)

Inert Ingredients: 99.80% (99.25%)

Total 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See the following section, **Precautionary Statements, Hazards To Humans and Domestic Animals**, for additional information.

EPA Reg. No. 60063-28

EPA Est. No. [insert] _____

NET WEIGHT: _____ pounds (_____ kg.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center, or doctor for further treatment advice.

Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turf grasses (excluding golf course putting greens) and lawns
- Landscape ornamentals
- Established perennials and wildflower plantings

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when it is activated by at least 1/2 inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown for; (1) Sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Do not apply this product through any type of irrigation system.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Kochia
Bluegrass, Annual (<i>Poa annua</i>)	Lambsquarter, common
Carpetweed	Lovegrass
Chickweed, common	Panicum (Texas, Fall, Browntop)
Chickweed, mouse ear (from seed)	Pigweed
Crabgrass (large, smooth)	Purslane, common
Crowfootgrass	Pusley, Florida
Cupgrass, Woolly	Rescuegrass ³
Foxtails, Annual	Shepherd's Purse ²
Goosegrass ¹	Signalgrass, Broadleaf
Henbit	Speedwell, Persian
Itchgrass	Sprangletop
Johnsongrass (from seed)	Spurge, Prostrate
Junglerice	Witchgrass
Knotweed	Woodsorrel, Yellow (from seed)

¹In many areas a single application of 0.65 lb. to 1.5 lb. active ingredient per acre (equal to xx lbs. to xxx lbs./A of this product per acre [or] equal to x.x to xx.x lbs./1000 sq.ft. of this product) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 0.65 lb. to 1.0 lb. a.i. per acre (equal to xx lbs. to xxx lbs./A of this product per acre [or] equal to x.x to xx.x lbs./1000 sq.ft. of this product) followed after 60 to 90 days by a second

application at doses that would not exceed those given in the **Maximum Annual Rate Table**. Do not exceed the maximum rate for turf grass species listed in the **Maximum Annual Rate Table** above.

²Applications for this weed should be made in late summer, fall, or winter prior to germination.

³Suppression only. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.


APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF:

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the **Maximum Annual Rates** section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES



USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year. (4) Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (6) Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate/Calendar Year of turf fertilizer by turf grass species ⁽¹⁾					
Turf Species:	Lbs. Product/A [or]		Lbs. Product /1000 sq.ft.		Lbs. ai/A
	[0.20% a.i.]	[0.75% a.i.]	[0.20% a.i.]	[0.75% a.i.]	
Creeping Bentgrass	325	86	<u>7.5</u>	<u>2.0</u>	0.65
Creeping Red Fescue	375	100	<u>8.6</u>	<u>2.3</u>	0.75
Buffalograss Kentucky Bluegrass Perennial Ryegrass	500	133	<u>11.5</u>	<u>3.1</u>	1.0
Bermudagrass ⁽²⁾ Bahia grass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysia	750	200	<u>17.2</u>	<u>4.6</u>	1.5

¹These are the maximum rates per calendar year by species limitations.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 0.5 lb. a.i./acre (equal to xx lbs./A of this product [or] equal to x.x to xx.x lbs./1000 sq.ft. of this product). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass, and Rescuegrass due to reduced product rates used in sprigging situations.

- Do not apply more than 1.5 lbs. a.i. per calendar year per acre (equal to xxx lbs. of this product [or] equal to x.x to xx.x lbs./1000 sq.ft. of this product).
- Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates Table.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the following table below for best overseeding/reseeding results.

Turf fertilizer + Prodiamine Rate (lb./A)		[or]	Turf fertilizer + Prodiamine Rate (lb./1000 sq.ft.)		ai/A	Months Before Overseeding		
[0.20%]	[0.75%]		[0.20%]	[0.75%]		North	Transition	South
250	67		<u>5.7</u>	<u>1.5</u>	0.50	4	4	4
325	86		<u>7.5</u>	<u>2.0</u>	0.65	5	4	4
375	100		<u>8.6</u>	<u>2.3</u>	0.75	6	5	5
400	106		<u>9.2</u>	<u>2.4</u>	0.80	-	6	6
500	133		<u>11.5</u>	<u>3.1</u>	1.00	-	7	7
570	152		<u>13.1</u>	<u>3.5</u>	1.14	-	-	9
650	173		<u>14.9</u>	<u>4.0</u>	1.30	-	-	10
750	200		<u>17.2</u>	<u>4.6</u>	1.50	-	-	12

Guaranteed Fertilizer Analysis:

Total Nitrogen (N) %

Available Phosphate (P₂O₅) %Soluble Potash (K₂O) %

Derived from: [Ureaformaldehyde, Urea, Sulfur-Coated Urea, Ammonium Phosphate, Muriate of Potash]

Spreader Settings for 20 lbs.	2,680 sq. ft.	X,XXX sq. ft.	10,000 sq. ft.
Scott's Evergreen 18"	# XX	# XX	# XX
Scott's Speedy Broadcast	# XX	# XX	# XX
Red Devil Broadcast	# XX	# XX	# XX
Republic EZ Drop 18"	# XX	# XX	# XX
Republic EZ Broadcast	# XX	# XX	# XX
Republic EZ Trctr. Broadcast	# XX	# XX	# XX

LANDSCAPE ORNAMENTALS**Including Established Perennials and Wildflower Plantings**

This product may be applied for residual preemergence weed control in ornamentals.

Use rates and When to Apply

Apply in fall and/or spring, 0.65 to 1.5 lbs. active ingredient per acre, which is equal to xx to xxx lbs. of this product per 1,000 sq.ft. of ornamental plantings; therefore 20 lbs of this product will treat x,xxx to x,xxx sq.ft. Use the higher rates of application for longer control periods. Sequential applications may be made as long as the total amount of product applied does not exceed the maximum annual application rate of 1.5 lbs. a.i. per acre.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate it into the soil.

Newly Transplanted Landscape Ornamentals

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when it is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific name	Common name (Tested varieties)
<i>Abies</i> spp.	Fir
<i>Abelia grandiflora</i>	Abelia; Sherwood
<i>Acer palmatum</i>	Japanese maple
<i>Acer platanoides</i>	Norway maple
<i>Achillea</i> sp.	Yarrow (King Edward)

Scientific name (cont.)	Common name (Tested varieties)
<i>Actinidia chinensis</i> *	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile; African Lily
<i>Akebia quintata</i>	Five-leaf or Chocolate vine
<i>Allium cernuum</i>	Lady's leek or Nodding onion
<i>Anemone hybrida</i>	Japanese anemone
<i>Aquilegia</i> spp.	Columbine (Red and gold)
<i>Arctostaphylos densiflora</i>	Vine Hill manzanita
<i>Arctotheca calendula</i>	Cape weed
<i>Artemisia</i> spp.	Wormwood (Silver mound; Castle)
<i>Aster</i> sp. or <i>Symphiotrichum</i> sp.	Aster (Bonny blue, Purple dome)
<i>Aster X frikartii</i>	
<i>Athyrium felix-femina</i>	Lady fern
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Begonia</i> spp.	Fibrous bigonia (Hardy Grandis)
<i>Berberis gladwynensis</i>	Barberry
<i>Berberis julianae</i>	Wintergreen barberry
<i>Berberis mentorensis</i>	Mentor barberry
<i>Berberis thunbergii</i>	Japanese barberry
<i>Berberis verruculosa</i>	Warty barberry
<i>Bergenia cordifolia</i>	
<i>Boltonia asteroides</i>	Snowbank
<i>Bougainvillea</i> spp.	Bougainvillea
<i>Buddleia davidii</i>	Butterfly-bush (Dwarf Blue, Royal Red)
<i>Buxus microphylla</i>	Japanese boxwood
<i>Callistemon viminalis</i>	Weeping bottlebrush
<i>Calluna vulgaris</i>	Scotch heather
<i>Campanula carpatica</i>	Tussock bellflower (White Clips)
<i>Campsis X tagliabuana</i>	Trumpet creeper; Trumpet flower (Madame Gaten)
<i>Carpobrotus edulis</i>	Hottentot fig; Ice plant
<i>Cassia artemisoides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Ceratostigma plumbaginoides</i>	
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Chrysanthemum nipponicum</i>	
<i>Cleyera japonica</i>	Cleyera
<i>Citrus</i> spp. *	Citrus species*
<i>Coreopsis</i> spp.	Coreopsis; Calliopsis (Early sunrise, Moonbeam)
<i>Cornus florida</i>	Flowering dogwood
<i>Cornus stolonifera</i>	American dogwood
<i>Cortaderia selloana</i>	Pampas grass
<i>Cotoneaster apiculatus</i>	Cranberry cotoneaster
<i>Cotoneaster buxifolius</i>	Cotoneaster
<i>Cotoneaster dammeri</i>	Bearberry cotoneaster
<i>Cotoneaster microphyllus</i>	Rockspray cotoneaster
<i>Crataegus</i> spp.	Hawthorne
<i>Crocasmia</i> spp.	Lucifer
<i>Cupressus sempervirens</i>	Italian cypress
<i>Delosperma alba</i>	White trailing ice plant
<i>Delphinium</i> spp.	Larkspur (Blue elf)

Scientific name (cont.)	Common name (Tested varieties)
<i>Dianthus deltoides</i>	Maiden pinks (Zing)
<i>Dianthus gratianopolitanus</i>	Cheddar pinks
<i>Dodonea viscosa</i>	Hop bush
<i>Echinacea purpurea</i>	Coneflower, purple (Magnus)
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus fortunei</i>	Wintercreeper
<i>Euonymus japonica</i>	Japanese spindle tree; Evergreen euonymus
<i>Euonymus kiautschovica</i>	Spreading euonymus
<i>Fatsia japonica</i>	Japanese aralia
<i>Forsythia intermedia</i>	Border forsythia
<i>Forsythia suspense</i>	Weeping forsythia
<i>Forsythia viridissima</i>	Greenstem forsythia
<i>Gaillardia</i> spp.	Blanket flower (Goblin)
<i>Gardenia jasminoides</i>	Gardenia, Cape-jasmine
<i>Gentiana dahurica</i>	Gentian
<i>Geranium cinereum</i>	Cranesbill
<i>Gladiolus</i> spp.**	Gladiolus
<i>Gypsophila repens</i>	Baby's breath
<i>Hedera helix</i>	English Ivy
<i>Helianthemum</i> spp.	Sunrose
<i>Hermerocallis</i> spp.	Daylily (Aztec gold, Stella d'Oro, Tender love)
<i>Hibiscus</i> spp.**	Rose of Sharon**
<i>Hibiscus</i> spp.	Mallow (Disco belle white)
<i>Hibiscus Rosa-sinensis</i> **	Chinese hibiscus**
<i>Hosta plantaginea</i>	Hosta, Plantain lily
<i>H. sieboldiana</i>	Hosta (Searsucker)
<i>Houttuynia cordata</i> var. <i>variegata</i>	
<i>Hydrangea macrophylla</i>	Bigleaf hydrangea
<i>Ilex cornuta</i> **	Chinese holly**
<i>Ilex crenata</i>	Japanese holly
<i>Ilex opaca</i>	American holly
<i>Ilex pernyi</i>	Holly
<i>Ilex vomitoria</i>	Yaupon holly
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-leaved iris (Jodlesong)
<i>Iris siberica</i>	Siberian iris (Cabernet)
<i>Iris</i> spp.	Iris **
<i>Jasminum nudiflorum</i>	Winter jasmine
<i>Juniperus chinensis</i>	Chinese juniper (Nick's compact)
<i>Juniperus conferta</i>	Shore juniper (Blue Pacific)
<i>Juniperus davurica</i>	Juniper
<i>Juniperus horizontalis</i>	Creeping juniper
<i>Juglans</i> spp.*	Walnut*
<i>Justicia brandegeana</i>	Shrimp plant
<i>Lagerstroemia indica</i>	Crape myrtle
<i>Lantana montevidensis</i>	Weeping lantana
<i>Lavender</i> spp.	Lavender (Munstead)
<i>Leontopodium alpinum</i>	Edelweiss
<i>Ligustrum amurense</i>	Amur privet

Scientific name (cont.)	Common name (Tested varieties)
<i>Ligustrum japonicum</i>	Japanese privet
<i>Ligustrum lucidum</i>	Glossy privet (Wax-Leaf)
<i>Ligustrum sinense</i>	Chinese privet (Variegata)
<i>Lilium spp.</i>	Lily (Jazz)
<i>Liriope muscari</i>	Liriope (Big blue, Lilyturf, Silver mound, Evergreen giant)
<i>Liriope muscari var. variegata</i>	Liriope, variegated
<i>Liriope spicata</i>	Liriope, creeping
<i>Lobelia cardinalis</i>	Cardinal flower
<i>Lonicera japonica</i>	Japanese honeysuckle
<i>Lonicera tatarica</i>	Tatarian honeysuckle
<i>Loropetalum chinense</i>	Burgundy
<i>Lythrum spp.</i>	Loosestrife (Modern Pink)
<i>Magnolia spp.**</i>	Magnolia**
<i>Maleophora luteola</i>	Ice plant
<i>Malus spp.*</i>	Crabapple*
<i>Miscanthus sinensis</i>	Yaku jima, Silberfeder
<i>Nandina domestica</i>	Heavenly bamboo
<i>Narcissus spp.**</i>	Narcissus**
<i>Nerium oleander</i>	Oleander
<i>Olea europaea*</i>	Olive*
<i>Ophiopogon japonicus**</i>	Mondo Grass**
<i>Osmanthus heterophyllus</i>	Osmanthus; False holly (Gulf tide)
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendron luteum</i>	Sourwood
<i>Persea americana</i>	Avocado*
<i>Photinia fraseri</i>	Frasier's photinia; Redtip
<i>Physostegia virginiana</i>	False dragonhead (Vivid)
<i>Picea spp.**</i>	Spruce species** (Colorado blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-valley shrub
<i>Pinus brutia</i>	Calabrian pine
<i>Pinus canariensis</i>	Canary island pine
<i>Pinus elliottii</i>	Slash pine
<i>Pinus halepensis</i>	Aleppo pine
<i>Pinus nigra</i>	Austrian black pine
<i>Pinus palustris</i>	Longleaf pine
<i>Pinus radiata</i>	Monterey pine
<i>Pinus strobus</i>	Eastern white pine
<i>Pinus sylvestris</i>	Scotch pine
<i>Pinus taeda</i>	Loblolly pine
<i>Pinus thunbergiana</i>	Japanese black pine
<i>Pinus virginiana</i>	Virginia pine
<i>Pistacia spp.*</i>	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland pittosporum
<i>Pittosporum tobira</i>	Japanese pittosporum
<i>Podocarpus macrophyllus</i>	Japanese yew
<i>Prunus laurocerasus</i>	English laurel
<i>Prunus spp.*</i>	Almond, Apricot, Nectarine, Peach, Plum, and Prune*
<i>Pseudotsuga menziesii**</i>	Douglas-fir**

12/14

Scientific name (cont.)	Common name (Tested varieties)
<i>Pyracantha coccinea</i>	Firethorn, scarlet
<i>Pyracantha fortuneana</i>	Firethorn
<i>Pyracantha koidzumii</i>	Firethorn
<i>Pyrus</i> spp.	Bradford pear
<i>Quercus rubra</i>	Red oak
<i>Quercus shumardii</i>	Shumard oak
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Rhododendron</i> (including Azalea)	Rhododendron & Azalea (Coral Bells, Delaware valley white, Flame creeper, George L. Tabor, PJM, Roseum Elegans, Formosa, Hino- crimson, Wakaebisu, Gumpo)
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i> *	Rosemary*
<i>Rudbeckia</i> spp.	Black-eyed Susan
<i>Rumohra adiantiformis</i>	Leatherleaf Fern
<i>Santolina virens</i>	Goldstrum Santolina
<i>Saxifraga</i> spp.	Saxifrage, Purple dome
<i>Sedum album</i>	Stonecrop
<i>Sedum caulicola</i>	Stonecrop, likadense
<i>Sedum dasyphyllum</i>	Stonecrop
<i>Syzygium paniculatum</i>	Japanese boxcherry
<i>Taxus cuspidata</i>	Japanese yew
<i>Taxus media</i>	Yew
<i>Teucrium</i> spp.	Germander
<i>Thalictrum dipterocarpum</i>	Meadow rue
<i>Thuja occidentalis</i>	American arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada hemlock
<i>Tulipa</i> spp.	Tulip
<i>Viburnum japonicum</i>	Japanese viburnum
<i>Viburnum odoratissimum</i>	Sweet viburnum
<i>Viburnum plicatum</i>	Japanese snowball
<i>Viburnum rigidum</i>	Canary Island Viburnum
<i>Viburnum tinus</i>	Laurustinus
<i>Viburnum trilobium</i>	Cranberry Bush
<i>Viburnum wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Greater periwinkle, Vinca
<i>Vinca minor</i>	Common periwinkle
<i>Vitis</i> spp.*	Grape*
<i>Weigela florida</i>	Old fashioned Weigela
<i>Yucca aloifolia</i>	Spanish bayonet
<i>Yucca filamentosa</i>	Yucca, Adam's needle

* Do not use on food producing trees, vines, or plants.

** Not for use on container grown plants

STORAGE AND DISPOSAL

STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available.

IF PARTLY FILLED: Call you local waster agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor ore outdoor drain.

SPILL: In case of spill, sweep up and dispose of material according to Product disposal directions above.

CONDITIONS OF SALE AND WARRANTY

The **Directions for Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Sipcam Agro USA or the Seller. All such risks shall be assumed by the Buyer. Sipcam Agro USA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions for Use** subject to the inherent risks referred to above. Sipcam Agro USA makes no other express or implied **Warranty of Fitness or Merchantability** or any other express or implied warranty. In no case shall Sipcam Agro USA or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this Product. Sipcam Agro USA and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Sipcam Agro USA.

SIPCAM AGRO USA, INC.
300 COLONIAL CENTER PKWY, SUITE 230
ROSWELL, GA 30076

11/13/02 EPA Stamped Label

3/4/08 Notification to add Lbs./1000 sq.ft.

[Chart showing each of the approved ai loads and rates for Turf Fertilizer with Prodiamine 20/75]

Maximum Application Rate/Calendar Year of turf fertilizer by turf grass species (1)

		Formulated Product a.i. Load													
		0.20%		0.28%		0.29%		0.37%		0.383%		0.42%		0.75%	
Turf Species:	Lbs. ai/A	Lbs/A	Lbs/1000 sq.ft.	Lbs/A	Lbs/1000 sq.ft.	Lbs/A	Lbs/1000 sq.ft.	Lbs/A	Lbs/1000 sq.ft.	Lbs/A	Lbs/1000 sq.ft.	Lbs/A	Lbs/1000 sq.ft.	Lbs/A	Lbs/1000 sq.ft.
Creeping Bentgrass	0.65	325	7.5	236	5.4	224	5.1	176	4.0	170	3.9	155	3.6	87	2.0
Creeping Red Fescue	0.75	375	8.6	273	6.3	259	5.9	203	4.7	196	4.5	179	4.1	100	2.3
Buffalograss Kentucky Bluegrass Perennial Ryegrass	1	500	11.5	364	8.3	345	7.9	270	6.2	261	6.0	238	5.5	133	3.1
Bermudagrass(2) Bahia grass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf- type) Zoysia	1.5	750	17.2	545	12.5	517	11.9	405	9.3	392	9.0	357	8.2	200	4.6
Rate Guidance Table	0.50	250	5.7	182	4.2	172	4.0	135	3.1	131	3.0	119	2.7	67	1.5
	0.65	325	7.5	236	5.4	224	5.1	176	4.0	170	3.9	155	3.6	86	2.0
	0.75	375	8.6	273	6.3	259	5.9	203	4.7	196	4.5	179	4.1	100	2.3
	0.80	400	9.2	291	6.7	276	6.3	216	5.0	209	4.8	190	4.4	106	2.4
	1.00	500	11.5	364	8.3	345	7.9	270	6.2	261	6.0	238	5.5	133	3.1
	1.14	570	13.1	415	9.5	393	9.0	308	7.1	298	6.8	271	6.2	152	3.5
	1.30	650	14.9	473	10.9	448	10.3	351	8.1	339	7.8	310	7.1	173	4.0
	1.50	750	17.2	545	12.5	517	11.9	405	9.3	392	9.0	357	8.2	200	4.6
20 lbs. treats up to ...sq. ft.		2,681		3,686		3,887		4,959		5,133		5,629		10,052	

b11/14