

60063-29

12/4/2009

10/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Joseph W. Burley, Ph.D.
Sipcam Agro USA, Inc.
2520 Meridian parkway, Ste 525
Durham, NC 27713

DEC 4 2009

Subject: Label Notification(s) for Pesticide Registration Notices 2007-4 and 98-10
1. Other minor revisions

Dear Dr. Burley:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notices (PRN) 2007-4 and 98-10 dated August 5, 2009 for:

| | |
|----------------------------------|---------------------------------|
| EPA Registration 60063-4 | Echo Zn |
| EPA Registration 60063-7 | Echo 720 |
| EPA Registration 60063-29 | Sipcam Tebuconazole 3.6F |
| EPA Registration 60063-31 | Torrent 2F |

The Registration Division (RD) has conducted a review of this request for applicability under PR Notices 2007-4 and 98-10 and finds that the label changes requested falls within the scope of PR Notices 2007-4 and 98-10. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Nicole Williams of my staff at 703-308-5551.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

20f11

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval Expires 05-31-98.



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

| | | |
|--|--|--|
| 1. Company/Product Number 60063-29 | 2. EPA Product Manager Mary Waller | 3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/Product Name Sipcam Tebuconazole 3.6F | PM# 21 | |
| 5. Name and Address of Applicant (Include ZIP Code) Sipcam Agro USA, Inc. 2520 Meridian Parkway, Suite 525 Durham, NC 27713 <input checked="" type="checkbox"/> Check if this is a new address | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____ | |

Section - II

| | | |
|--|--|---|
| <input type="checkbox"/> Amendment - Explain below. | <input type="checkbox"/> Final printed labels in response to Agency letter dated | NOTIFICATION DEC - 4 2009 |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application. | |
| <input checked="" type="checkbox"/> Notification - Explain below. | <input type="checkbox"/> Other - Explain below. | |

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146 and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this project except as certified in the paragraph below. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146 and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. [See attached page for further certification statement.]

Section - III

| | | | |
|---|---|---|---|
| 1. Material This Product Will Be Packaged In: | | | |
| Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No | Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If "Yes" Unit Packaging wgt. No. per container | Water Soluble Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No If "Yes" Unit Packaging wgt. No. per container | 2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____ |
| * Certification must be submitted | | | |
| 3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container | 4. Size(s) Retail Container 2.5 gal. | 5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product | |
| 6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled | | | |

Section - IV

| | | |
|---|-----------------------------|---|
| 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) | | |
| Name Joseph W. Burley | Title Regulatory Manager | Telephone No. (Include Area Code) (919) 226 1297 |

| | | |
|--|--------------------------------|--|
| Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. | | 6. Date Application Received (Stamped) |
| 2. Signature | 3. Title Regulatory Manager | |
| 4. Typed Name Joseph W. Burley | 5. Date 08/05/2009 | |

30/11

Section II Explanation (continued):

Notification of addition of EPA approved language regarding certified adjuvants; and of addition (as per PR Notice 2001-5) of voluntary resistance management labeling per PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the Confidential Statement of Formula except as certified in the paragraph above. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.



40411
✓

August 7, 2009

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Subject: **Sipcam Tebuconazole 3.6F**
EPA Reg. No. 60063-29
Notifications per PR Notice 2007-4 and PR Notice 98-10

Dear Sir or Madam:

In support of the above notifications, please find the following documents enclosed:


- Application for Pesticide Notification, Form 8570-1
- One copy of the product labeling, which is annotated to enable the Agency to differentiate the changes which have been made to that label
- One clean copy of the label, containing the incorporated changes
- Copy of a letter from Mr. Donald Stubbs which acknowledges that the CPDA language appropriate to approved adjuvants can be added to a registrant's label by notification under the procedures set forth in PR Notice 98-10.

For ease of review by the Agency, please note that changes to this product label are limited to those found on pages 1, 4, 6 and 7. Those changes have been underlined and highlighted in the annotated copy.

If you have any questions about this submission, please feel free to call me at 770-594-6356

Sincerely,

Joseph W. Burley PhD
Regulatory Manager
Sipcam Agro USA, Inc.



50f11



SIPCAM AGRO USA, INC.

NOTIFICATION

DEC - 4 2009

Sipcam Tebuconazole 3.6F

| | | |
|-------|---|-----------|
| GROUP | 3 | FUNGICIDE |
|-------|---|-----------|

Active Ingredients:

Tebuconazole {alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1,-dimethylethyl-1H-1,2,4-triazole-1-ethanol)} .. 38.7%

Other Ingredients: 61.3%

Total: 100.0%


Contains 3.6 pounds tebuconazole per gallon

Keep Out of Reach of Children

CAUTION

| FIRST AID | |
|--|--|
| IF SWALLOWED | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have affected person sip a glass of water if able to swallow. • Do not induce vomiting unless told by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| IF ON SKIN OR CLOTHING | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| IF INHALED | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. • Call a poison control center or doctor for further treatment advice. |
| IF IN EYES | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. | |
| Emergency phone numbers | (800) 222-1222 Poison Control Center (800) 858-7378 NPIC (human and animal health) (800) 424-9300 CHEMTREC (transportation and spills) |
| NOTE TO PHYSICIAN: No specific antidote. Treat symptomatically. This product may be a dermal sensitizer. | |

Manufactured for
 Sipcam Agro USA, Inc.
 2520 Meridian Parkway, Suite 525
 Durham, NC 27713



EPA Reg. No. 60063-29

Net Contents: _____ gallons

EPA Est. No. _____

60911

Precautionary Statements Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Shoes and socks;
- Chemical-resistant gloves such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:


- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours. This pesticide is toxic to estuarine and marine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.



7 of 11

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

General Precautions and Restrictions

Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, and estuaries:

- Apply only during alternate years in fields adjacent to aquatic areas listed above.
- Do not apply by ground or air within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

Spray drift management: For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.


Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Rotational crops: Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

Mixing: Dilute this product into water, then apply by means of spraying equipment which is properly engineered and calibrated to deliver uniform deposit onto surfaces of the crops which are approved for use of this product. Add specified amount of Sipcam Tebuconazole 3.6 F into the mixing tank while filling with water to the desired level.



80f11

Operate the agitator while mixing. If other materials are added to the mixing and/or sprayer tank, this product should be thoroughly dispersed into the water carrier prior to the addition of other materials.

Spray Volume: Apply this product in a minimum of 10 gallons of water by ground sprayer, or in a minimum of 5 gallons of water per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary for complete crop coverage, increase the volume rate of water carrier.


Chemigation: Do not apply this product through any type of irrigation system.

Compatibility:

- To determine the physical compatibility of Sipcam Tebuconazole 3.6 F with other products, the following procedure should be followed: Pour the specified proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains uniformly mixed or can be re-mixed readily, the mixture is considered physically compatible.
- Biological compatibility, i.e. effectiveness for disease control or lack of phytotoxicity, must be determined on a case-by-case basis. Do not mix this product with any other pesticide, fertilizer, adjuvant or any other materials except water unless you know that such mixtures will be effective and non-injurious to your crop under your conditions of use. When an adjuvant is to be used with this product, Sipcam Advan USA recommends the use of a Chemical Producers and Distributors Association (CPDA) certified adjuvant.

Approved Crop Uses

| CROP | PHI (DAYS) | DISEASES | RATE OF PRODUCT PER ACRE | APPLICATION DIRECTIONS |
|------------------------|------------|--|--|--|
| Grasses Grown for Seed | 4 | Rusts (<i>Puccinia</i> spp.), Powdery mildew | 4 to 8 fl oz per application Maximum 16 fl oz per crop season | Apply when weather conditions are favorable for disease development or when first signs of disease are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals. |
| | | <p>General directions: Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.</p> <p>For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with this product. Chaff, screenings and straw from treated areas may be used for feed purposes, however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.</p> | | |



9 of 11

| CROP | PHI (DAYS) | DISEASES | RATE OF PRODUCT PER ACRE | APPLICATION DIRECTIONS |
|--|------------|--|---|---|
| Peanut | 14 | <p>Soilborne: Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only)</p> <p>Foliar: Early leaf spot, Late leaf spot, Leaf rust, Web blotch (<i>Phoma</i>), Pepper spot (<i>Leptosphaerulina</i>)</p> | <p>7.2 fl oz per acre</p> <p>Maximum 28.8 fl oz per crop season</p> | <p>FOUR-APPLICATION PROGRAM: Apply this product consecutively four times per growing season within a preventive schedule of fungicide applications that are typically used for the control of peanut foliar diseases. Repeat applications of this product at 14-day intervals. For optimal control of soilborne diseases, the first application of this product should be made not later than the beginning of pegging, approximately 45 to 60 days after planting. Applications of this product should be preceded by and followed by products containing chlorothalonil active ingredient, such as Echo[®], in accordance with their label(s).</p> <p>LEAF SPOT ADVISORY SCHEDULE: For control of soilborne diseases in an advisory schedule, apply this product in the first advisory spray in July and continue applications at 14-day intervals. Applications after mid-August should be tank mixed with a product containing chlorothalonil for resistance management purposes.</p> |
| <p>General directions: For optimum control of the specified soilborne diseases, four consecutive applications of this product must be made at 14-day intervals. Do not feed hay or threshings to livestock, or allow livestock to graze in treated areas. Tebuconazole, the active ingredient in this product, is a sterol demethylation inhibitor (DMI) fungicide. A chlorothalonil product may be tank mixed at the rate of 0.75 lbs of active ingredient with this product as a leaf spot resistance management strategy. A spray surfactant is not necessary when this product is tank mixed with a chlorothalonil product. Mixing or alternating this product with other sterol DMI fungicides may lead to resistance. The active ingredient in this product must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i>. Drought conditions may decrease the effectiveness of this product against root and pod rots. Use this product in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.</p> | | | | |
| CROP | PHI (DAYS) | DISEASES | RATE OF PRODUCT PER ACRE | APPLICATION DIRECTIONS |
| Soybeans | 21 | Rust (<i>Phakopsora pachyrhizii</i>) | <p>3 to 4 fl oz per acre</p> <p>Maximum 12 fl oz per acre per crop season</p> | <p>At the first sign of rust pustules on foliage make a spray application of this product. If environmental conditions are favorable to continued disease development, make a second application after 10 to 21 days dependent upon the severity of disease pressure. Do not apply more than 0.225 pounds of the active ingredient, tebuconazole, per acre per year and make no more than 3 applications of this product per season. Apply this product in a minimum of 10 gallons of spray suspension per acre by ground sprayer or in a minimum of 5 gallons of spray suspension per acre by aircraft.</p> |
| <p>General directions: Observe fields closely for early disease symptoms. For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with this product. Do not enter treated fields within 12 hr of application.</p> | | | | |
| CROP | PHI | | RATE OF | APPLICATION DIRECTIONS |

| | (DAYS) | DISEASES | PRODUCT PER ACRE | |
|--|------------|--|--|--|
| Sunflowers | 50 | Rust (<i>Puccinia spp.</i>) | 6 fl oz per acre Maximum 16 fl oz per crop season | <p>Make a foliar spray application of this product (6 fl oz per acre) at the first sign of rust infection. Repeat this application after 14 days if the severity of the infection warrants treatment or if conditions conducive to infection prevail. Do not exceed the maximum specified treatment rate (16 fl oz) per crop season.</p> <p>Apply the specified treatment rate in a minimum quantity of 20 gallons of water per acre when making application of this product with ground sprayers. When making applications by aircraft use a minimum quantity of 5 gallons of water to disperse the specified application rate. Do not exceed the maximum specified application rate per crop per season.</p> |
| <p>General directions: Observe fields closely for early disease symptoms. For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with this product. Thorough coverage is important for optimum disease control.</p> | | | | |
| CROP | PHI (DAYS) | DISEASES | RATE OF PRODUCT PER ACRE | APPLICATION DIRECTIONS |
| Wheat, Barley | 30 | Rusts (leaf, stem, and stripe) (<i>Puccinia spp.</i>) Scab or head blight (<i>Fusarium graminearum</i>) - suppression | 4 fl oz per acre Maximum 4 fl oz per acre per crop season | <p>Rusts: Make an application of this product at the first sign of rust pustules on foliage. Fusarium head blight: The best time to apply this product to suppress Fusarium head blight is at the beginning of anthesis (flowering) of the main head, Feekes growth stage 10.5.1. Apply this product in a minimum of 10 gallons of spray suspension per acre by ground sprayer or in a minimum of 5 gallons of spray suspension per acre by aircraft.</p> |
| <p>General directions: Observe fields closely for early disease symptoms under prolonged conditions favorable for disease development and/or when varieties susceptible to these diseases are planted. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after application of this product. Straw cut after harvest may be fed to livestock or used for bedding. For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with this product. The plant will absorb the active ingredient systemically after two to four hours following application prior to irrigation or rain; after which, this product will resist weathering. Tebuconazole, the active ingredient in this product, is a sterol demethylation inhibitor (DMI) fungicide.</p> | | | | |

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10



seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND LIMITATION OF DAMAGES

CONDITIONS OF SALE: To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions stated on the label. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. **SIPCAM AGRO USA, INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS OR MERCHANTABILITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SIPCAM AGRO USA, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND SIPCAM AGRO USA, INC.'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. BUYER AND USER ACKNOWLEDGE, AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. SIPCAM AGRO USA, INC. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.**

Echo is a registered trademark of Sipcam Agro USA Inc.

