

60063-51

7/23/2012

10914



U S ENVIRONMENTAL PROTECTION AGENCY

Office of Chemical Safety and Pollution Prevention
Office of Pesticide Programs
Registration Division (7504P)
1200 Pennsylvania Ave N W
Washington DC 20460

EPA Reg Number

60063 51

Date of Issuance

JUL 23 2012

Term of Issuance

Unconditional

Name of Pesticide Product

Iprodione ETQ

NOTICE OF PESTICIDE

- X Registration
- Reregistration
Under FIFRA as amended

Name and Address of Registrant (include ZIP Code)

Sipcam Agro, USA, Inc
2520 Meridian Parkway, Suite 525
Durham, NC 27713

Mailed to

Joseph W Burley
Registration Manager

Note Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant the above named pesticide is hereby registered under the Federal Insecticide Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment the Administrator on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA section 3(c)(5) provided that you

- 1 Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data and submit acceptable responses required for reregistration of your product under FIFRA section 4

Page 1 of 2

Signature of Approving Official

Mary L Waller

Date

7/23/2012

Mary L Waller Product Manager (21)
Fungicide Branch/Registration Division/OPP/OCSP (7504P)

2014

- 2 You must submit the following data before the due date of 1/31/2014
  - a Storage Stability (830 6317) and Corrosion Characteristics (830 6320) studies
- 3 Make the following changes to the label
  - a At the bottom of page 1 change the product registration number to EPA Registration No 60063 51
- 4 Submit one copy of the revised final printed label for the record before the product is released for shipment

Your release for shipment of the product constitutes acceptance of these conditions

A copy of the label stamped Accepted with Comments is enclosed for your records



Mary L Waller  
Product Manager (21)  
Fungicide Branch  
Registration Division (7504P)

Enclosure

Label stamped Accepted with Comments  
Product Chemistry Review DP401596 dated June 19 2012  
Acute Toxicity Review DP401597 dated June 19 2012

40414

ACCEPTED  
with COMMENTS  
In EPA Letter Dated:

7/23/2012

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No. 60063-51

# Iprodione ETQ

For control of listed diseases on ornamentals and turf grasses

## ACTIVE INGREDIENT

Iprodione 3 (3 5 dichlorophenyl) N (1 methylethyl) 2 4 dioxo 1  
imidazolincarboxamide 23 3%

**OTHER INGREDIENTS** 76 7%

**TOTAL** 100 0%

Contains 2 pounds Iprodione per gallon

## KEEP OUT OF REACH OF CHILDREN

### CAUTION/PRECAUCION

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en  
detalle (If you do not understand the label find someone to explain it to you in detail )

#### FIRST AID

<b>If swallowed</b>	Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by the poison control center or doctor Do not give anything by mouth to an unconscious person
<b>If in eyes</b>	Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
<b>If on skin or clothing</b>	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice
<b>If inhaled</b>	Move person to fresh air If person is not breathing call 911 or an ambulance then give artificial respiration preferably mouth to mouth if possible Call a poison control center or doctor for further treatment advice

Have the product container or label with you when calling a poison control center or doctor or going for treatment For Emergency Medical Assistance call the National Pesticide Information Center 1 800 858 7378

For chemical emergency spill leak fire exposure or accident call CHEMTREC 1 800 424 9300

Manufactured for  
Sipcam Agro USA Inc  
2520 Meridian Parkway Suite 525  
Durham NC 27713

EPA Registration No 60063 LR

EPA Est No \_\_\_\_\_

Net Contents

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION/PRECAUCION**

Harmful if swallowed or inhaled Causes moderate eye irritation Avoid contact with eyes or clothing Wash thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Mixers, loaders, others exposed to the concentrate and applicators applying as a dip treatment must wear**

- Long sleeved shirt and long pants
- Chemical resistant gloves such as barrier laminate nitrile rubber neoprene rubber or Viton
- Chemical resistant apron
- Chemical resistant footwear plus socks

**Applicators using hand held equipment must wear**

- Coveralls over long sleeved shirt and long pants
- Chemical resistant gloves such as barrier laminate nitrile rubber neoprene rubber or Viton
- Chemical resistant footwear plus socks
- Chemical resistant headgear for overhead exposure
- NIOSH approved respirator with any R P or HE filter

**Applicators using aircraft or mechanical ground equipment and flaggers for aerial application must wear**

- Long sleeved shirt and long pants
- Shoes plus socks

Applicators using truck mounted equipment with a handgun at the end of a hose (i e for commercial turfgrass or ornamental applications) and all other handlers not specified above must wear

- Long sleeved shirt and long pants
- Chemical resistant gloves such as barrier laminate nitrile rubber neoprene rubber or Viton
- Shoes plus socks

Follow manufacturer s instructions for cleaning/maintaining PPE If no instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

Discard clothing or other materials that have been drenched or heavily contaminated w th this products concentrate Do not reuse them

When applicators use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(5)] the handler PPE requirements may be reduced or modified as specified in the WPS

**USER SAFETY RECOMMENDATIONS**

**Users should**

- Wash thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

**ENVIRONMENTAL HAZARDS**

This chemical can contaminate surface water through aerial and ground spray applications Under some conditions it may also have a high potential for runoff into surface water after application These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters frequently flooded areas areas overlaying extremely shallow ground water areas with in field canals or ditches that drain to surface water areas not separated from adjacent surface waters with vegetated filter strips and areas overlaying tile drainage systems that drain to surface water

This pesticide is toxic to invertebrates Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Drift or runoff from treated areas is hazardous to aquatic invertebrates in neighboring areas Do not contaminate water when disposing of equipment wash water or rinsate

**DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling** Do not apply this product in a way that will contact workers or other persons either directly or indirectly through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS) 40 CFR part 170 This Standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval The requirements in this box only apply to uses of this product that are covered by the WPS

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for ornamental uses. The REI for all other uses is 24 hrs

PPE required for early entry into treated areas that is permitted under the WPS and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves such as barrier laminate, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, or viton  $\geq$  14 mils
- Shoes plus socks

**NON AGRICULTURAL USE REQUIREMENTS**

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated areas until sprays have dried.

**USE PRECAUTIONS AND RESTRICTIONS**

Read the entire Directions for Use and Conditions of Sale before using this product.

- Use of this product at residential sites is prohibited.
- Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh, or natural pond, estuary, or commercial fish pond, there must be at least a 25ft vegetative buffer strip between the water body and the point of application.
- For golf courses only, do not apply to turf cut higher than 1" on golf courses where water bodies are present.
- **Do not** apply this product when the wind direction is toward aquatic areas.

**SPRAY DRIFT MANAGEMENT**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they must be observed.

**Aerial Drift Advisory Information**  
**INFORMATION ON DROPLET SIZE**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and

control Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly or under unfavorable conditions (see Wind Temperature)

#### CONTROLLING DROPLET SIZE

**Volume** Use high flow rate nozzles to apply the highest practical spray volume

Nozzles with higher rated flows produce larger droplets

**Pressure** Do not exceed the nozzle manufacturer's recommended pressures For many nozzle types lower pressure produces larger droplets When higher flow rates are needed use higher flow rate nozzles instead of increasing pressure

**Number of nozzles** Use the minimum number of nozzles that provide uniform coverage

**Nozzle orientation** Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice Significant deflection from horizontal will reduce droplet size and increase drift potential

**Nozzle type** Use a nozzle type that is designed for the intended application With most nozzle types narrower spray angles produce larger droplets Consider using low drift nozzles Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift potential

#### BOOM LENGTH

For some use patterns reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width

#### APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind

#### SWATH ADJUSTMENT

When applications are made with a crosswind the swath will be displaced downwind Therefore on the up and downwind edges of the field the applicator must compensate for this displacement by adjusting the path of the aircraft upwind Swath adjustment distance should increase with increasing drift potential (higher wind small drops etc )

#### WIND

Drift potential is lowest between wind speeds of 2-10 mph However many factors including droplet size and equipment type determine drift potential at any given speed Application should be avoided below 2 mph due to variable wind direction and high inversion potential NOTE Local terrain can influence wind patterns Every applicator should be familiar with local wind patterns and how they affect spray drift

#### TEMPERATURE AND HUMIDITY

When making applications in low relative humidity set up equipment to produce larger droplets to compensate for evaporation Droplet evaporation is most severe when conditions are both hot and dry

#### TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high Temperature inversions restrict vertical air mixing which causes small suspended droplets to remain in a concentrated cloud This cloud can move in unpredictable directions due to the light variable winds common during inversions Temperature inversions are characterized by increasing temperatures with altitude and are common on

nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS**

Application through sprinkler irrigation systems in CA is prohibited.

Apply this product only through sprinkler irrigation system(s) including center pivot. DO NOT apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

**DO NOT** apply this product through irrigation systems connected to a public water system. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments should the need arise.

The irrigation water pipeline must be fitted with a functional automatic quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low-pressure drain located between the irrigation water pump and the check valve to prevent back-siphoning of treated irrigation water into the water source.

Always inject Iprodione ETQ into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional normally closed solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.



The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected

Spray mixture in the chemical supply tank must be agitated at all times otherwise settling and uneven application may occur DO NOT apply when wind speed favors drift beyond the area intended for treatment

For injection of pesticides a metering pump such as a positive displacement injection pump of either diaphragm or piston type constructed of materials that are compatible with pesticides fitted with a system interlock and capable of injection at pressures approximately 2 to 3 times those encountered within the irrigation water line must be used

## TURF

### **Golf Courses, Sod Farms and Institutional Areas**

**Do not** prepare more solution than can be used in 12 hours to minimize potential degradation of active ingredient

Initiate applications when disease presence is detected or if weather conditions favor disease development

Under severe disease pressure apply the highest rate at the shortest interval of applications for all diseases For light to moderate disease pressure apply the lowest rate at the longest interval

Apply the rates indicated in the table in 0.5 - 10 gallons of water per 1000 sq ft

**Do not** drench the foliage to the point of runoff

**Do not** apply more than 35 fl oz of this product per 1000 sq ft per year (24 lbs a.i. per acre)

**Do not** make more than 6 applications per year

**Do not** mix with any sticker extender or wetting agent

**Do not** mow or irrigate treated areas until the foliage is completely dry Wait 24 hours following treatment

**Do not** graze animals on treated turf and do not feed clippings from treated turf to livestock or poultry

### Application Rates and Directions

TARGET PEST	RATE (fl oz /1000 sq ft )	APPLICATION DIRECTIONS [Retreatment Interval (days)]
Dollar spot ( <i>Lanzia spp and Moellerodiscus spp</i> )	3 4	Greens & tees [14 21]
Brown patch ( <i>Rhizoctonia solani</i> )	2 4	Fairways & other turf areas [14 28]
Leaf spot ( <i>Drechslera spp</i> )		
Large patch* ( <i>Rhizoctonia solani</i> )	4	Make first application in fall when conditions favor disease development but no symptoms are visible [14 21]
Fusarium blight ( <i>Fusarium spp</i> )	8	Use only preventative foliar application when conditions first become favorable for disease development [28 minimum]
Necrotic ring spot* ( <i>Leptosphaeria korrae</i> )		
Fusarium patch ( <i>Microdochium nivalis</i> ) [Pacific Northwest Only West of the Cascade Mountains]	4 8	[14 21]
Gray snow mold ( <i>Typhula spp</i> )	4 8 (see tank mixes for additional information)	Make one application before first permanent snow cover and a second during a mid winter thaw
Pink snow mold ( <i>Fusarium nivale</i> )		
Corticium read thread ( <i>Laetisaria fuciformosis</i> )	4	As required for prevention [14 minimum]
Curvularia ( <i>Curvularia spp</i> ) [Burmegrass only]	4	As required for prevention [14 minimum]
Anthracnose ( <i>Colletotrichum</i> )	4 8	Combine this product with appropriately labeled and registered trifloxystrobin or fosetyl al products or other anthracnose control fungicides
Note suppression only		
Pythium blight	See tank mixes below	

\* Not registered for this use in California

### TANK MIXTURES FOR TURF APPLICATIONS

To expand the spectrum of pests controlled tank mix this product with most commonly used fungicides containing flutolanil trifloxystrobin and azoxystrobin When tank mixing products be sure to follow the most restrictive instructions on all product labels Do not tank mix with any product that contains a prohibition on tank mixing

### Broad Spectrum Disease Control and Resistance Management

Tank mixing this product with an appropriately labeled and registered thiophanate methyl product provides effective broad spectrum turf disease control and also serves as a useful

tank mixture in the resistance management program required for other resistance sensitive fungicides

Disease Pressure	Product Rate (fl oz /1000 sq ft )	Thiophate Methyl Rate (fl oz /1000 sq ft )
Low to Medium	3	1.0
High	3	2.0

**Summer Stress Complex/Summer Decline**

Mix 2 to 4 oz per 1000 sq ft of this product with an appropriately labeled and registered fosetyl al containing product at the labeled rate

**Pythium Blight**

Pythium blight will be controlled by the tank mixing of fosetyl al or propamcarb hydrochloride with this product. If using a tank mixture follow label directions for the use of that product and apply at the specified rate for control of the target disease organism

**Gray Snow Mold**

In areas where continuous snow cover occurs use 4 to 8 fl oz of this product tank mixed with an appropriately labeled and registered chlorothalonil product at the labeled rate. Make applications in the fall before snow cover occurs and use the higher rate listed if the turf remains frozen prior to snow cover. Apply with 1 to 5 gallons of spray solution per 1000 sq ft. For best results reapply if loss of snow cover occurs during a winter thaw

**ORNAMENTALS**

**Field, Landscape and Greenhouse Ornamentals and Conifer Nurseries**

For Use in Commercial Nurseries only Not for use in Residential Areas

As it is not possible to test every species or variety of ornamental plant for tolerance the user should test for phytotoxic responses in plants not listed in this label prior to widespread application

This product has been tested on the following ornamentals

- |              |                     |                     |                        |
|--------------|---------------------|---------------------|------------------------|
| Ageratum     | Ajuga               | Almond (ornamental) | Alyssum                |
| Andromeda    | Aphelandra          | Artemisia           | Aster                  |
| Azalea       | Boxwood             | Cactus              | Calendula              |
| Carnation    | Cherry (ornamental) | Chrysanthemum       | Cineraria              |
| Cistena Plum | Coleus              | Columbine           | Coral Bells (Heuchera) |

Crape Myrtle	Crassula	Croton	Cyclamen
Daffodils	Dahlia	Delphinium	Deutzia
Dianthus	Dieffenbachia	Dizygotheca	Dogwood
Dracena	English Ivy	Episcia	Euonymous
Ficus	Forsythia	Gazania	Geranium
Gladiolus	Gloxinia	Gypsophila	Hawthorn
Holly	Hoya	Hydrangea	Impatiens
Iris	Juniper	Kalanchoe	Lilies
Lipstick Vine	Marigold	Monarda (Bee Bahm)	Pachysandra
Palm	Pansy	Peach (ornamental)	Peperomia
Periwinkle	Philodendron	Phlox	Pilea
Pine	Pitosporum	Plum (ornamental)	Poinsettia
Poppy	Pothos	Primrose	Privet
Protea	Pyracantha	Rhododendron	Rose
Rose Tree of China	Salvia	Schefflera	Snapdragon
Statice	Tree Ivy	Tulip	Viburnum
Violet	Zinnia		

**Note** Do not apply this product to Peace Lily or White Anthurium (*Spathiphyllum*)

Use the following table to determine the diseases controlled and the appropriate application method

Disease	Can be Applied to	Foliar Spray	Drench	Dip
Aerial web blight ( <i>Rhizoctonia spp</i> )	All	X		
Alternaria leaf blight ( <i>Alternaria euphorbiae</i> )	All	X		
Alternaria leaf spot ( <i>Alternaria panax Alternaria tenuissima</i> )	All	X		
Botrytis blight ( <i>Botrytis spp</i> )	All	X		
Fusarium leaf spot ( <i>Fusarium moniliforme</i> )	All	X		
Helminthosporium leaf spot ( <i>Helminthosporium spp</i> )	All	X		
Rhizoctonia stem and root rot ( <i>Rhizoctonia spp</i> )	All except Impatiens and Pothos		X	
Ink spot ( <i>Drechslera iridis</i> )	Iris	X		
Tulip fire ( <i>Botrytis tulipea</i> )	Tulip	X		
Alternaria leaf blight ( <i>Alternaria zinniae</i> )	Zinnia	X		
Ray blight ( <i>Ascochyta chrysanthami</i> )	Chrysanthemum	X		
Fusarium corm rot ( <i>Fusarium oxysporum</i> )	Gladiolus			^

Daffodil leaf scorch ( <i>Stagnospora curtissi</i> )	Daffodilis	X		
Blossom blight ( <i>Monolinia fructicola</i> )	Cistena Plum/Ornamental Plum	X		
Botrytis storage rot ( <i>Botrytis spp</i> )	Rose			X
Cylindrocladium blight and wilt ( <i>Cylindrocladium scoparium</i> )	Azalea and Rhododendron			X

### Foliar Spray Applications

Apply 1 to 2.5 quarts of this product in 100 gallons of water per acre every 7-14 days until disease pressure is within acceptable levels. Make no more than 4 applications per crop per year. Spray plants to the point of run off to ensure thorough coverage. Apply no more than 2.5 quarts of this product per acre per application. Apply no more than 10 quarts of this product per acre per year. Applications at the highest rate and shortest interval are appropriate for all diseases when conditions are severe. Applications at the lower rates and longer intervals are appropriate when disease pressure is light to moderate.

### Drench Applications

To control *Rhizoctonia* stem and root rot (*Rhizoctonia spp*) mix 13 fl oz in 100 gallons of water. Apply 1-2 pints of this solution per sq ft of soil at seeding or transplanting. Repeat application every 14 days as disease pressure warrants. Apply no more than 35 fl oz of this product per 1000 sq ft per year. Make no more than 6 applications per year. Applications at the highest rate and shortest interval are appropriate for all diseases when conditions are severe. Applications at the lower rates and longer intervals are appropriate when disease pressure is light to moderate.

**Do not** use this product as a drench on Impatiens and Pothos

### Dip Applications

For dip applications disperse 1 quart of this product in 100 gallons of water.

Roses To control Botrytis Storage Rot (*Botrytis spp*) dip bare root for 5 min prior to cold storage.

Azalea and Rhododendron To control Cylindrocladium Blight and Wilt\* (*Cylindrocladium scoparium*) dip cuttings for 5 min before planting.

Gladiolus To control Fusarium Corm Rot (*Fusarium oxysporum*) dip corms for 5 min prior to storage.

\* Not registered for this use in California

**STORAGE AND DISPOSAL**

Do not contaminate water food or feed by storage or disposal

**Pesticide Storage** Store in a cool dry place and in such a manner as to prevent cross contamination with other pesticides fertilizers food and feed Store in original container and out of the reach of children preferably in a locked storage area

**Pesticide Disposal** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

**Container Disposal Non refillable container** Do not reuse or refill this container Triple rinse or pressure rinse container (or equivalent) promptly after emptying Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container ¼ full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank for later use or disposal Drain for 10 seconds after flow begins to drip Repeat this procedure two more times Offer for recycling if available or puncture and dispose of in a sanitary landfill or incineration or if allowed by State and local authorities by burning If burned stay out of smoke

**WARRANTY AND LIMITATION OF DAMAGES**

**CONDITIONS OF SALE** To the extent consistent with applicable law Sipcam Agro USA Inc warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to Sipcam Agro USA Inc **SIPCAM AGRO USA, INC DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED** To the extent consistent with applicable law **SIPCAM AGRO USA, INC SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND SIPCAM AGRO USA, INC 'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE** To the extent consistent with applicable law, **BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT SIPCAM AGRO USA, INC DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT**

Initial Registration Application 3/6/12