



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

60063-55

Date of Issuance:

6/8/15

NOTICE OF PESTICIDE:

Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

CTL + Ipro Turf and Ornamental Fungicide

Name and Address of Registrant (include ZIP Code):

Sipcam Agro USA, Inc.  
Jennifer Yentel, Registration Manager  
2525 Meridian Parkway, Suite 350  
Durham, NC 27713

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Tony Kish, Product Manager 22  
Fungicide Branch, Registration Division (7505P)

Date:

6/8/15

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 60063-55.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF, dated 10/22/2014
- Alternate CSFs 1-9, dated 10/22/2014

If you have any questions, please contact Tony Kish by phone at 703 308-9443, or via email at [kish.tony@epa.gov](mailto:kish.tony@epa.gov); or Craig Reeves by phone at 703 347-0486, or via email at [reeves.craig@epa.gov](mailto:reeves.craig@epa.gov).

Sincerely,



Tony Kish, Product Manager 22  
Fungicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure: Stamped Accepted Label



**SIPCAM AGRO USA, INC.**

# CTL + Ipro Turf and Ornamental Fungicide

**Active Ingredients:**

Chlorothalonil: tetrachloroisophthalonitrile .....	28.0%
Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolinecarboxamide .....	14.0%
Other Ingredients: .....	58.0%
Total: .....	100.0%

Contains 3 pounds chlorothalonil per gallon.

Contains 1.5 pounds iprodione per gallon.

## Keep Out of Reach of Children DANGER – PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF INHALED	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF SWALLOWED	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 222-1222 Poison Control Center
NOTES TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Persons having a temporary allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.	


EPA Reg. No. 60063-xx

Net Contents: XXXXXXXXXX

EPA Est. No.: XXXXXXXXXX

See additional Precautionary Statements and Directions for Use inside label booklet.

Manufactured for:  
Sipcam Agro USA, Inc.  
2525 Meridian Parkway, Suite 350  
Durham, NC 27713



**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**DANGER – PELIGRO**

Corrosive. Causes irreversible eye damage. May be fatal if inhaled. Causes skin irritation. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**Note to User:** This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Person having allergic reaction should contact a physician.

**Personal Protective Equipment (PPE):**

Applicators, mixers, loaders and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made of waterproof material, such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyethylene, polyvinyl chloride, or viton; If you want more options follow the instructions for Category A on an EPA chemical-resistance category selection chart.
- Chemical-resistant footwear plus socks
- Protective eye-wear such as goggles, safety glasses, or face shield
- A dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC21C)
- Chemical-resistant headgear for overhead exposures
- A chemical resistant apron when cleaning equipment, mixing, or loading

**User Safety Requirements:**

Discard clothing or other materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls:**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to wildlife, fish, invertebrates, and marine/estuarine organisms. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This product can contaminate surface water through spray drift. Do not apply when weather conditions favor drift from treated areas. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not

separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

Chlorothalonil, one of the active ingredients in this product, is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Attention: This product contains a chemical known to the State of California to cause cancer.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read the entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval (REI). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the REI of 48 hours.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water) is:

Coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear.

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the restricted-entry interval expires after 12 hours, for the next 6 1/2 days entry is permitted only when the following safety measures are provided:

- (1) At least one container designed specifically for flushing eyes must be available in operating condition at the WPS-required decontamination site intended for workers entering the treated area.
- (2) Workers must be informed, in a manner that they can understand:
  - that residues in the treated area may be highly irritating to their eyes,
  - that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes,
  - that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site or using other readily available clean water, and
  - how to operate the eyeflush container.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter the treated areas until sprays have dried.

### General Use Precautions and Restrictions

- Use of this product at residential sites is prohibited.
- Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.
- For golf courses only, do not apply to turf cut higher than 1" on golf holes where water bodies are present.
- Do not apply this product when the wind direction is toward aquatic areas.
- Apply with ground equipment only.
- Do not apply more than 24 lbs. of iprodione active ingredient per acre annually.
- Do not make more than 6 applications of this product per year.
- Do not mix this product with any sticker, extender or wetting agents.

### Mixing, Loading, and Applying Instructions

Partially fill the spray tank with clean water. Measure the required amount of CTL + Ipro Turf and Ornamental Fungicide and pre-mix with a small volume of water then add this to the tank. Agitate to ensure thorough mixing while filling tank with remaining water. Maintain agitation during application and apply with properly calibrated application equipment. **Do not prepare more solution than can be used within a 12 hour period to minimize degradation of active ingredient, particularly in water with a high pH.** The spray solution should be buffered to a pH of 5.0 – 7.0. Mixing with very acidic products may result in precipitation of CTL + Ipro Turf and Ornamental Fungicide.

**If you are unsure about disease conditions, contact your local extension agent.**

### TURFGRASSES

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, recreational park athletic fields, athletic fields located on or next to school (e.g., elementary, middle, and high schools), campgrounds, churches, and theme parks.

Do not apply more than the following totals of chlorothalonil active ingredient from all registered product sources to the indicated types of turfgrass:

TYPE OF TURFGRASS	TOTAL CHLOROTHALONIL ACTIVE INGREDIENT PER ACRE PER YEAR
Golf Course Greens	73 lbs
Golf Course Tees	52 lbs
Golf Course Fairways	26 lbs
Other Turf	26 lbs

Do not drench the foliage to the point of runoff.

Do not mow or irrigate treated areas until the spray deposit on turfgrass is completely dry. Wait 24 hours following treatment.

Do not graze animals on treated turf and do not feed clippings from treated turf to livestock or poultry.

Initiate applications when disease presence is detected or if weather conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease development apply the highest rate at the shortest interval of applications for all diseases. For light to moderate disease pressure apply the lowest rate at the longest interval.

Apply the rates indicated in the table in 0.5 to 10 gallons of water per 1,000 sq. ft.

### Application Rates and Directions

Disease	Rates		Application Interval (days)
	Golf Course Greens and Tees (Rate/1000 sq. ft)	Golf Course Fairways, Lawns, & Other Turfgrasses (Rate/Acre)	
Dollar Spot ( <i>Sclerotinia homoeocarpa</i> ; <i>Lanzia spp.</i> and <i>Moellerodiscus spp.</i> )	4 – 6 fl. oz.	10 7/8 – 16 1/4 pt.	Golf course greens and tees: 14 - 21
Brown Patch ( <i>Rhizoctonia solani</i> , <i>R. zeae</i> , <i>R. cerealis</i> )			
Leaf spot, Melting-out, Brown blight ( <i>Drechslera spp.</i> (including <i>D. poae</i> , <i>D. siccans</i> ), <i>Bipolaris sorokiniana</i> , <i>Curvularia spp.</i> )	4 – 6 fl. oz.	10 7/8 – 16 1/4 pt.	Golf course fairways and other turf areas: 14 - 28
Large patch* ( <i>Rhizoctonia solanii</i> )	5 fl. oz.	13 5/8 pt.	14 - 21
	Make first application in fall when conditions favor disease development but no symptoms are visible.		
Gray leaf spot ( <i>Pyricularia grisea</i> , <i>P. oryzae</i> )	4 – 6 fl. oz.	10 7/8 – 16 1/4 pt.	14 - 28
Red thread ( <i>Laetisaria fuciformis</i> )	5 – 8 fl. oz.	13 5/8 – 21 3/4 pt.	14 minimum
Anthracnose ( <i>Colletotrichum graminicola</i> )	6 – 8 fl. oz.	16 1/4 - 21 3/4 pt.	14 minimum
Copper spot ( <i>Gloeocercospora sorghi</i> )	8 fl. oz.	21 3/4 pt.	14 minimum
Stem rust (bluegrass) ( <i>Puccinia graminis</i> )	8 fl. oz.	21 3/4 pt.	14 minimum
Dichondra leaf spot ( <i>Alternaria spp.</i> )	8 fl. oz.	21 3/4 pt.	14 minimum
Large patch* ( <i>Rhizoctonia solanii</i> )	4 – 7 fl. oz.	10 7/8 – 19 1/8 pt.	14 minimum
Fusarium blight ( <i>Fusarium spp.</i> ) Necrotic ring spot* ( <i>Leptosphaeria korrae</i> )	10 fl. oz.	27 1/8 pt.	28 minimum
	Use only preventative foliar application when conditions first become favorable for disease development.		
Fusarium patch ( <i>Microdochium nivalis</i> ) [Pacific Northwest Only – West of the Cascade Mountains]	8 – 10 fl. oz.	21 3/4 - 27 1/8 pt.	21 - 28
Gray snow mold ( <i>Typhula spp.</i> ) Pink snow mold ( <i>Fusarium nivale</i> )	8 – 10 fl. oz.	21 3/4 - 27 1/8 pt.	Make one application before first permanent snow cover and a second during a mid-winter thaw.
Corticium read thread ( <i>Laetisaria fuciformis</i> )	5.5 fl. oz.	14 7/8 pt.	14 minimum
	Make applications as required for prevention.		
Curvularia ( <i>Curvularia spp.</i> ) [Bermudagrass only]	5.5 fl. oz.	14 7/8 pt.	14 minimum
	Make applications as required for prevention.		

\*Not registered for use in California

### ORNAMENTALS

For field, landscape and greenhouse ornamentals.

For use in commercial nurseries only. Do not use in residential areas.  
Do not apply to ornamentals which can bear harvestable fruit or nuts.

Fruits and other structures which may be borne on treated plants **MUST NOT BE EATEN.**

This product may be used in greenhouses. Do not use mistblowers or high pressure spray equipment when making applications of this product in greenhouses.

CTL + Ipro Turf and Ornamental Fungicide is recommended for use on the ornamentals listed in the table below for control of fungal diseases referred to by numbers in the Diseases Controlled column. As it is not possible to test every species or variety of ornamental plant for tolerance, the user should test this product for phytotoxic responses, using the listed rates on each type of ornamental plant on a small area prior to widespread use. Applications made during bloom may damage flowers and/or fruits.

Apply CTL + Ipro Turf and Ornamental Fungicide at a rate of 2 3/4 pints per 100 gallons of water per acre unless other directions are given in the tables below. Apply enough diluted spray per acre to provide through coverage of all plants that are intended to be protected from disease, generally ranging from 20 to 150 gallons per acre. Repeat applications at 7 to 14 day intervals until conditions are no longer favorable for disease. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, apply CTL + Ipro Turf and Ornamental Fungicide at 7 day intervals.

Do not apply more than 36.4 lbs. of chlorothalonil active ingredient per acre per year on field-grown ornamentals.

Do not apply more than 24 lbs. of iprodione active ingredient per acre per year on ornamental trees and plants with a maximum of 6 applications.

<b>Broadleaf Shrubs and Trees</b>			
	<b>Diseases Controlled</b>		<b>Diseases Controlled</b>
Almond (non-bearing)	1, 2	Holly	1
Andromeda (Pieris)	3	Peach (non-bearing)	1, 2
Azalea	1, 2, 3	Plum (non-bearing)	1, 2
Cherry (non-bearing)	1, 2	Privet (Ligustrum)	1
Dogwood	1	Rhododendron	1, 2, 3
Euonymus	1	Viburnum	4
Hawthorne	1		
<b>Flowering<sup>1</sup> Plants and Bulbs</b>			
	<b>Diseases Controlled</b>		<b>Diseases Controlled</b>
Arabian Violet	2	Marigold	1
Carnation	1, 2	Pansy	1
Chrysanthemum	1, 2	Phlox	1
Daffodil	1	Poinsettia <sup>2</sup>	1
Geranium	1, 5	Rose <sup>3</sup>	1
Gladiolus	1, 2	Statice	1
Hydrangea (foliage only)	1, 5	Tulip	1
Iris	1, 2	Zinnia	1, 4
Lily	1		
<sup>1</sup> Avoid application during bloom period on plants where flower injury is unacceptable.			
<sup>2</sup> Discontinue applications prior to bract formation; phytotoxicity is possible on the bracts.			
<sup>3</sup> Use 2 pints of CTL + Ipro Turf and Ornamental Fungicide per 100 gallons of water.			
<b>Foliage Plants</b>			
	<b>Diseases Controlled</b>		<b>Diseases Controlled</b>
Artemesia	1	Pachysandra <sup>4</sup>	1
Dieffenbachia	1	Parlor palm	1
Dracaena	1	Peperomia	1
Ficus (non-bearing)	1	Philodendron	1, 3



Lipstick plant	1	
<sup>4</sup> Use 5 ½ pints of CTL + Ipro Turf and Ornamental Fungicide per 100 gallons of water.		

NOTE: Do not apply this product to Peace Lily or White Anthurium (Spathiphyllum).

<b>1. Leafspots and Foliar Blights:</b>	
Actinopelte leafspot Alternaria leafspot or leaf blight Anthracnose (Gnomnia, Glomerella, Colletotrichum, Driscula) blights Black spot (Diplocarpon) Botrytis blights Cephalosporium leafspot Cercospora leafspot Shothole (Stigmata) Corynespora stem & leafspot Curvularia leafspot Dactylaria leafspot Didymellina leafspot Drechslera (Bipolaris) leafspots, inkspot Fabraea (Entomosporium) leafspot Fusarium (Gibberella) leafspot	Gloeosporium black leafspot Helminthosporium leaf spot (Helminthosporium spp.) Marssonina leafspot Monilinia blossom blight, twig blight Mycosphaerella ray blight Myrothecium leafspot, brown rot Phyllosticta leafspot Ramularia leafspot Rhizoctonia web blight Scab (Venturia) Septoria leafspot Sphaeropsis leafspot Stagonospora leaf scorch Tan leafspot (Curvularia) Volutella leaf blight
<b>2. Flower Spots and Blights:</b>	
Botrytis flower spot, flower blight Curvularia flower spot Monilinia blossom blight	Ovulinia flower blight Rhizopus blossom blight Sclerotinia flower blight
<b>3. Phytophthora leaf blight, dieback</b>	
<b>4. Powder mildews:</b>	
Erysiphe cichoracearum Sphaerotheca fuliginea	Microsphaera spp.
<b>5. Rusts:</b>	
Gymnosporangium spp. Pucciniastrum hydrangea	Puccinia spp.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross-contamination with other pesticides, fertilizers, food, and feed.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:**

**Containers < 5 Gallons:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Containers ≥ 5 Gallons:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents

into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Bulk Containers:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When the container is empty, replace the cap and seal all openings that have been opened during use; and return to the point of purchase, or to a designated location named at the time of purchase of this product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged or leaking, call CHEMTREC at (800) 424-9300. If the container is damaged and leaking or material has been spilled, follow these procedures:

- Cover spill with absorbent material.
- Sweep into disposal container.
- Wash area with detergent and water and follow with clean water rinse.
- Do not allow to contaminate water supplies.
- Dispose of according to instructions.

If not returned to the point of purchase or to a designated location, clean empty container as instructed above and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

#### **WARRANTY AND LIMITATION OF DAMAGES**

**CONDITIONS OF SALE:** To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. **SIPCAM AGRO USA, INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED.** To the extent consistent with applicable law, **SIPCAM AGRO USA, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND SIPCAM AGRO USA, INC.'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. SIPCAM AGRO USA, INC. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.**

Chlorothalonil + Iprodione (EPA Reg. No. 60063-xx) (Application to Register to EPA 10-22-14) (REV 06-04-2015) (REV 06-08-2015)