



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 20, 2020

Patricia McFadden
Registration Manager
Sipcam Agro USA, Inc.
2525 Meridian Parkway
Durham, NC 27713

Subject: Label Amendment – Removal of CDPR turf restrictions optional language; Making CDPR ornamental and conifer restrictions optional language; Addition of turf diseases; Addition of application rate tables for ornamentals; Creation of crop table for christmas trees; Incorporation of ID mitigation language; Updated use site restrictions and crop group names; Inclusion of equivalent active ingredient rates for product rates; Minor label changes
Product Name: Azoxystrobin 250 SC
EPA Registration Number: 60063-59
Application Dates: June 17, 2019 and August 31, 2020
Decision Numbers: 566708 and 566435

Dear Ms. McFadden:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false

Page 2 of 2
EPA Reg. No. 60063-59
Decision No. 566708 and 566435

or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Lindsay DeMers via email at demers.lindsay@epa.gov.

Sincerely,

A handwritten signature in black ink, reading "Shaja B. Joyner". The signature is written in a cursive style with a large initial "S".

Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Enclosure



Azoxystrobin	Group	11	Fungicide
--------------	-------	-----------	-----------

Azoxystrobin 250 SC (Alternate Brand Name: Endow 2SC, Arius 250)

[For control of listed diseases on [Almonds;] [Artichoke, globe;] [Asparagus;] [Barley;] [Berry, low growing subgroup 13-07G (except cranberry);] [Brassica head & stem subgroup;] [Brassica, leafy greens subgroup;] [Bulb vegetables crop group 3-07;] [Carrots;] [Corn, field, pop, sweet (includes seed production);] [Cucurbits;] [Fruiting vegetables crop group 8-10;] [Grapes and other small fruit vine climbing subgroup 13-07F (except fuzzy kiwifruit);] [Herbs & spices (except black pepper), crop group 19;] [Leafy vegetables (except brassica);] [Legume vegetables, dry & succulent and legume vegetables, foliage of any cultivar of bean (*phaseolus* spp.) and field pea (*pisium* spp.);] [Peanuts;] [Pecans;] [Pistachios;] [Potatoes;] [Rice;] [Soybean;] [Stone fruit;] [Tomatoes;] [Tree nuts;] [Vegetables, leaves of root and tuber group and root subgroup;] [Vegetables, tuberous and corm subgroup;] [Wheat and triticale;] [Wild rice;] [Turf;] [Golf courses;] [Lawns and landscape areas around residential, institutional, public, commercial and industrial buildings;] [Parks;] [Recreational areas;] [Athletic fields;] [Ornamentals;] [Conifers including Christmas trees;] [and] [Roses, commercial rose production]]]

Active Ingredient:

Azoxystrobin: methyl (E)-2-{2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl}-3-methoxyacrylate* 22.93%

Other Ingredients: 77.07%

Total: 100.0%

*IUPAC

Contains 2.08 lbs. Azoxystrobin per gallon.

Keep Out of Reach of Children WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 222-1222 Poison Control Center

Net Contents: ___ [gallons] [gal.] [(Liters)]
EPA Est. No.: _____
Lot number [begins with] [ends with] xx]

EPA Reg. No. 60063-59
[lot number / label date code]

Read the [entire] label [carefully] before [opening the container] [using this product.]
See additional Precautionary Statements and Directions for Use inside booklet.

Manufactured for:
Sipcam Agro USA, Inc.
2525 Meridian Parkway
Durham, NC 27713

Optional Language for label

[Pull][Peel] back [book] [label] here]
[Application Type [AG] [Agriculture]]
[Application Type T[/O] [Turf] [& Ornamental]]
[Formulated in the United States of America, with U.S. and imported ingredients.]
[Product of _____] [Note: if manufactured in a country other than U.S., country name will appear here]
[Fungicide]

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Warning. May be fatal if swallowed. Harmful if absorbed through the skin. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Human flagging is prohibited.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, or viton \geq 14 mils
- In addition, mixers/loaders/applicators using mechanically pressurized handwands except when applying to Christmas tree farms, nursery ornamentals, landscaping, must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

Respirator fit testing, medical qualification, and training

Using a program that conforms to OSHA's requirements (see 29 CFR Part), employers must verify that any handler who uses a respirator is:

- Fit-tested and fit-checked,
- Trained, and
- Examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn. A qualified medical practitioner is a physician or other licensed health care professional who will evaluate the ability of a worker to wear a respirator. The initial evaluation consists of a questionnaire that asks about medical conditions (such as a heart condition) that would be problematic for respirator use. If concerns are identified, then additional evaluations, such as a physical exam, might be necessary. The initial evaluation must be done before respirator use begins. Handlers must be reexamined by a qualified medical practitioner if their health status or respirator style or use conditions change.

Upon request by local/state/federal/tribal enforcement personnel, employers must provide documentation demonstrating how they have complied with these requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

Azoxystrobin is toxic to freshwater and estuarine/marine fish and aquatic invertebrates. Azoxystrobin can be persistent for several months or longer.

DO NOT discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. **DO NOT** discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

For terrestrial uses **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

Azoxystrobin and a degradate of azoxystrobin are known to leach through soil to ground water under certain conditions as a result of label use. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of azoxystrobin and a degradate of azoxystrobin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Sipcam Agro USA, Inc. immediately if you observe any adverse environmental effects due to use of this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use of this product through airblast application equipment on grapes is prohibited in the following townships and boroughs of Erie County, Pennsylvania: North East, Harborcreek, Lawrence Park, Erie, Presque Isle, Millcreek, Fairview, Girard, and Springfield.

This prohibition is intended to help eliminate phytotoxicity problems with apples observed in this geographic location.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval and notification to workers. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area being treated must be vacated by unprotected persons.

DO NOT treat while unprotected humans or domestic animals are present in the treatment areas. Because certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information.

DO NOT allow entry into treatment area until the area that was treated with this product is dried.

PRODUCT INFORMATION

This product is a broad spectrum, preventative fungicide with systemic and curative properties. This product may be applied as a foliar spray in alternating spray programs or in tank mixes with other registered crop protection products. All applications must be made according to the use directions that follow.

RESTRICTIONS

- **DO NOT** graze or feed clippings from treated turf areas to animals.
- This product is extremely phytotoxic to certain apple varieties. **AVOID SPRAY DRIFT**. Extreme care must be used to prevent injury to apple trees and apple fruit. **DO NOT** spray this product where spray drift may reach apple trees. **DO NOT** use spray equipment which has been previously used to apply this product to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.
- Aerial and/or chemigation application to sod is prohibited.

SPRAY DRIFT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft. above the ground or crop canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzles that deliver Medium to coarse spray droplets in accordance with ASABE Standard S-572.1.
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

Groundboom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Airblast Applications:

- Sprays must be directed into the canopy.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer rows.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT - Ground Boom

- For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

- Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

- When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

- Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

- Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.
- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

MIXING, LOADING AND APPLYING

This product is intended to be diluted into water and then applied to crops by typical agricultural spraying techniques. Always apply this product in sufficient water to obtain thorough, uniform coverage of foliage and crop surfaces intended to be protected from disease. Spray volume to be used will vary with crop and amount of plant growth. Spray volume must normally range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute sprays and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are acceptable unless specific directions are given for a crop.

Slowly invert container several times to assure uniform mixture. Measure the required amount of this product and pour into the spray tank during filling. Keep agitator running when filling spray tank and during spray operations.

It is necessary to thoroughly apply the product in order to provide good disease control. **DO NOT** prepare more spray solution than is needed for application. Avoiding spray overlap will reduce the potential for crop injury.

Tank Mixing

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

When tank mixing this product with other pesticides, observe the more restrictive label limitations and precautions. **DO NOT** exceed any label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

DO NOT combine this product in the sprayer tank with pesticides, surfactants or fertilizers, unless prior use has shown the combination to be physically compatible, non-injurious and effective under similar use conditions. **DO NOT** combine this product with other products that contain *Bacillus thuringiensis* subsp. *kurstaki* strain ABTS-351, as the combination may result in phytotoxicity when applied to the crops listed on this label. **DO NOT** tank mix this product with oil or with any adjuvants which contain oil as their principal ingredient.

When mixed with EC (emulsifiable concentrate) formulations, this product may be phytotoxic to other crops listed on this label, especially when applied during cool, cloudy conditions that last for several days. Adjuvants containing

silicone could also have phytotoxic effects. When an adjuvant is used with this product, use a Council of Producers and Distributors of Agrotechnology (CPDA) certified adjuvant.

Applications through Sprinkler Irrigation Systems (Chemigation)

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set and portable (wheel move, side roll, end tow, or hand move) irrigation system(s). **DO NOT** apply this product through any other type of irrigation system. Use only on crops specifically designated in the **Crop Use Directions**.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

DO NOT apply this product through irrigation systems connected to a public water system. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation must be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject this product into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

This product may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a metering pump, such as a positive displacement injection pump of either diaphragm or piston type, constructed of materials that are compatible with pesticides, fitted with a system interlock, and capable of injection at pressures approximately 2 to 3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Fill chemical supply tank of injection equipment with water. Operate system for one complete revolution or run across the field, measuring time required, amount of water injected, and acreage covered. Thoroughly mix the specified amount of this product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously

agitated during the injection run. Shut off injection equipment after one revolution or run but continue to operate irrigation system until this product has been cleared from the last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a 30 – 45 minute period. Mix desired amount of this product for acreage to be covered with water so that the total mixture of this plus water in the injection tank is equal to the quantity of water used during calibration and operate entire system at normal pressures specified by the manufacturer of injection equipment used for the amount of time established during calibration. No agitation should be required. This product can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until this product has been cleared from the last sprinkler head.

INTEGRATED PEST / DISEASE MANAGEMENT

This product provides excellent control of fungal diseases when used according to label directions for control of a broad spectrum of plant diseases. This product is for use in programs that are compatible with the principals of Integrated Pest Management (IPM), including the use of disease resistant crop varieties, cultural practices, pest scouting, and disease forecasting systems which reduce unnecessary applications of pesticides.

RESISTANCE MANAGEMENT

For resistance management, Azoxystrobin 250 SC contains azoxystrobin, a QoI Group 11 fungicide. Any fungal population may contain individuals naturally resistant to Azoxystrobin 250 SC and other QoI Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Fungal isolates with acquired resistance to Group 11 may eventually dominate the fungal population if Group 11 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. Cross resistance has been shown between all members of the QoI fungicides. Since QoI fungicides are a high risk for resistance, this may result in partial or total loss of control of those species.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 11 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance - management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your local Sipcam Agro representative. You can also contact your pesticide distributor or university extension specialist to report resistance.

Follow the crop specific resistance management guidance listed in the Crop Use Directions table. If resistance management guidance is not specified in the Crop Use Directions table, then follow the guidance provided in the table below.

Total fungicide applications planned per crop	1	2	3	4	5	6	7	8	9	10	11	12
Applications of QoI fungicides applied alone	1	1	2	2	2	2	2	3	3	3	3	4
Applications of QoI fungicides applied in mixture (tank-mix or formulated)	1	2	2	2	2	3	3	4	4	5	5	6

When multiple applications are required during the year, spray programs for Group 11 (QoI) fungicides must be developed. When two sequential applications of Group 11 fungicides are made, they must be alternated with two or more applications of a fungicide that is not a Group 11 fungicide. If more than 12 applications are made during the year, observe these guidelines:

- When applying Group 11 (QoI) fungicides alone, the number of applications must not exceed more than 1/3 of the total number of fungicide applications per year.
- When applying Group 11 (QoI) fungicides in tank mixes or premixes with mixing partners of different modes of action, the number of QoI containing applications must not exceed more than 1/2 of the total number of fungicide applications per year.
- When applying Group 11 (QoI) fungicides both alone and in mixtures, the number of QoI containing applications must not exceed 50% of the total number of fungicide applications per year.

When applying a Group 11 fungicide to seed or soil, wait at least 3 weeks before making another application with a Group 11 fungicide.

SOILBORNE/SEEDLING DISEASE CONTROL

If applied early in the growing season, this product provides control of many soilborne diseases for those crops that list application directions for soilborne disease control. To control of pre- or post-emergence damping off and diseases that infect plants and the soil-plant interface, apply this product either in-furrow or as banded applications over the row, shortly after seedling emergence or during herbicides application or cultivation.

Regional cultural practices determine the application type used. The success of application types varies by region and depends on the timing and scope of the disease. In-furrow applications provide seedling disease control while banded applications are more effective at controlling soilborne diseases that develop later in the year. Consult your local extension agent for guidance on best application type for your situation.

Crop injury can occur when this product is applied as a soil directed application during cool, wet conditions.

Banded Applications

- Apply this product as a directed spray to the soil, prior to infection. Use single or multiple nozzles to provide thorough coverage of lower stems and soil surface surrounding the plants.
- Limit band width to 7 inches or less.
- Apply this product at a rate of 0.40 – 0.80 fl. oz. product. (0.10 – 0.20 oz. a.i.)/1000 row feet. For banded applications on 22-inch rows, the maximum application rate is 0.70 fl. oz/1000 row feet.
- Since banded applications come into contact with the foliage, they are considered to be foliar applications when following resistance management guidance.
- Make banded applications during cultivation or hilling operations to provide soil incorporation.

In-furrow Applications

- Apply this product as an in-furrow spray in 3-15 gallons of water at planting.
- Mount the spray nozzle so the spray is directed into the furrow just before the seeds are covered.
- Use the higher rate when the weather conditions are expected to be conducive to disease development, if the field has a history of *Pythium* problems, or if minimum/low till programs are in place.

Application Rates								
Rate per 1000 row-feet		Row Spacing (inches)						
fl. oz. product	oz. AI	22"	30"	32"	34"	36"	38"	40"
		rows	rows	rows	rows	rows	rows	rows
Product per acre (fl. oz.)								
0.40	0.10	9.5	7.0	6.5	6.1	5.8	5.5	5.2
0.60	0.15	14.3	10.5	9.8	9.2	8.7	8.3	7.8
0.80	0.20		13.9	13.1	12.3	11.6	11.0	10.5
1.00	0.25					14.5	13.8	13.1

Row-feet per acre: 22" = 23,760 row ft.; 30" = 17,424 row ft.; 32" = 16,335 row ft.; 34" = 15,374 row ft.; 36" = 14,520 row ft.; 38" = 13,756 row ft.; 40" = 13,068 row ft.

DO NOT apply more than 15 fl. oz. (0.24 lb. AI) of this product per acre.

Drip

Refer to the **Applications through Sprinkler Irrigation Systems (Chemigation)** section of this label.

Crop Rotation Restrictions

Refer to the table below for the minimum time intervals required between the last application of this product and a new crop planting.

Crop	Rotational Interval (in days)
Buckwheat, millet	12 months
All other crops with azoxystrobin registered uses	0 days

CROP USE DIRECTIONS

During conditions which are favorable to prolonged periods of fungal infection use another registered fungicide for additional applications if maximum amount of this product has been applied. Efficacy for certain diseases may be reduced if resistant isolates to Group 11 fungicides are present. Use this product in an IPM program, alternating fungicides with different modes of action. Use the table in the “Resistance Management” section of this label to determine the number of applications of this product that can be made before alternating with fungicides with a mode of action other than QoI Group 11. When environmental conditions are favorable to disease, during period of heavy disease pressure, or with highly susceptible varieties, use the higher listed rates in the rate range and/or shorter spray intervals.

ALMONDS

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Alternaria Leaf and Fruit Spot <i>(Alternaria alternata)</i> Anthracnose <i>(Colletotrichum acutatum)</i> Brown Rot Blossom Blight <i>(Monilinia laxa, M. fructicola)</i> Leaf Blight (<i>Seimatosporium lichenicola</i>) Leaf Rust (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium carpophilum</i>) Shot Hole <i>(Wilsonomyces carpophilus)</i>	12.0 - 15.5 (0.20 - 0.25)	<p>Apply this product prior to disease outbreak or when conditions are favorable to disease development. Continue applications throughout the season following the resistance management guidelines</p> <p>Apply this product by ground in adequate water to provide complete coverage, by air in a minimum of 15 gallons of water per acre or by chemigation. Apply this product by air only at growth stages prior to and including 5 weeks after petal fall. If an adjuvant is used, add it at the manufacturer's specified rates.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p> <p>Anthracnose, scab and shot hole: Make first application prior to disease outbreak or when conditions are favorable to disease development. Make a second application 7 to 14 days after the first application, depending on the severity of disease pressure.</p> <p>Blossom blight: Make first application at early bloom and continue through petal fall.</p>
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 28 days 		

ARTICHOKE, GLOBE

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Ramularia Leaf Spot (<i>Ramularia cynarae</i>)	11.0 – 15.5 (0.18 – 0.25)	<p>Apply this product before disease outbreak or in the early stages of diseases. If environmental conditions are favorable to continued disease development, continue applications at 14- to 21-day intervals until harvest day.</p> <p>Apply this product by ground in 50-200 gallons of water per acre, by air in a minimum of 15 gallons of water per acre, or by chemigation.</p> <p>DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 0 days 		

ASPARAGUS

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Stemphyllium Purple Spot (<i>Stemphyllium vesicarium</i>)	6.0 – 15.5 (0.10 – 0.25)	<p>Apply this product before disease outbreak. If environmental conditions are favorable to disease development, make additional applications at 7- to 14-day intervals.</p> <p>Apply this product by ground in a minimum of 10 gallons of water per acre, by air in a minimum of 3 gallons of water per acre, or by chemigation.</p> <p>DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 100 days 		

BARLEY

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Kernel Blight or Black Point (<i>Alternaria</i> spp., <i>Cochiobolus sativus</i>) Leaf Rust (<i>Puccinia hordei</i> , <i>P. recondita</i>)	6.0 - 12.0 (0.10-0.20)	Apply this product before disease outbreak and until inflorescence has fully emerged.
Barley Stripe (<i>Drechslera graminea</i> = <i>Pyrenophora graminea</i>)	9.0 - 12.0 (0.15-0.20)	Apply this product by ground, air, or chemigation. If using a crop oil concentrate adjuvant, add at 1.0% v/v to optimize efficacy.
Powdery Mildew (<i>Erysiphe graminis f. sp. hordei</i>) Stagonospora Blotch (<i>Stagonospora nodorum</i>)	12.0 (0.20)	DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply after Feekes 10.54. • DO NOT apply more than 12 fl. oz. (0.20 lbs. AI) of this product per acre per application. • DO NOT apply more than 24 fl. oz. (0.40 lbs. AI) of this product per acre per year. • DO NOT make more than two (2) applications of this product at the high rate (0.20 lbs. AI / acre) per year. • Minimum Retreatment Interval: 14 days • DO NOT apply within 7 days of grazing or harvest (7-day PHI) for forage and hay. 		

BERRY, LOW GROWING SUBGROUP 13-07G (EXCEPT CRANBERRY)

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Anthracnose <i>(Colletotrichum fragariae)</i> Leather Rot <i>(Phytophthora cactorum)</i> Powdery Mildew <i>(Sphaerotheca macularis)</i> Suppression of Botrytis on Foliage <i>(Botrytis cinerea)</i>	6.0–15.5 (0.10 – 0.25)	Apply this product before disease outbreak. If environmental conditions are favorable to disease development, make additional applications at 7- to 10-day intervals throughout the year. Apply this product by ground, air, or chemigation. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11. Leather Rot: Make 2 applications on a 7-day schedule from late bloom through harvest. Field Nurseries: Apply to young plants in field nurseries by ground, drip, or overhead chemigation. If applying through drip irrigation, calculate the rate as a band application with a band width equal to the root zone width. Inject this product into the irrigation water. Root and crown rot suppression <i>(Colletotrichum spp.)</i> , at transplanting for commercial berry production. Dip plants for 2-5 minutes in a mix of 5-8 fl.oz. of this product per 100 gallons of water. Before dipping, remove excess soil. Treated plants must be transplanted as quickly as possible. Follow with foliar applications beginning 2-3 weeks after transplant.
Soilborne Diseases: Seedling Root Rot, Basal Stem Rot <i>(Rhizoctonia solani)</i>	0.40 – 0.80 fl. oz. per 1000 row feet	See directions and rates under the Soilborne/Seedling Disease Control section of this label.
CROP LIST: Bearberry, Bilberry, Cloudberry, Muntries, Partridgeberry, Strawberry, cultivars, varieties and/or hybrids of these.		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 61.5 fl. oz. (1.0 lbs. AI) of this product per acre per year. • DO NOT make more than four (4) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • DO NOT use in plant propagation nurseries. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 0 days 		

BRASSICA HEAD & STEM SUBGROUP 5A

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Alternaria Leaf Spot (<i>Alternaria</i> spp.) Downy Mildew (<i>Peronospora parasitica</i>) Pin Rot (<i>Alternaria</i> spp.)	6.0 - 15.5 (0.10 - 0.25)	Apply this product before disease outbreak. If environmental conditions are favorable to disease development, make additional applications at 7- to 14-day intervals throughout the year. Apply this product by ground in a minimum of 10 gallons of water per acre, by air in a minimum of 3 gallons of water per acre, or by chemigation. If an adjuvant is used, add it at the manufacturer's specified rates. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than Qol Group 11.
CROP LIST: Broccoli; Chinese broccoli (gai lan); Brussels sprouts; Cabbage; Chinese cabbage (napa); Chinese mustard (gai choy); Cauliflower; Cavalo broccolo; Kohlrabi; cultivars, varieties and/or hybrids of these		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of product per acre per year • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 0 days 		

BRASSICA, LEAFY GREENS SUBGROUP 5B

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Alternaria Leaf Spot (<i>Alternaria</i> spp.) Black Spot (<i>Alternaria</i> spp.) Cercospora Leaf Spot (<i>Cercospora</i> spp.) White Rust (<i>Albugo candida</i>)	6.0 - 15.5 (0.10 - 0.25)	Apply this product before disease outbreak. If environmental conditions are favorable to disease development, make additional applications at 7- to 14-day intervals throughout the year. Apply by ground, air or chemigation. If an adjuvant is used, add it at the manufacturer's specified rates. DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than Qol Group 11.
Soilborne Diseases Seedling Root Rot, Basal Stem Rot (<i>Rhizoctonia solani</i>)	0.40 - 0.80 fl. oz./ 1,000 row ft.	See directions and rates under the Soilborne/Seedling Disease Control section of this label.
CROP LIST: Broccoli raab; Cabbage, chinese; collards; kale; mizuna; mustard greens; mustard spinach; rape greens; cultivars, varieties and/or hybrids of these		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 46.5 fl. oz. (0.75 lbs. AI) of product per acre per year • DO NOT make more than three (3) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 0 days 		

BULB VEGETABLES CROP GROUP 3-07 (See list below)

Target Diseases	Rate fl. oz./Acre (lbs.ai/A)	Application Directions
Foliar Diseases Cladosporium Leaf Blotch (<i>Cladosporium allii</i>) Purple Blotch and Leaf Blight (<i>Alternaria porri</i>) (<i>Stemphylium vesicarium</i>) Rust (<i>Puccinia allii</i>)	6.0-12.0 (0.10-0.20)	Downy Mildew: Apply this product before disease outbreak on a 5- to 7-day interval. All other diseases: Apply this product before disease outbreak. If environmental conditions are favorable to disease development, make additional applications at 7- to 14-day intervals. Apply this product by ground, air or chemigation. Mixtures of this product with insecticides and silicone adjuvants must be tested for crop safety before application to the crop.
Botrytis Leaf Blight (<i>Botrytis aclada</i>) Downy Mildew (<i>Peronospora destructor</i>)	9.0 - 15.5 (0.15-0.25)	DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than QoI Group 11.
Soilborne Diseases Rhizoctonia Damping-Off (<i>Rhizoctonia solani</i>)	0.40 - 0.80 fl. oz./ 1,000 row ft.	See directions under the Soilborne/Seedling Disease Control section of this label. When making an in-furrow application, the spray must be made just prior to seeding in order to place the majority of the chemical under the seed. This practice reduces the potential for phytotoxicity, especially if fertilizer is added to the application.
CROP LIST: Chive, fresh leaves; chive, Chinese, fresh leaves; daylily, bulb; elegans hosta; fritillaria, bulb; fritillaria, leaves; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; kurrat; lady's leek; leek; leek, wild; lily, bulb; onion, Beltsville bunching; onion, bulb; onion, Chinese, bulb; onion, fresh; onion, green; onion, macrostem; onion, pearl; onion, potato, bulb; onion, tree, tops; onion, Welsh, tops; shallot, bulb; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of product per acre per year • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 5 days • Pre-harvest Interval (PHI): 0 days 		

CARROTS

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Early Blight (<i>Cercospora carotae</i>) Late Blight (<i>Alternaria dauci</i>) White Mold (<i>Sclerotium rolfsii</i>) Additional target diseases listed in the Vegetables, Root, subgroup	9.0 – 15.5 (0.15 – 0.25)	Apply this product before disease outbreak or when conditions are favorable to disease development. Apply every 7 to 14 days following resistance management practices. Apply the higher listed rate and shorter application intervals when disease pressure is severe. Apply this product by ground, air or chemigation. DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than QoI Group 11.
Soilborne Diseases Rhizoctonia Root Rot (<i>Rhizoctonia solani</i>)	0.40- 0.80 fl. oz. / 1,000 row ft.	See direction and rates under the Soilborne/Seedling Disease Control section of this label.
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 124 fl. oz. (2.0 lbs. AI) of this product per acre per year. • DO NOT make more than eight (8) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 0 days 		

CORN, FIELD, POP, SWEET (INCLUDES SEED PRODUCTION)

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Rust (<i>Puccinia sorghi</i>)	6.0 - 9.0 (0.10 – 0.15)	<p>Gray leaf spot: Apply this product when infection begins. Make a second application 14 days later if disease pressure persists.</p> <p>All other diseases: Apply this product before disease outbreak. If environmental conditions are favorable to disease development, make additional applications at 7- to 14-day intervals.</p> <p>Apply this product by ground, air or chemigation.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
Anthracnose Leaf Blight (<i>Colletotrichum graminicola</i>)		
Eye Spot (<i>Aureobasidium zeae</i>)		
Gray Leaf Spot (<i>Cercospora sorghi</i>)		
Northern Corn Leaf Blight (<i>Setosphaeria turcica</i>)	6.0 - 15.5 (0.10 - 0.25)	
Northern Corn Leaf Spot (<i>Cochliobolus carbonum</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>)		
Soilborne Diseases Rhizoctonia Root and Stalk Rot (<i>Rhizoctonia solani</i>)	0.40 - 0.80 fl. oz./ 1,000 row ft.	See directions and rates under the Soilborne/Seedling Disease Control section of this label.
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. (For rust, DO NOT apply more than 9.0 fl.oz. (0.15 lbs. AI) of this product per acre per application.) • For popcorn and sweet corn, DO NOT apply more than 124 fl. oz. (2.0 lbs. AI) of this product per acre per year. • For popcorn and sweet corn, DO NOT make more than eight (8) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • For field corn and field corn grown for seed, DO NOT apply more than 31 fl. oz. (0.5 lbs. AI) of this product per acre per year. • For field corn and field corn grown for seed, DO NOT make more than two applications per year at the high rate (0.25 lbs. AI). • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 7 days 		

CUCURBITS, CROP GROUP 9

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Alternaria Blight (<i>Alternaria cucumerina</i>) Anthracnose (<i>Colletotrichum lagenarium</i>) Belly Rot (<i>Rhizoctonia solani</i>) Cercospora Leaf Spot (<i>Cercospora citrulina</i>) Plectosporium blight (<i>Plectosporium tabacinum</i>) Downy Mildew (<i>Pseudoperonospora cubensis</i>) Gummy Stem Blight (<i>Didymella bryoniae</i>) Leaf Spots (<i>Alternaria</i> spp., <i>Cercospora</i> spp.) Myrothecium Canker (<i>Myrothecium roridum</i>) Plectosporium Blight (<i>Plectosporium tabacinum</i>) Powdery Mildew (<i>Sphaerotheca filiginea</i> , <i>Erysiphe cichoracearum</i>) Target Leaf Spot (<i>Corynespora cassicola</i>) Ulocladium Leaf Spot (<i>Ulocladium cucurbitae</i>)	12.0 – 15.5 (0.20 - 0.25)	<p>Downy and Powdery Mildew: Apply this product before disease outbreak or when conditions are favorable to disease development. Repeat at 5- to 7-day intervals for as long as conditions favor disease. Use the higher listed rate and a 5-day application interval when conditions are favorable to disease development.</p> <p>Belly rot control: Apply this product at the 1-3 leaf crop stage followed by a second application just prior to vine tip over or 10 to 14 days later, whichever occurs first.</p> <p>Other diseases: Apply this product before disease outbreak or when conditions are favorable to disease development. Repeat at 7- to 14-day intervals for as long as conditions favor disease. Use the higher listed rate and a 7 day application interval when conditions are favorable to disease development.</p> <p>Apply this product by ground, air or chemigation.</p> <p>DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
Soilborne Diseases Rhizoctonia Root Rot (<i>Rhizoctonia solani</i>)	0.40 - 0.80 fl. oz./ 1,000 row ft.	See directions and rates under the Soilborne / Seedling Disease Control section of this label.
CROP LIST: Cantaloupe; chayote; Chinese waxgourd; cucumber; gourds; honeydew; melons; <i>Momordica</i> spp. (bitter melon, balsam apple); muskmelon; watermelon; pumpkin; squash; zucchini; cultivars, varieties and/or hybrids of these		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 5 days • Pre-harvest Interval (PHI): 1 days • DO NOT tank mix this product with crop oil concentrates (COC), methylated spray oil (MSO), or silicon adjuvants. • DO NOT tank mix with products containing malathion, methomyl, chlorpyrifos, potassium salts of fatty acid, or dicloran. 		

FRUITING VEGETABLES CROP GROUP 8-10

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Anthracnose <i>(Colletotrichum spp.)</i> Powdery Mildew <i>(Sphaerotheca spp.)</i>	6.0 - 15.5 (0.10 - 0.25)	Apply this product before disease outbreak. If environmental conditions are favorable to disease development, make additional applications at 7- to 14-day intervals throughout the year. Apply this product by ground, air or chemigation. DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than Qol Group 11.
Soilborne Diseases Rhizoctonia Seedling Rot <i>(Rhizoctonia solani)</i>	0.40 - 0.80 fl. oz/ 1,000 row feet	See directions and rates under the Soilborne/Seedling Disease Control section of this label.
CROP LISTS: Pepper, bell; pepper, non-bell; pepper, sweet non-bell; eggplant, african; eggplant, pea; eggplant, scarlet; okra; pepino, tomatillo; cultivars, varieties and/or hybrids of these		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 62 fl. oz. (1.0 lbs. AI) of this product per acre per year. • DO NOT make more than four (4) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 0 days 		

**GRAPES AND OTHER SMALL FRUIT VINE CLIMBING (EXCEPT FUZZY KIWIFRUIT)
 SUBGROUP 13-07F**

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Black Rot (<i>Guignardia bidwellii</i>) Downy Mildew (<i>Plasmopara viticola</i>) Phomopsis Cane and Leaf Spot (<i>Phomopsis viticola</i>) Powdery Mildew (<i>Uncinula necator</i>) Suppression only: Botrytis Bunch Rot (<i>Botrytis cinerea</i>)	10.0 – 15.5 (0.16 - 0.25)	<p>Apply this product before disease outbreak. If environmental conditions are favorable to disease development, make additional applications at 7 to 14-day intervals throughout the year.</p> <p>Apply this product by ground, air or chemigation.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p> <p>This product is extremely phytotoxic to certain apple varieties. AVOID SPRAY DRIFT. Extreme care must be made to avoid injury to apple trees and apple fruit. DO NOT spray this product where spray drift may reach apple trees. DO NOT use spray equipment which has been previously used to apply this product to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.</p> <p>AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.</p>
<p>CROP LISTS: Amur River Grape; Kiwifruit; hardy maypop; muscadines; schisandra berry); cultivars, varieties and/or hybrids of these</p>		
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 14 days 		

HERBS & SPICES (EXCEPT BLACK PEPPER), CROP GROUP 19 (See list below)

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Corynespora Blight (<i>Corynespora cassiicola</i>) Dill Blight (<i>Cercosporidium punctum</i>) Phoma Blight (<i>Passalora puncta</i>)	6.0 - 15.5 (0.10 - 0.25)	Apply this product when disease outbreak begins and continue throughout the year at 7-day intervals. Apply this product by ground only in a minimum of 30 gallons of water per acre. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.
CROP LIST: Allspice; Angelica; Anise (seed); Anise, star; Annatto; Balm; Basil; Borage; Burnet; Chamomile; Caper (buds); Caraway; Caraway, black; Cardamom; Cassia (buds); Catnip; Celery Seed; Chervil (dried); Chive; Chive, Chinese; Cinnamon; Clary; Clove (buds); Coriander (cilantro or Chinese parsley) (leaf); Coriander (seed); Costmary; Culantro (leaf and seed); Cumin; Curry (leaf); Dill (seed); Dillweed; Fennel, Common; Fennel, Florence (seed); Fenugreek; Grains of Paradise; Horehound; Hyssop; Juniper (berry); Lavender; Lemongrass; Lovage (leaf and seed); Mace; Marigold; Marjoram; Mustard (seed); Nasturtium; Nutmeg; Parsley (dried); Pennyroyal; Pepper, White; Poppy Seed; Rosemary; Rue; Saffron; Sage; Savory, Summer and Winter Sweet Bay; Tansy; Tarragon; Thyme; Vanilla; Wintergreen; Woodruff; Wormwood		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 0 days 		

LEAFY VEGETABLES (EXCEPT BRASSICA), CROP GROUP 4

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
<p>Foliar Diseases Alternaria Leaf Spot (<i>Alternaria sonchi</i>, <i>A. spp.</i>) Anthracnose (<i>Microdochium panattonianum</i>, <i>Colletotrichum dematium</i>) Ascochyta Leaf Spot (<i>Ascochyta spp.</i>) Cercospora Leaf Spot (<i>Cercospora spp.</i>) Rust (<i>Puccinia spp.</i>), (<i>Uromyces spp.</i>) Septoria Leaf Spot (<i>Septoria petroselini</i>) White Rust (<i>Albugo occidentalis</i>)</p>	<p>6.0 - 15.5 (0.10 - 0.25)</p>	<p>Apply this product before disease outbreak. If environmental conditions are favorable to disease development, make additional applications at 10 to 14-day intervals throughout the year.</p> <p>Apply this product by ground, air or chemigation.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p> <p>ATTENTION: A tank mix of this product with other pesticides and adjuvants could result in phytotoxicity. DO NOT mix this product with any other pesticides and/or adjuvants that can increase foliar penetration including, but not limited to: products containing permethrin, foseetyl-AI, or lambda-cyhalothrin, , and silicone wetters.</p>
<p>Downy Mildew (<i>Bremia lactucae</i>) Powdery Mildew (<i>Eyrisiphe cichoracearum</i>)</p>	<p>12.0 - 15.5 (0.20 - 0.25)</p>	<p>Apply this product before disease outbreak or when conditions are favorable to disease development. Repeat at 5- to 7-day intervals for as long as conditions favor disease. Use the higher rate and a 5-day application interval when conditions are favorable to disease development.</p> <p>Apply this product by ground, air or chemigation.</p> <p>DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
<p>Soilborne Diseases Webb Blight, Bottom Rot, Crater Rot , Root Rot (<i>Rhizoctonia solani</i>)</p>	<p>0.40 - 0.80 fl. oz./ 1,000 row ft.</p>	<p>See directions and rates under the Soilborne/Seedling Disease Control section of this label.</p>
<p>CROP LIST: Amaranth; Arugula; Cardoon; Celery; Celtuce; Chervil; Chrysanthemum, Edible; Corn Salad; Cress; Dandelion; Dock; Endive; Fennel; Lettuce, Head and Leaf; Orach; Parsley; Purslane; Radicchio; Rhubarb; Spinach; Swiss Chard; cultivars, varieties and/or hybrids of these</p>		
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 5 days • Pre-harvest Interval (PHI): 0 days 		

**LEGUME VEGETABLES, DRY & SUCCULENT, CROP GROUP 6
 LEGUME VEGETABLES, FOLIAGE OF ANY CULTIVAR OF BEAN (*PHASEOLUS* SPP.)
 AND FIELD PEA (*PISIUM* SPP.), CROP GROUP 7**

(See crop list below)

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Bean Rust <i>(Uromyces appendiculatus)</i> Alternaria Blight (<i>Alternaria</i> spp.) Alternaria Leaf Spot (<i>Alternaria alternate</i>) Anthracnose (<i>Colletotrichum lindemuthianum</i>) Ascochyta Blight (<i>Mycosphaerella pinodes</i>) Ascochyta Leaf and Pod Spot (<i>Ascochyta</i> spp.) Ascochyta Leaf Spot (<i>Ascochyta phaseolorum</i>) Rust (<i>Phakopsora</i> spp.) Southern Blight (<i>Sclerotium rolfsii</i>) Web Blight (<i>Rhizoctonia solani</i>)	6.0 (0.10) 6.0 - 15.5 (0.10 - 0.25)	Apply this product before disease outbreak. If environmental conditions are favorable to disease development, make additional applications at 7- to 14-day intervals throughout the year. In case of severe disease pressure, use higher rate and shorter spray intervals. Apply this product by ground, air or chemigation. If an adjuvant is used, add it at the manufacturer's specified rates. If rust is present, use a non-ionic surfactant. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.
Soilborne Diseases Rhizoctonia Root Rot <i>(Rhizoctonia solani)</i>	0.40 - 0.80 fl. oz./ 1,000 row ft.	See directions and rates under the Soilborne / Seedling Disease Control section of this label. Apply this product to the furrow and covering soil at planting time in a 7 inch band. Avoid a concentrated stream directly on the seed or delayed emergence may occur. If using a narrow spray as an in-furrow spray, adjust the spray stream to hit the soil next to the seed but not hit the seed. NOTE: Conduct a seed safety test with your crop before making in-furrow applications.
CROP LIST: Bean (<i>Lupinus</i> spp.) (Includes grain lupin, sweet lupin, white lupin, and white sweet lupin); Bean (<i>Phaseolus</i> spp.) (Includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); Bean (<i>Vigna</i> spp.) (Includes adzuki bean, asparagus bean, blackeyed pea, cowpea, catjang, Chinese longbean, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); Bean (Glycine max) Soybean, Immature Seed (edamame); Broad bean (fava bean) (<i>Vicia faba</i>); Chickpea (garbanzo bean) (<i>Cicer arietinum</i>); Guar (<i>Cyamopsis tetragonoloba</i>); Jackbean (<i>Canavalia ensiformis</i>); Lablab Bean (hyacinth bean) (<i>Lablab purpureus</i>); Lentil (<i>Lens esculenta</i>); Pea (<i>Pisum</i> spp.) (includes dwarf pea, edible-pod pea, English pea, garden pea, green pea, field pea, snow pea, sugar snap pea); Pigeon Pea (<i>Cajanus cajan</i>); Sword Bean (<i>Canavalia gladiata</i>)		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-Harvest Interval (PHI) for dry legume vegetables (dry bean and dry pea seeds): 14 days • Pre-Harvest Interval (PHI) for succulent beans and peas: 0 days <p>For use on soybeans, please refer to the soybean crop use directions.</p>		

PEANUTS

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
<p>Soilborne Diseases. Early season in-furrow application Aspergillus Crown Rot (<i>Aspergillus niger</i>) Pythium Damping Off (<i>Pythium</i> spp.)</p> <p>Suppression only: Stem Rot/White Mold (<i>Sclerotium rolfsii</i>)</p>	<p>0.40-0.80 fl. oz./ 1,000 row ft.</p>	<p>For control of several seed/seedling diseases including early season suppression of stem rot, apply this product in-furrow at planting.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
<p>Soilborne Diseases – mid-late season Rhizoctonia Peg and Pod Rot (<i>Rhizoctonia solani</i>) Stem Rot/White Mold (<i>Sclerotium rolfsii</i>)</p> <p>Suppression Only: Cylindrocladium Black Rot (<i>Cylindrocladium crotalariae</i>) Pythium Pod Rot (<i>Pythium myriotylum</i>)</p>	<p>18.5 – 24.5 (0.30 - 0.40)</p>	<p>Use this product in a typical preventative fungicide program for control of soilborne diseases. Apply approximately 60 and 90 days after planting. Adjust application timing if local conditions favor early disease outbreak.</p> <p>For control of Pythium, a rate of 24.5 fl. oz./A is required.</p> <p>Apply this product by ground, air or chemigation.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p> <p>Use this product in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, including proper crop rotation practices. Consult with your Extension Service representatives for guidance on the proper use of this product in programs which attempt to minimize the occurrence of disease resistance to fungicides.</p>
<p>Foliar Diseases Early Leaf Spot (<i>Cercospora arachidicola</i>) Late Leaf Spot (<i>Cercosporidium personatum</i>) Rust (<i>Puccinia arachidis</i>) Web Blotch (<i>Phoma arachidicola</i>)</p>	<p>6.0 - 18.5 (0.10 - 0.30)</p>	<p>Apply this product when conditions favor disease, when leaf wetness first occurs or 30 to 40 days after planting. Repeat applications at 10- to 14-day intervals if conditions remain favorable for disease.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 24.5 fl. oz. (0.40 lbs. AI) of this product per acre per application. • DO NOT apply more than 49 fl. oz. (0.80 lbs. AI) of this product per acre per year. • DO NOT make more than two (2) applications of this product at the high rate (0.40 lbs. AI / acre) per year. • Minimum Retreatment Interval: 10 days • Pre-harvest Interval (PHI): 14 days 		

PECANS

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Anthracnose <i>(Glomerella cingulata)</i> Scab <i>(Cladosporium caryigenum)</i>	6.0 - 12.0 (0.10 - 0.20)	Apply this product before disease outbreak or when conditions are favorable to disease development. If environmental conditions are favorable to continued disease development, make a second application after 7 to 14 days, dependent upon the severity of disease pressure. Apply the higher rate and shorter application intervals when disease pressure is severe. Apply this product by ground, air or chemigation. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than Qol Group 11.

RESTRICTIONS:

- **DO NOT** apply more than 12.0 fl. oz. (0.20 lbs. AI) of this product per acre per application.
- **DO NOT** apply more than 72 fl. oz. (1.2 lbs. AI) of this product per acre per year.
- **DO NOT** make more than six (6) applications of this product at the high rate (0.20 lbs. AI / acre) per year.
- **Minimum Retreatment Interval:** 7 days
- **Pre-harvest Interval (PHI):** 45 days

PISTACHIOS

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Alternaria Late Blight <i>(Alternaria alternata)</i> Botryosphaeria Panicle and Shoot Blight <i>(Botryosphaeria dothidea)</i> Septoria Leaf Spot <i>(Septoria pistaciarum)</i>	6.0 - 15.5 (0.10 - 0.25)	Apply this product before disease outbreak or when conditions are favorable to disease development. If environmental conditions are favorable to continued disease development, make a second application after 7 to 21 days, dependent upon the severity of disease pressure. Apply the higher rate and shorter application intervals when disease pressure is severe. Apply this product by ground, air or chemigation. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than Qol Group 11.

RESTRICTIONS:

- **DO NOT** apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application.
- **DO NOT** apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year.
- **DO NOT** make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year.
- **Minimum Retreatment Interval:** 7 days
- **Pre-harvest Interval (PHI):** 7 days

POTATOES

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Black Dot <i>(Colletotrichum coccodes)</i> Early Blight <i>(Alternaria solani)</i> Late Blight <i>(Phytophthora infestans)</i> Powdery Mildew <i>(Erysiphe cichoracearum)</i>	6.5 – 20.0 (0.11 – 0.33)	<p>Apply this product before disease outbreak or when conditions are favorable to diseases development.</p> <p>Early Blight: Apply every 7 to 14 days following resistance management practices. Use the higher rate and a 7-day application interval when conditions are favorable to disease development.</p> <p>Late Blight: Apply a minimum of 12.0 fl. oz./A on a 7-day schedule. If late blight symptoms develop or conditions favor disease, switch immediately to a non-Group 11 fungicide, using a 5-day schedule. Addition of spreader/sticker may improve coverage.</p> <p>Other diseases: Apply this product prior to disease development and continue throughout the season every 7 to 14 days. Use the higher rate and the shorter interval if disease epidemics are severe.</p> <p>Apply this product by ground, air or chemigation.</p> <p>Tank mixtures of this product with other pesticides and adjuvants must be tested on a small scale for crop safety prior to application to the entire crop.</p> <p>DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
Soilborne Diseases Black Dot <i>(Colletotrichum coccodes)</i> Black Scurf <i>(Rhizoctonia solani)</i> Silver Scurf <i>(Helminthosporium solani)</i>	0.40 - 0.80 fl. oz./ 1,000 row ft.	<p>See directions and rates under the Soilborne / Seedling Disease Control section of this label.</p>
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 20 fl. oz. (0.33 lbs. AI) of this product per acre per application. • DO NOT apply more than 120 fl. oz. (2.0 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.33 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 14 days 		

RICE

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Sheath / Stem Diseases Sheath Blight <i>(Rhizoctonia solani)</i>	9.0 - 18.5 (0.15 - 0.30)	Apply this product before disease outbreak or when conditions are favorable to disease development. Sheath Blight: Application rates may vary from 9.0 to 12.0 fl. oz./A depending on the growth stage of the rice and the severity of the disease. Consult with your local extension representative for the technical bulletin on sheath blight control.
Aggregate Sheath Spot <i>(Ceratobasidium oryzae-sativae = Rhizoctonia oryzae-sativae)</i> Black Sheath Rot <i>(Gaeumannomyces graminis var. graminis)</i> Sheath Spot <i>(Rhizoctonia oryzae)</i> Stem Rot <i>(Magnaporthe salvinii = Nakateae sigmoidea)</i>	12.5 - 15.5 (0.20 - 0.25)	Apply this product by ground, by air at 5-10 gallons of water per acre or by chemigation. If an adjuvant is used, add it at the manufacturer's specified rates. For other stem/sheath diseases including stem rot, black sheath rot, aggregate sheath spot and sheath spot, make one application when disease is less than 4 inches above the water level typically between panicle differentiation (PD) +5 days to PD +10 days or at early detection of the disease. A second application may be necessary under heavy disease pressure and conditions favorable for disease. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than Qol Group 11.
Foliar Diseases Brown Leaf Spot <i>(Cochliobolus miyabeanus)</i> Leaf Smut <i>(Entyloma oryzae)</i> Narrow Brown Leaf Spot <i>(Cercospora janseana = Cercospora oryzae)</i>	12.5 - 15.5 (0.20-0.25)	Apply this product before disease outbreak or prior to favorable conditions for blast development. For panicle blast, apply at mid-boot to boot-split but prior to full head emergence. Apply second application when panicles are approximately 60-90% emerged from the boot (7-14 days later). Apply this product by ground, air or chemigation. In case there is no rotation to other crops, DO NOT make more than two sequential applications of this product in 2 crop seasons for Panicle Blast before alternating with a fungicide with a different mode of action than Qol Group 11.
Panicle Diseases Kernel Smut <i>(Tilletia barclayana = Neovossia barclayana)</i> Panicle Blast <i>(Pyricularia grisea)</i>		DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than Qol Group 11.
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 18.5 fl. oz. (0.30 lbs. AI) of this product per acre per application. • DO NOT apply more than 42 fl. oz. (0.70 lbs. AI) of this product per acre per year. • DO NOT make more than two (2) applications of this product at the high rate (0.30 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 28 days • DO NOT treat rice fields used for aquaculture of fish and crustaceans. • DO NOT apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Applicators must use care in making applications near non-target aquatic habitats. • DO NOT allow release of irrigation or flood water for at least 14 days after the last application. 		

SOYBEAN

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Aerial Blight (<i>Rhizoctonia solani</i>) Alternaria Leaf Spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum truncatum</i>) Brown Spot (<i>Septoria glycines</i>) Cercospora Blight and Leaf Spot (<i>Cercospora kikuchii</i>) Frogeye Leaf Spot (<i>Cercospora sojina</i>) Pod and Stem Blight (<i>Diaporthe phaseolorum</i>) Rust (<i>Phakopsora</i> spp.)	6.0 - 15.5 (0.10 - 0.25)	Apply this product before disease outbreak or when conditions are favorable to disease development. If environmental conditions are favorable to continued disease development, make a second application after 14 to 21 days, dependent upon the severity of disease pressure. Apply the higher rate and shorter application intervals when disease pressure is severe. Apply this product by ground, air or chemigation. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.
Soilborne Diseases Rhizoctonia Solani (<i>Rhizoctonia solani</i>) Southern Blight (<i>Sclerotium rolfsii</i>)	0.40-0.80 fl. oz./ 1,000 row ft.	See directions and rates under the Soilborne / Seedling Disease Control section of this label.
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT make more than one application at 15.5 fl. oz (0.25 lbs. AI) product per acre per year to soybean forage and hay. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 14 days • Pre-harvest Interval (PHI) for harvest of soybeans: 14 days • Pre-harvest Interval (PHI) for harvest of soybean forage and hay: 0 days 		

- **DO NOT** apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application.
- For soybean forage and hay, **DO NOT** apply more than 15.5 fl. oz (0.25 lbs. AI) product per acre per year.
- For soybean forage and hay, **DO NOT** make more than one application.
- For soybean seed, **DO NOT** apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year.
- For soybean seed, **DO NOT** make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year.
- **Minimum Retreatment Interval:** 14 days
- **Pre-harvest Interval (PHI) for harvest of soybean forage and hay:** 0 days
- **Pre-harvest Interval (PHI) for harvest of soybeans:** 14 days

STONE FRUIT, CROP GROUP 12-12

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Brown Rot Blossom Blight and Fruit Rot (<i>Monilinia fructicola</i> , <i>M. laxa</i>)	12.0 - 15.5 (0.20 - 0.25)	DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.
Scab (<i>Cladosporium carpophilum</i>) Alternaria Spot and Fruit Rot (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum prunicola</i> , <i>C. gloeosporioides</i>) Leaf Rust (<i>Tranzschelia discolor</i>) Powdery Mildew (<i>Sphaerotheca pannosa</i> , <i>Podosphaera clandestina</i>) Shot Hole (<i>Wilsonomyces carpophilus</i>)	12.0 - 15.5 (0.20 - 0.25)	<p>Brown Rot Blossom Blight: Begin applications at early bloom and continue through petal fall.</p> <p>Brown Rot Fruit Rot: Apply this product to fruit up to the day of harvest.</p> <p>Scab: Begin applications at petal fall and continue at 7 to 14 day intervals. Peaches: apply 9.0 – 15.5 fl.oz. per acre.</p> <p>All other diseases: Apply this product at the onset of disease and continue on a 7 to 14 day schedule. Apply this product by ground, air or chemigation.</p>
<p>CROP LIST: Apricot; Cherry, Sweet; Cherry, Tart; Nectarine; Peach; Plum; Plumcot; Prune; cultivars, varieties and/or hybrids of these</p>		
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 93 fl. oz. (1.5 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 0 days 		

TOMATOES, SUBGROUP 8-10A

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Anthracnose (<i>Colletotrichum coccodes</i>) Black Mold (<i>Alternaria alternata</i>) Buckeye Rot (<i>Phytophthora</i> spp.) Early Blight (<i>Alternaria solani</i>) Powdery Mildew (<i>Oidiopsis sicula</i>) Septoria Leaf Spot (<i>Septoria lycopersici</i>) Target Spot (<i>Corynespora cassiicola</i>)	5.0 - 6.2 (0.08 - 0.10)	Apply this product before disease outbreak or when conditions are favorable to disease development. Repeat applications if conditions remain favorable for disease. When disease pressure is severe, apply the higher rate. Late Blight: Apply this product at 5 to 7 day intervals. All other diseases: Apply this product at 7 to 21 day intervals. Under certain weather conditions (particularly high temperatures) this product, in combination with high rates of silicone-based or oil containing (petroleum or crop) additives or adjuvants, may cause injury. DO NOT exceed 0.125% adjuvant (v/v).
Late Blight (<i>Phytophthora infestans</i>)	6.2 (0.10)	A tank mixture with Dimethoate may cause crop injury. On fresh market tomatoes, DO NOT use adjuvants or tank mix this product with any emulsifiable concentrate (EC) product. Apply this product by ground, air or chemigation. DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than QoI Group 11.
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 6.2 fl. oz. (0.1 lbs. AI) of this product per acre per application. • DO NOT apply more than 37.2 fl. oz. (0.6 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.10 lbs. AI / acre) per year. • Minimum Retreatment Interval: 5 days • Pre-harvest Interval (PHI): 0 days 		

TREE NUTS, CROP GROUP 12-14 See specific instructions for Almonds and Pistachios.

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Alternaria Leaf and Fruit Spot (<i>Alternaria alternata</i>) Blossom Blight (<i>Monilinia laxa</i> , <i>M. fructicola</i>) Late Blight (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum acutatum</i> , <i>Glomerella cingulata</i>) Eastern Filbert Blight (<i>Anisogramma anomale</i>) Scab (<i>Cladosporium carpophilum</i>) Septoria Leaf Spot (<i>Septoria pistaciarum</i>) Shot Hole (<i>Wilsonomyces carpophilus</i>)	12.0 (0.20)	Apply this product before disease outbreak or when conditions are favorable to disease development. If environmental conditions are favorable to continued disease development, make a second application after 7 to 21 days, dependent upon the severity of disease pressure. Blossom blight: Apply this product at early bloom and continue through petal fall. Apply this product by ground, air or chemigation. If an adjuvant is used, add it at the manufacturer's specified rates. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.
CROP LIST: Beechnut; Brazil Nut; Butternut; Cashew; Chestnut; Chinquapin; Filbert (hazelnut); Hickory Nut; Macadamia Nut; Pecan; Walnut, English and black		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 12.0 fl. oz. (0.20 lbs. AI) of this product per acre per application. • DO NOT apply more than 72 fl. oz. (1.2 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.20 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 45 days 		

**ROOT VEGETABLES, (EXCEPT SUGAR BEET) SUBGROUP 1B
 LEAVES OF ROOT AND TUBER VEGETABLES, GROUP 2**

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Foliar Diseases Alternaria Leaf Spot (<i>Alternaria</i> spp., <i>A. alternata</i>) Ascochyta Leaf Spot (<i>Ascochyta cynarae</i>) Rust (<i>Uromyces betae</i> , <i>Puccinia helianthi</i>) White Rust (<i>Albugo tragopogonis</i>)	6.0 - 20.0 (0.10 - 0.33)	Apply this product before disease outbreak or when conditions are favorable to disease development. Powdery Mildew: Apply every 5 to 7 days. All other diseases: Apply every 7 to 14 days. Apply the higher rate and shorter application intervals when disease pressure is severe. Apply this product by ground, air or chemigation. Apply as in in-furrow spray in a minimum of 10 gallons per acre.
Cercospora Leaf Spot (<i>Cercospora betae</i> , <i>C. pastinaceae</i>) Powdery Mildew (<i>Erysiphe polygoni</i> , <i>Leveillula taurica</i>)	9.0 - 15.5 (0.15 - 0.25)	Apply as in in-furrow spray in a minimum of 10 gallons per acre. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.
Soilborne Diseases Circular Spot, Southern Blight (<i>Sclerotium rolfsii</i>) Pythium Root Rot (<i>Pythium aphanidermatum</i>) Rhizoctonia Stem Canker, Crown Root Rot (<i>Rhizoctonia solani</i>)	0.40 - 0.80 fl. oz./ 1,000 row ft.	See directions and rates under the Soilborne / Seedling Disease Control section of this label. For sugar beets apply 3-7 inch banded applications in a minimum of 10 gallons per acre at the 2 to 8 leaf stage. DO NOT apply as a dribble application over the seed row. Tank mixtures of this product with crop oil concentrates (COC) or methylated spray oil (MSO) may result in crop injury. If cool soil conditions are expected after planting, which could result in an extended period of plant emergence, this product must not be applied in-furrow. If using this product at the time of planting, DO NOT use a starter fertilizer with it.
CROP LIST: Beet, Garden ^{1,2} ; Beet, Sugar ² ; Burdock, edible ^{1,2} ; Carrot ^{1,2} ; Cassava, Bitter and Sweet ² ; Celery (celery root) ^{1,2} ; Chervil, Turnip-Rooted ^{1,2} ; Chicory ^{1,2} ; Dasheen (taro) ² ; Ginseng ¹ ; Horseradish ¹ ; Parsley, turnip-Rooted ¹ ; Parsnip ^{1,2} ; Radish ^{1,2} ; Radish, Oriental (daikon) ^{1,2} ; Rutabaga ^{1,2} ; Salsify ¹ ; Salsify, Black ^{1,2} ; Salsify, Spanish ¹ ; Skirret ¹ ; Sweet Potato ² ; Tanier ² ; Turnip ^{1,2} ; Yam, True ² ¹ = Root vegetables, (except sugar beet) Subgroup 1B ² = Leaves of root and tuber vegetables Group 2		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 20.0 fl. oz. (0.33 lbs. AI) of this product per acre per application. • DO NOT apply more than 123 fl. oz. (2.0 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.33 lbs. AI / acre) per year. • Minimum Retreatment Interval: 5 days • Pre-harvest Interval (PHI): 0 days 		

VEGETABLES, TUBEROUS AND CORM SUBGROUP 1C

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Foliar Diseases Alternaria Leaf Spot (<i>Alternaria</i> spp., <i>A. alternata</i>) Ascochyta Leaf Spot (<i>Ascochyta cynarae</i>) Rust (<i>Uromyces betae</i> , <i>Puccinia helianthi</i>) White Rust (<i>Albugo tragopogonis</i>)	6.0 - 20.0 (0.10 - 0.33)	Apply this product before disease outbreak or when conditions are favorable to disease development. Powdery Mildew: Apply every 5 to 7 days following resistance management practices. All other diseases: Apply every 7 to 14 days following resistance management practices.
Cercospora Leaf Spot (<i>Cercospora betae</i> , <i>C. pastinaceae</i>) Powdery Mildew (<i>Erysiphe polygoni</i> , <i>Leveillula taurica</i>)	9.0 - 15.5 (0.15 - 0.25)	Apply the higher rate and shorter application intervals when disease pressure is severe. Apply this product by ground, air or chemigation. DO NOT make more than one application of this product before alternating with fungicides with a mode of action other than QoI Group 11.
Soilborne Diseases Circular Spot, Southern Blight (<i>Sclerotium rolfsii</i>) Rhizoctonia Stem Canker, Crown Rot (<i>Rhizoctonia solani</i>) Pythium Root Rot (<i>Pythium aphanidermatum</i>)	0.40 - 0.80 fl. oz./ 1,000 row ft.	See directions and rates under the Soilborne/Seedling Disease Control section of this label.
CROP LIST: Arracacha; Arrowroot; Artichoke, Chinese and Jerusalem; Canna, Edible; Cassava, Edible; Cassava, Bitter and Sweet; Chayote (root); Chufa; Dasheen (Taro); Ginger; Leren; Potato; Sweet Potato; Tanier; Turmeric; Yam, Bean; Yam, True		
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 20.0 fl. oz. (0.33 lbs. AI) of this product per acre per application. • DO NOT apply more than 123 fl. oz. (2.0 lbs. AI) of this product per acre per year. • DO NOT make more than six (6) applications of this product at the high rate (0.33 lbs. AI / acre) per year. • Minimum Retreatment Interval: 5 days • Pre-harvest Interval (PHI): 14 days 		

WHEAT AND TRITICALE

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Leaf Rust (<i>Puccinia triticina</i> = <i>Puccinia recondita</i> f.sp <i>tritici</i>) Septoria Leaf and Glume Blotch (<i>Septoria tritici</i> , <i>Septoria nodorum</i>) Stem Rust (<i>Puccinia graminis</i>) Stripe Rust (<i>Puccinia striiformis</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>)	4.0 - 12.0 (0.07 - 0.20)	Apply this product before disease outbreak or when conditions are favorable to disease development. Apply this product by ground, air or chemigation. A crop oil concentrate adjuvant may be added at 1.0% v/v to optimize efficacy.
Powdery Mildew (<i>Erysiphe graminis</i>)	7.5 - 11.0 (0.125-0.175)	DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.

RESTRICTIONS:

- **DO NOT** apply after Feekes 10.54.
- **DO NOT** apply more than 12.0 fl. oz. (0.20 lbs. AI) of this product per acre per application.
- **DO NOT** apply more than 24 fl. oz. (0.40 lb. AI) of this product per acre per year.
- **DO NOT** make more than two (2) applications of this product at the high rate (0.20 lbs. AI / acre) per year.
- **Minimum Retreatment Interval:** 14 days
- **Pre-harvest Interval (PHI) for forage and hay:** 7 days
- **Pre-harvest Interval (PHI) for grazing:** 14 days

WILD RICE

Target Diseases	Rate fl. oz./Acre (lbs.AI/A)	Application Directions
Brown Spot (<i>Bipolaris oryzae</i> or <i>Bipolaris sorokiana</i>) Also known as <i>elminthosporium oryzae</i> and <i>H. sativum</i> Stem rot (<i>Nakataea sigmoidea</i>)	12.5 – 15.5 (0.20 – 0.25)	Apply this product before disease outbreak or when conditions are favorable to disease development. Apply this product by ground, by air in 5-10 gallons of water per acre, or by chemigation. If an adjuvant is used, add it at the manufacturer’s specified rates. For foliar diseases, apply this product preventively before disease outbreak. Apply during tillering, boot, early heading, or at early detection of disease. Make a second application under heavy disease pressure and when conditions are favorable for disease development. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.
<p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 42 fl. oz. (0.70 lbs. AI) of this product per acre per year. • DO NOT make more than two (2) applications of this product at the high rate (0.25 lbs. AI / acre) per year. • Minimum Retreatment Interval: 7 days • Pre-harvest Interval (PHI): 28 days • DO NOT allow release of irrigation or flood water for at least 14 days after the last application. • DO NOT treat wild rice fields used for aquaculture of fish and crustaceans. • DO NOT apply when weather conditions favor drift from treat areas to non-target aquatic habitat. Applicators must use care in making applications near non-target aquatic habitats. 		

TURF

This product is for control of labeled diseases on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Integrated Pest (Disease) Management

Integrate this product into an overall disease and pest management strategy whenever the use of a fungicide is required. Follow cultural practices known to reduce disease development, including varieties with disease resistance, removal of plant debris in which inoculums overwinter, and proper timing and placement of irrigation. Consult your agricultural authorities for additional IPM strategies established for your area. This product may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development.

Resistance Management

Some turf diseases are known to have developed resistance to products used repeatedly for their control. Apply this product in a tank mix or alternation program with other registered fungicides with different modes of action and to which pathogen resistance has not developed.

DO NOT make more than two sequential applications of this product for control of *Pythium* spp. and Gray Leaf Spot. For all other diseases when *Pythium* spp. is not present, **DO NOT** make more than three sequential applications of this product.

Application Directions

Apply this product prior to disease development. Mix with the required amount of water and apply as a dilute spray application in 2-4 gallons of water per 1000 sq. ft. (87–174 gallons per acre). Repeat at specified application intervals

until disease is under control or until the maximum treatment rate has been reached. See **Rate Conversion Chart for Turf** for maximum number of applications at each of the specified rates.

For spot treatments, use 0.4 fl. oz. of this product per 1 to 2 gallons of water.

Apply the higher rate and shorter application intervals when prolonged disease conditions exist.

Note: This product does not control Dollar Spot. It is compatible in tank mixes with many other fungicides that do control Dollar Spot. Always tank mix this product with another fungicide that controls Dollar Spot when this disease is present.

Follow directions under Tank Mixes below.

RESTRICTIONS

- Apply by ground (includes any hand-held and mechanically pressurized handwands) only.
- **DO NOT** use more than the maximum allowed use rate down in the “Application Directions for Turf” chart below.
- **DO NOT** apply more than 8.8 quarts of product per acre per year (6.46 fl. oz. product/1000 sq. ft./year) (0.52 lbs. AI/acre/year). See **Rate Conversion Chart for Turf** for maximum number of applications at each of the specified rates.
- **DO NOT** make more than two sequential applications of this product for control of *Pythium* spp. For all other diseases, **DO NOT** make more than three sequential applications of this product.
- Follow the Application Interval requirements in the “Application Directions for Turf” chart below.

Tank Mixes

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 qt. water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

When mixed with EC (emulsifiable concentrate) formulations, this product may be phytotoxic, especially when applied during cool, cloudy conditions that last for several days. Adjuvants containing silicone could also have phytotoxic effects.

APPLICATION DIRECTIONS FOR TURF DISEASES

Target Diseases	Use Rate (fl. oz./ 1000 sq. ft.)	Application Interval (days)	Application Directions
Anthracnose (<i>Colletotrichum graminicola</i>) Brown Patch (<i>Rhizoctonia solani</i>) Fusarium Patch (<i>Microdochium nivale</i>) Gray Leaf Spot (<i>Pyricularia grisea</i>) Red Thread (<i>Laetisaria fuciformis</i>) Southern Blight (<i>Sclerotium rolfsii</i>) Summer Patch	0.38-0.77	14 – 28	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days

Target Diseases	Use Rate (fl. oz./ 1000 sq. ft.)	Application Interval (days)	Application Directions
<i>(Magnaporthe poae)</i>			
Cool Weather Brown Patch Yellow Patch <i>(Rhizoctonia cerealis)</i>	0.77	28	Make one or two applications in fall or when conditions are favorable for disease development. Minimum Retreatment Interval: 28 days
Fairy Ring <i>(Lycoperdon spp., Agrocybe pediades, and Bovistra plumbea)</i>	0.77	28	Apply this product as soon as possible after fairy ring symptoms develop. Irrigate with at least 0.5 inches water within 24 hours after application if rainfall is not expected. Severely damaged or thin turf may require reseeding. Fairy ring symptoms may take 2 to 3 weeks to disappear following application. Make an additional application 14-28 days if needed. Minimum Retreatment Interval: 28 days
Gray Snow Mold* Typhula Blight <i>(Typhula incarnata, T. ishikariensis)</i>	1.33	Single Application	Make a single application of 1.33 fl. oz. or two applications of 0.67 fl.oz. at 14-day intervals in late fall just before snow cover. When disease pressure is severe, tank mix with another snow mold fungicide to enhance control. Minimum Retreatment Interval: 28 days *NOT for use in California.
	0.67	28	
Leaf Spot <i>(Bipolaris sorokiniana)</i> Melting Out <i>(Drechslera poae)</i>	0.38 – 0.77	14 – 21	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Leaf Rust <i>(Puccinia spp.)</i>	0.38 – 0.77	14 – 28	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Necrotic Ring Spot <i>(Leptosphaeria korrae)</i>	0.77	14 – 28	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Pink Patch <i>(Limonomyses roseipellis)</i>	0.38 - 0.77	14 – 28	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Pink Snow Mold* <i>(Microdochium nivale)</i>	1.33	Single Application	Make a single application of 1.33 fl. oz. or two applications of 0.67 at 14-day intervals in late fall just before snow cover. When disease pressure is severe, tank mix with another snow mold fungicide to enhance control. Minimum Retreatment Interval: 14 days *NOT for use in California.
	0.67	14	
Powdery Mildew <i>(Blumeria graminis)</i>	0.38 – 0.77	14 – 28	Apply this product before disease outbreak or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Pythium Blight Pythium Root Rot <i>(Pythium aphanidermatum, Pythium spp.)</i>	0.77	10 – 14	Apply this product before disease is present. During periods of prolonged favorable conditions, treat on the 10 day application interval. This product can be used on newly seeded as well as established turf.

Target Diseases	Use Rate (fl. oz./ 1000 sq. ft.)	Application Interval (days)	Application Directions
			Minimum Retreatment Interval: 10 days
Pythium Root Dysfunction (<i>Pythium volutum</i>)	0.77	21 – 28	Apply preventatively when mean daily soil temperatures are between 55°F and 70°F. Irrigate with 0.1 to 0.2 inches within 24 hours after application to facilitate movement into the root zone. Minimum Retreatment Interval: 21 days
Rhizoctonia Large Patch** (<i>Rhizoctonia solani</i>)	0.38 - 0.77	14 – 28	Make one or two applications in fall or when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days **Not for use at the lower rate in California.
Rhizoctonia Leaf Spot (<i>Rhizoctonia zeae</i>)	0.77	14 – 28	Apply when conditions are favorable for disease development. Minimum Retreatment Interval: 14 days
Spring Dead Spot (<i>Leptosphaeria korrae</i>) or (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>) or (<i>Ophiosphaerella herpotricha</i>)	0.77	28	Make one or two applications in fall or when conditions are favorable for disease development. Minimum Retreatment Interval: 28 days
Take-all Patch (<i>Gaeumannomyces graminis</i> var. <i>avenae</i>)	0.77	28	Make two applications at 28-day intervals in the spring and two applications at 28-day intervals in the fall. Minimum Retreatment Interval: 28 days
Zoysia Patch (<i>Rhizoctonia solani</i> and/or <i>Gaeumannomyces incrustana</i>)	0.38 - 0.77	14 – 28	Make one or two applications in late fall before snow cover or when conditions are favorable for disease development. DO NOT apply on top of snow. Minimum Retreatment Interval: 14 days

Rate Conversion Chart for Turf

Fluid Ounces Product/1000 sq. ft.	Ounces AI /1000 sq. ft.	Fluid Ounces Product/Acre	Pints of Product / Acre	Maximum number of applications at specified rate
0.38	0.100	16.6	1.03	17
0.5	0.130	21.8	1.4	13
0.6	0.156	26.1	1.6	12
0.67	0.174	29.2	1.8	10
0.77	0.200	33.5	2.1	8
1.33	0.34	57.9	3.6	5

Amount to Mix per 100 Gallons for Turf Applications

Use Rate (fl. oz.)	Spray Volume (gallons/1000 square feet)		
	2.0 gals.(fl. oz.)	3.0 gals. (fl. oz.)	4.0 gals. (fl. oz.)
0.38	19	12.7	9.5
0.5	25	17	13
0.6	30	20	15
0.67	33.5	22.4	16.8
0.77	38.5	25.7	19.3
1.33	66.5	44.3	33.2

ORNAMENTALS [(Restriction: Not for use in California)]

This product is for control of certain pathogens causing foliar, aerial, and root diseases, including leaf, tip and flower blights, leaf spots, downy mildew, powdery mildew, anthracnose, and rusts of ornamental plants. Use this product to control certain diseases of container, bench, flat, plug, bed or field-grown ornamentals in greenhouses, shade-houses, outdoor nurseries, retail nurseries and other landscape areas.

Integrated Pest (Disease) Management

Integrate this product into an overall disease and pest management strategy whenever the use of a fungicide is required. Follow cultural practices known to reduce disease development, including selection varieties with disease resistance, optimum plant populations, proper fertilization, winter and/or spring pruning, plant residue management and proper timing and placement of irrigation. Consult your agricultural authorities for additional IPM strategies established for your area. This product may be used in State Agricultural Extension advisory (disease forecasting) programs which specify application timing based on environmental factors favorable for disease development.

Application Directions

Apply this product as a broadcast or banded spray, targeting the foliage or crown of the plant. Apply to runoff in sufficient water, ensuring complete coverage of the target plant. Best control of targeted diseases is attained with sufficient coverage and wetting of foliage. Refer to the specific use directions for control of certain diseases. Repeat at specified application intervals (plus alternations for resistance management) until disease is under control or until the maximum treatment rate has been reached. (see **Application Rates** chart below).

Begin applications of this product prior to disease development and continue throughout the season at the specified intervals following resistance management guidelines. Use this product as part of a preventative disease management program.

This product may not provide adequate disease control when applied after disease outbreak.

Application Rates

See charts below for the specific rate range for this product. This product can be applied every 7 – 28 days (or as otherwise specified on this label for a specific plant or disease). Adding a non-silicone based wetting/sticking agent at the specified use rate may enhance coverage on hard-to-wet plant foliage.

Field-grown or Nursery ornamentals (ground and chemigation applications)

Conditions	Rate Range	Retreatment Intervals	Maximum Number of Applications
For most conditions and for most diseases	3.85 – 5.75 fl. oz. / 100 gallons	7 – 14 days	DO NOT apply more than 0.36 gallons (46 fl. oz. or 0.75 lbs.AI) of this product per acre per year or 8 applications per year.
	1.9 – 2.88 fl. oz. / 50 gallons	7 – 14 days	
When disease pressure is light to moderate	1.9 – 2.88 fl. oz. / 100 gallons	7 – 14 days	
	0.95 – 1.44 fl. oz. / 50 gallons	7 – 14 days	
	5.75 fl.oz. / 100 gallons	14 – 28 days	
	2.85 – 2.88 fl. oz. / 50 gallons	14 – 28 days	
When environmental conditions are favorable to severe disease development	5.75 fl.oz. / 100 gallons	7 – 14 days	
	2.85 – 2.88 fl. oz./50 gallons	7 – 14 days	

Nursery ornamentals broadcast application and Landscape (plants, flowers, trees, turf), (any hand-held applications)

Greenhouse ornamentals (mechanically pressurized handwand applications)

Conditions	Rate Range	Retreatment Intervals	Maximum Number of Applications
For most conditions and for most diseases	3.85 – 5.75 fl. oz. / 100 gallons	7 – 14 days	DO NOT apply more than 0.0012 gallons (0.15 fl. oz. or 0.0025 lbs.AI) of this product per acre per year or 8 applications per year.
	1.9 – 2.88 fl. oz. / 50 gallons	7 – 14 days	
When disease pressure is light to moderate	1.9 – 2.88 fl. oz. / 100 gallons	7 – 14 days	
	0.95 – 1.44 fl. oz. / 50 gallons	7 – 14 days	
	5.75 fl.oz. / 100 gallons	14 – 28 days	
	2.85 – 2.88 fl. oz. / 50 gallons	14 – 28 days	
When environmental conditions are favorable to severe disease development	5.75 fl.oz. / 100 gallons	7 – 14 days	
	2.85 – 2.88 fl. oz./50 gallons	7 – 14 days	

Ornamental Uses (except for Field-grown or Nursery ornamentals (ground and chemigation applications), any hand-held applications and mechanically pressurized handwand applications)

Conditions	Rate Range	Retreatment Intervals	Maximum Number of Applications
For most conditions and for most diseases	3.85 – 7.7 fl. oz. / 100 gallons	7 – 14 days	DO NOT apply more than 2.4 gallons (5.0 lbs. AI) of this product per acre per year or 8 applications per year.
	1.9 – 3.85 fl. oz. / 50 gallons	7 – 14 days	
When disease pressure is light to moderate	1.9 – 3.85 fl. oz. / 100 gallons	7 – 14 days	
	0.95 – 1.9 fl. oz. / 50 gallons	7 – 14 days	
	5.75 – 7.7 fl.oz. / 100 gallons	14 – 28 days	
	2.85 – 3.85 fl. oz. / 50 gallons	14 – 28 days	
When environmental conditions are favorable to severe disease development	5.75 – 7.7 fl.oz. / 100 gallons	7 – 14 days	
	2.85 – 3.85 fl. oz. / 50 gallons	7 – 14 days	

Application Rate Conversion Chart

Fl. Oz. Rate	Lbs. AI		Fl. Oz. Rate	Lbs. AI
0.38 fl. oz.	0.006		3.85 fl. oz.	0.063
0.95 fl. oz.	0.015		5.75 fl. oz.	0.093
1.44 fl. oz.	0.023		7.7 fl. oz.	0.125
1.75 fl. oz.	0.028		15.4 fl. oz.	0.250
1.9. fl. oz.	0.031		30.75 fl. oz.	0.500
2.85 fl. oz.	0.046		46.0 fl. oz.	0.748
2.88 fl. oz.	0.047			

Drench Application

Apply this product as a preventative, drench treatment prior to infection to control soilborne, seedling, and crown diseases of production ornamentals (greenhouse, shadehouse, and container grown). Good coverage of the pre-infection area (root zone, root ball, crown, etc.) is necessary for satisfactory control.

This product may be applied as a drench to container grown ornamentals using 0.38 – 1.75 fl. oz. / 100 gallons of water prior to infection as healthy roots are necessary to optimize product uptake, systemic translocation and disease protection. Apply 1-2 pints of the solution per square foot surface area at 7-28 day intervals.

Drench applications may cause phytotoxicity in small bedding plants in the seedling/plug stage. Test this product on a small number of plants before applying on a larger scale.

Drip Irritation

Apply this product through drip irrigation systems to potted ornamentals or to bedded, field grown ornamentals for soil-borne disease control. Apply at the rate of 3.85 – 30.75 fl. oz. per acre as a preventative disease application. Ensure that the soil or potting media has adequate moisture before making the drip application.

Drip irrigation must be terminated when the fungicide is depleted from the main feed supply tank or 6 hours after starting irrigation, whichever is shorter. Delay a subsequent irrigation (water only) for at least 24 hours following the drip application to ensure maximum efficacy.

ORNAMENTAL USE PRECAUTIONS

- This product may be applied to certain varieties of crabapple for control of apple scab. It has been shown to be safer when applied to the species and varieties listed in Table 3, however, due to the large number of genera, species, and varieties of crabapple, it is impossible to test every one for tolerance to this product. The professional user must conduct small scale testing to ensure plant safety prior to broad scale commercial use on plant genera and species.
- Use only surfactants approved for use on ornamental plants in combination with this product. Prior to broad scale use, test tank mixes on a small group of representative plants.
- This product may not provide adequate disease control when applied after disease outbreak.
- Test this product on a small number of plants before applying on a larger scale.

ORNAMENTAL USE RESTRICTIONS

- For all ornamental Uses (NOT for Field-grown or Nursery ornamentals (ground and chemigation applications), any hand-held applications and mechanically pressurized handwand applications):
 - **DO NOT** apply more than 2.4 gallons (5.0 lbs. AI) of this product per acre per year.
 - **DO NOT** apply more than 8 applications per year.
 - The Maximum Single Application rate is indicated in the chart by ^M following the rate (for example 7.7^M fl.oz.)
 - The Minimum Retreatment Interval is indicated in the chart by ^R following the interval range (for example 7^R – 28 days).
- For Field-grown or Nursery ornamentals (ground and chemigation applications):
 - **DO NOT** apply more than 0.36 gallons (46 fl. oz. or 0.75 lbs. AI) of this product per acre per year.
 - **DO NOT** apply more than 8 applications per year.
- For Nursery ornamentals broadcast application and Landscape (plants, flowers, trees, turf), (any hand-held applications) and Greenhouse ornamentals (mechanically pressurized handwand applications):
 - **DO NOT** apply more than 0.0012 gallons (0.15 fl. oz. or 0.0025 lbs. AI) of this product per acre per year.
 - **DO NOT** apply more than 8 applications per year.
- **DO NOT** apply more than 600 gallons spray volume per acre for foliar applications.
- **DO NOT** apply more than 2 pints per square foot for drench and crown applications.
- Apply by ground (includes any hand-held and mechanically pressurized handwands) **only**.
- **DO NOT** apply this product to apple or cherry trees (flowering, Yoshina variety) due to possible phytotoxicity. **DO NOT** use spray equipment that has applied this product for use on these sensitive crops due to possible phytotoxicity from residue remaining in the sprayer.
- **DO NOT** use silicone based products with this product due to possible phytotoxicity.
- **DO NOT** tank mix this product with other pesticides, fertilizers, adjuvants, etc., unless testing or local knowledge indicates that the tank mixture is safe when used on ornamental plants.
- **DO NOT** make more than three sequential drench applications of this product before alternating with a fungicide of a different mode of action.
- **DO NOT** alternate with other strobilurin fungicides.

TABLE 1 - DISEASES CONTROLLED

When used according to the label directions, this product provides control of the following diseases of ornamental plants:

Target Diseases	Use Rates and Application Directions	
	8 oz. and larger Containers fl. oz. product per 100 gallons	4 oz. Containers fl. oz. product per 50 gallons
1. Conifer Blights		
1a. Phomopsis Blight (<i>Phomopsis juniperovora</i>)	Apply 1.9 – 7.7 ^M fl. oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl. oz. at 7 ^R to 28-day intervals.
1b. Tip Blight (<i>Sirococcus strobilinus</i>)		
2. Leaf Blights/Leaf Spots		
2a. Alternaria Leaf Spot (<i>Alternaria</i> spp.) 2b. Anthracnose (<i>Colletotrichum</i> spp., <i>Elsinoe</i> spp.) 2c. Entomosporium Leaf Spot (<i>Entomosporium mespili</i>) 2d. Leaf spot (<i>Cladosporium echinulatum</i>) 2e. Downy Mildew of Bedding Plants (<i>Peronospora</i> spp.) 2f. Cercospora Leaf Spot (<i>Cercospora</i> spp.)	Apply 1.9 – 7.7 ^M fl. oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl. oz. at 7 ^R to 28-day intervals.
2h. Marssonina Leaf Spot (<i>Marssonina</i> spp.)	Apply 1.9 – 7.7 ^M fl. oz. at 14 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl. oz. at 14 ^R to 28-day intervals.
2i. Downy Mildew of Rose (<i>Peronospora sparsa</i>)	Apply 3.85 – 7.7 ^M fl. oz. at 7 ^R to 21-day intervals during periods of active plant growth and prior to dormancy or severe infection.	Apply 1.95 – 3.85 ^M fl. oz. at 7 ^R to 21-day intervals during periods of active plant growth and prior to dormancy or severe infection.
2j. Iris Leaf Spot (<i>Mycosphaerella macrospora</i>) 2k. Myrothecium leaf spot (<i>Myrothecium</i> spp.)	Apply 3.85 – 7.7 ^M fl. oz. at 7 ^R to 21-day intervals.	Apply 1.95 – 3.85 ^M fl. oz. at 7 ^R to 21-day intervals.
2l. Rose Blackspot (<i>Diplocarpon rosea</i>)	Apply 7.7 – 15.4 ^M fl. oz. at 7 ^R to 14-day intervals. If disease pressure is light, apply at 7-day intervals. This product may be tank-mixed with another Rose Blackspot fungicide if disease conditions are severe. DO NOT exceed 46 fl. oz. of product per acre.	Apply 3.85 – 7.7 ^M fl. oz. every 7 ^R -14 days. If disease pressure is light, apply at 7-day intervals. This product may be tank-mixed with another Rose Blackspot fungicide if disease conditions are severe. DO NOT exceed 46 fl. oz. of product per acre.
2m. Scab (<i>Venturia inaequalis</i>)	Apply 1.9 – 7.7 ^M fl. oz. at 10 ^R to 28-day intervals. DO NOT apply to apple trees. Refer to Table 3 for tolerant species of crabapples.	Apply 0.95 – 3.85 ^M fl. oz. at 10 ^R to 28-day intervals. DO NOT apply to apple trees. Refer to Table 3 for tolerant species of crabapples.
3. Powdery Mildew		
3a. <i>Erysiphe pannosa</i> , <i>E.</i> spp. 3b. <i>Microsphaera azalea</i> 3c. <i>Sphaerotheca pannosa</i>	Apply 1.9 – 7.7 ^M fl. oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl. oz. at 7 ^R to 28-day intervals.

Target Diseases	Use Rates and Application Directions	
	8 oz. and larger Containers fl. oz. product per 100 gallons	4 oz. Containers fl. oz. product per 50 gallons
4. Rusts		
4a. Needle Rust (<i>Melampsora occidentalis</i>) 4b. <i>Phragmidium</i> spp. 4c. <i>Puccinia</i> spp. 4d. <i>Gymnosporangium</i> spp.	Apply 1.9 – 7.7 ^M fl. oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl. oz. at 7 ^R to 28-day intervals.
5. Flower Blights		
5a. Anthracnose (<i>Collectotrichum</i> spp., <i>Elsinoe</i> spp.)	Apply 1.9 – 7.7 ^M fl. oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 3.85 ^M fl. oz. at 7 ^R to 28-day intervals.
5b. Botrytis Slight (<i>Botrytis cinerea</i>) – Suppression only	Apply 7.7 – 15.4 ^M fl. oz. at 7 ^R to 21-day intervals. DO NOT exceed 46 fl. oz. of product per acre.	Apply 3.85 – 7.7 ^M fl. oz. at 7 ^R to 21-day intervals. DO NOT exceed 46 fl. oz. of product per acre.
6. Shoot/Stem Diseases		
6a. Aerial/Shoot Blight (<i>Phytophthora</i> spp.)	Apply 1.9 – 3.85 ^M fl. oz. at 7 ^R to 28-day intervals.	Apply 0.95 – 1.9 ^M fl. oz. at 7 ^R to 28-day intervals.
7. Soilborne Diseases (Directed Spray)		
7a. <i>Rhizoctonia solani</i> 7b. <i>Sclerotium rolfsii</i> 7c. <i>Rosarium</i> spp.	Apply 1.9 – 7.7 ^M fl. oz. at 7 ^R to 21-day intervals.	Apply 0.95 – 3.85 ^M fl. oz. at 7 ^R to 21-day intervals.
8. Soilborne Diseases (Drench)		
8a. <i>Rhizoctonia solani</i> 8b. <i>Sclerotium rolfsii</i> 8c. <i>Fusarium</i> spp.	Apply 0.35 – 1.75 ^M fl. oz., 1 -2 pints of the solution per square foot of surface area, at 7 ^R to 28-day intervals.	Apply 0.19 – 0.95 ^M fl. oz., 1-2 pints of the solution per square foot of surface area, at 7 ^R to 28-day intervals.

^M The Maximum Single Application rate is indicated in the chart by ^M following the rate (for example 7.7^M fl.oz.)

^R The Minimum Retreatment Interval is indicated in the chart by ^R following the interval range (for example 7^R – 28 days).

PLANT SAFETY

This product has been shown to be safe when applied to the ornamental plants listed in Tables 2 and 3. However due to the large number of genera, species and varieties of ornamental and nursery plants; it is impossible to test every one for tolerance to this product. Neither the manufacturer nor the seller has determined whether or not this product can be used safely on all genera, species, or varieties of ornamental and nursery plants. The professional user must conduct small scale testing to ensure plant safety prior to broad scale commercial use on plant genera and species not listed in this label.

In addition, **DO NOT** tank mix this product with other pesticides, fertilizers, adjuvants, etc, unless local experience indicates that the tank mix is safe to ornamental plants.

DO NOT apply or use spray equipment used to make applications of this product to certain apple, crabapple or cherry trees and other sensitive crops due to possible phytotoxicity.

Table 2 - Tolerant Ornamental Plants and Diseases Controlled

When applied to the plants listed in Tables 2 and 3 at the listed rates and according to the application directions on this label, this product has been found to be safe and effective at controlling the listed diseases.

Common Name	Botanical Name	Diseases/Pathogens (Refer to Table 1)
Abelia	Abelia spp.	2
Alder (White), Clethra	Clethra alnifolia	2
Arborvitae	Thuja spp.	2
Aspen trees	Populus spp.	2
Aster, Starwort	Aster, spp.	4
Azalea, Glacier	Rhododendron spp.	2b, 3, 6, 7
Azaleas, Rhododendron	Rhododendron spp.	2b, 3, 6, 7
Barberry	Berberis thunbergii	3, 4
Australian Laurel	Pittosporum spp.	3, 4
Baby Rubber-plant	Peperomia spp.	2, 7
Begonia	Begonia spp. (except Reiger begonia)	2, 3
Birch (River)	Betula nigra	3, 4
Black-eyed Susan	Rudbeckia hirta	2
Blanket-Flower	Gaillardia spp.	2
Bougainvillea	Bougainvillea spp.	2
Boxwood	Buxus sempervirens	2, 7a
Bradford's Pear	Pyrus cafleryana	3
Buddleia, Butterfly Bush	Buddleia davidii	2
Bugle, Bugleweed	Ajuga reptans	3
Burning Bush	Euonymus alatus	2
Caladium	Caladium spp.	7
Camellia	Camellia japonica	2
Carnation	Dianthus caryophyllus	3, 4
Cedar (Atlas)	Cedrus atlantica	2, 4
Cedar (Red)	Juniperus virginiana	1a, 4
Cedar, Western Red	Thuja plicata	4
Cedar (White)	Cedrus spp.	2, 4
Cherry	Prunus pumila	2, 5
Chinese evergreen	Aglaonema spp.	2, 4
Chrysanthemums	Chrysanthemum spp.	2, 7c
Cinquefoil	Potentilla spp.	2

Common Name	Botanical Name	Diseases/Pathogens (Refer to Table 1)
Cotoneaster (Creeping)	Cotoneaster adpressus	7
Cotoneaster (Variegated Rockspray)	Cotoneaster horizontalis	7
Crabapple (See Table 3 for variety list)	Malus spp.	2i
Cranesbill	Geranium spp.	5b
Crapemyrtle	Lagerstroemia indica	2, 3
Creeping thyme	Thymus sagahyifam	4
Cyclamen	Cyclamen spp.	7c
Cyperus	Cyperus spp.	1
Cypress (Sawara)	Chamaecyparis pisifera	1
Cypress, Leyland cypress	Chamaecyparis spp.	1
Daisy (Gerber, Transvaal)	Gerbera jamesonii	3
Dogwood	Cornus florida	2b, 3
Dogwood, Pink Dogwood, Flowering Dogwood	Cornus spp.	2b, 3
Dwarf Pampas Grass	Phelans spp.	3
Dumb cane	Dieffenbachia spp.	2
Euonymus (Dwarf Winged)	Euonymus alata	2
Euonymus (Evergreen)	Euonymus japonicas	2
Fatsia (Japanese), Paper-plant	Fatsia japonica	2
Fig	Ficus spp.	2
Fir, Douglas	Pseudotsuga spp.	1, 4
Fir (Fraser)	Abies fraseri	1, 4
Fir (Noble)	Abies procera	1, 4
Floss-flower	Ageratum spp.	3, 4
Forsythia	Forsythia viridissima	2
Foxglove	Digitalis spp.	2, 3
French hydrangea	Hydrangea macrophylla	2, 3
Gardenia	Gardenia jasminoides	3
Geranium	Pelargonium spp.	3, 4, 5b
Grass	Permisetum alopecuriodes	2
Hydrangea	Hydrangea spp.	2, 3
Heather	Erica dareyensis	2
Hibiscus	Hibiscus moscheutos	2, 3
Hemlock	Tsuga spp.	4
Hibiscus	Hibiscus rosa-sinensis	2, 3
Holiday cactus	Schlumbergera	2, 7
Holly, Winterberry, Yaupon	Ilex spp.	3
Hosta	Hosta spp.	2
Impatiens, Balsam	Impatiens spp.	2a, 7a
Indian Hawthorn	Phaphiplepsisindica	2, 3, 4
Iris (African, Butterfly)	Dietes iridiodes	4c
Iris (bulbous, Spanish, Dutch)	Iris xiphium	2e
Ivy (Algerian)	Hedera algeriensis	2
Ivy (English)	Hedera helix	2
Ivy, Swedish Coleus	Plectranthus spp.	2
Japanese Andromeda	Pieris japonica	2, 7
Japanese aucuba, Japanese laurel	Aucuba japonica	7
Juniper	Juniperus procumbens	1a, 4
Juniper	Juniperus scopulorum	1a, 4
Juniper	Juniperus spp.	1a, 4

Common Name	Botanical Name	Diseases/Pathogens (Refer to Table 1)
Larkspur	Delphinium spp.	2
Laurel	Laurus nobilis	3
Lilac (wild)	Ceanothus sanguineus	3
Lily (Asiatic)	Lilium spp.	2
Lily-turf	Liriope muscari	2
Live-forever, House-Leek	Sempervivum spp.	2
Magnolia, Southern	Magnolia grandiflora	2
Magnolia, Saucer	Magnolia soulangiana	2
Magnolia	Magnolia spp.	2
Maple (Japanese)	Acer palmatum	2
Maple (Sugar)	Acer saccharum	2
Marigold	Tagetes spp.	2a
Mock-orange	Philadelphus	3, 4
Muhgo pine	Pinus muhgo	1b, 4
Mugwort, Sagebrush	Artemisia spp.	2
Nandina	Nandina domestica	2
Oak, pin	Quercus palustris	2, 3
Oak, red	Quercus falcate	2, 3
Oleander, Rose-bay	Nerium oleander	2
Orpine, Stonecrop	Sedum spp.	2i
Palm, date	Phoenix dactylifera	2, 7
Palm (Parlor)	Chamaedora elegans	7
Palm, Queen	Syagrus romanzollianum	2
Palm, Roebelin's	Phoenix roebelenii	2, 7
Palm (Sago)	Caryota urens	2, 7
Pampas Grass	Cortaderia selloana	3
Peace lily	Spathiphyllum floribundium	2, 7
Periwinkle	Vinca spp.	2, 6a
Petunia	Petunia spp.	6a
Philodendron	Philodendron spp.	2j
Phlox	Phlox spp.	3
Pine, Black	Pinus nigra	1b, 4
Pine, Eastern White	Pinus strobes	1b, 4
Pine, Scotch	Pinus silvestris	1, 4
Pine	Pinus spp.	1b, 4
Pink	Dianthus spp.	3, 4
Plum, Flowering; Purple-leaf	Prunus spp.	2, 5
Poinsettia	Euphorbia spp.	2a
Poplar	Populus trichocarpa	4
Pothos	Epipremnum spp.	2
Primrose	Primula spp.	2
Pussy's Foot	Ageratum spp.	3, 4
Redbud (Western)	Cercis occidentalis	2
Red tip photinia	Photinia glabra	2, 3, 4
Ribbon Grass	Setaria spp.	2, 3
Rose	Rosa spp.	2a, 2c, 3c, 4b
Rose of Sharon	Hibiscus syriacus	2, 3
Rosemary (prostrate)	Rosmarinus spp.	2
Rubber-tree, Umbrella-tree	Brassaia actinophylla	2, 7
Sage	Salvia spp.	3, 4j

Common Name	Botanical Name	Diseases/Pathogens (Refer to Table 1)
Snapdragon	Antirrhinum spp.	3, 4
Snowball, Ceanothus, California lilac	Ceanothus spp.	3
Spirea	Spirea budalda	3
Spirea	Spirea japonica	3
Spreading yew	Taxus baccata	7
Spruce, Blue	Picea purtgens	1
Spruce, Norway	Picea abies	1
Spruce, White	Picea glauca	1
Sweet Alyssum	Lobularia maritima	7
Verbena	Verbena spp.	3
Vervain	Verbena spp.	3
Viburnum	Viburnum spp.	2, 3, 4
Vinca	Catharanthus roseus	2
Viola, Pansy	Viola spp.	1, 2
Virginia Willow	Itea virginica	3, 4
Vinca	Catharanthus roseus	2
Viola, Pansy	Viola spp.	1, 2
Virginia Willow	Itea virginica	3, 4
Western hemlock	Tsuga heiarophyllia	4
Wiegela (Pink)	Wiegela florida	2
Wormwood	Artemisia spp.	2
Yucca	Yucca spp.	7
Zebra Plant	Aphelandra spp.	2
Zinnia	Zinnia sp.	2a, 3

* **DO NOT** exceed 3.85 fl. oz./100 gallons on these species.

TABLE 3. Tolerant Varieties of Crabapple Species (Genus Malus)

Arkansas Black	Eleyi	Mary Potter	Seiboldii
Atrosanguinea	Enterprise	Molten Lava	Selkirk
Baccafa	Evereste	New Centennial	Sentinel
Baccata var. jackii	Eyeiynn	Ormiston Roy	Silver Moon
Baccata var. mandshurica	Floribunda	Pink Satin	Siiverdrift
Callaway	Gloriosa	Prairie Maid	Sinai Fire
Candymint Sargent	Golden Delicious	Prairifire	Spectabfis
Christmas Holly	Golden Raindrops	Profusion	Sugar Tyme
Coronaria	Hopa	Pumila	Van Eseltine
David	Indian Magic	Ralph Shay	White Angel
Dolgo	Island	Red Jade	Williams Pride
Donald Wyman	Katherine	Red Baron	Winter Gold
Dorothea	Lancelot	Sargent	Yellow Delicious
Doublings	Louisa	Sargentii	Zumi Calocarpa

TABLE 4. Intolerant Plants. DO NOT apply this product to these species or varieties.

COMMON NAME	BOTANICAL NAME
Apple	Malus domestics
Crabapple - Flame variety	Malus spp.
Crabapple - Brandywine variety	Mals spp.
Crabapple - Novamac variety	Malus spp.
Cherry, Flowering - Yoshina variety	Prunus yedoensis
Leatherleaf Fern and Other Ferns for cut foliage	Rumohra adianformis and other species for cut foliage
Privet	Ligusirum spp.

CONIFERS INCLUDING CHRISTMAS TREES, COMMERCIAL PRODUCTION ROSES [(Restriction: Not for use in California)]

Use this product to control diseases on conifers in production (indoor and outdoor) and in landscape situations. Please see the Ornamental Section above for more detailed directions for use in landscape situations.

CONIFERS, including Christmas Trees		
Target Diseases	Rate per Acre (lbs. AI/A)	Application Directions
Diplodia tip blight (<i>Diplodia pinea</i>) Lophodermium Needlecast (<i>Lophodermium pinastri</i>) Swiss Needlecast (<i>Phaeocryptopus gaumannif</i>)	6.0 – 15.5 (0.10 – 0.25)	Apply this product before disease outbreak and continue throughout the season at 7- to 21-day intervals following resistance management guidelines. Apply this product by ground, air or chemigation. If an adjuvant is used, add it at the manufacturer’s specified rates. Include this product in an IPM program, which includes alternating fungicides with different modes of action and selections of varieties with disease resistance and removal of plant debris in which inoculum may overwinter. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 123 fluid ounces (2.0 lbs. AI) of this product per acre per year. • DO NOT make more than eight applications of this product per acre per year at the high rate (0.25 lbs. AI). • Minimum Retreatment Interval: 7 days 		

CHRISTMAS TREES (Any hand-held applications)		
Target Diseases	Rate per Acre (lbs. AI/A)	Application Directions
Diplodia tip blight (<i>Diplodia pinea</i>) Lophodermium Needlecast (<i>Lophodermium pinastri</i>) Swiss Needlecast (<i>Phaeocryptopus gaumannif</i>)	0.005 – 0.01 fl. oz. (0.0001 – 0.0002)	Apply this product before disease outbreak and continue throughout the season at 7- to 21-day intervals following resistance management guidelines. If an adjuvant is used, add it at the manufacturer’s specified rates. Include this product in an IPM program, which includes alternating fungicides with different modes of action and selections of varieties with disease resistance and removal of plant debris in which inoculum may overwinter. DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.
RESTRICTIONS: <ul style="list-style-type: none"> • DO NOT apply more than 0.01 fl. oz. (0.0002 lbs. AI) of this product per acre per application. • DO NOT apply more than 0.08 fluid ounces (0.00125 lbs. AI) of this product per acre per year. • DO NOT make more than eight applications of this product per acre per year at the high rate (0.0002 lbs. AI). • Minimum Retreatment Interval: 7 days 		

ROSES, Commercial Rose Production		
Target Diseases	Rate per Acre fl. oz./Acre (lbs. AI/A)	Application Directions
Downy Mildew (Peronospora sparsa) Powdery Mildew (Sperotheca pannosa) Rust (Phragmidium mucronatum, P. tubercalutum, and other Phragmidium spp.) Septoria Leaf Spot (Septoria rosea) Alternaria Leaf Spot (Alternaria alternate)	3.0 – 15.5 (0.05 – 0.25)	<p>Apply this product before disease outbreak and continue throughout the year at 7– to 21 day intervals following resistance management guidelines.</p> <p>Apply this product by ground, air or chemigation. If an adjuvant is used, add it at the manufacturer’s specified rates.</p> <p>Include this product in an IPM program, which includes alternate fungicides with different modes of action and selection of varieties with disease resistance, proper fertilizer application, winter and/or spring pruning, management of plant residue, and proper irrigation timing and application.</p> <p>Azoxystrobin has been shown to be safe when applied to roses, however, all varieties of roses have not been tested. Test the product first on a smaller scale to ensure its safety prior to making a broadscale application.</p> <p>DO NOT tank mix this product with other pesticides, fertilizers, etc. unless testing or local knowledge indicates that the tank mixture is safe when used on roses.</p> <p>DO NOT make more than two applications of this product before alternating with fungicides with a mode of action other than QoI Group 11.</p>
<p>Restrictions:</p> <ul style="list-style-type: none"> • DO NOT apply more than 15.5 fl. oz. (0.25 lbs. AI) of this product per acre per application. • DO NOT apply more than 123 fluid ounces (2.0 lbs. AI) of this product per acre per year. • DO NOT make more than eight applications of this product per acre per year at the high rate (0.25 lbs. AI). • Minimum Retreatment Interval: 7 days 		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

STORAGE

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER HANDLING [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

CONTAINER HANDLING [Bulk/Mini-Bulk] [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER HANDLING [Bulk/Mini-Bulk] [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container or pressure rinse promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full of water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

WARRANTY AND LIMITATION OF DAMAGES

CONDITIONS OF SALE: To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. SIPCAM AGRO USA, INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED. To the extent consistent with applicable law, SIPCAM AGRO USA, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND SIPCAM AGRO USA, INC.'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. SIPCAM AGRO USA, INC. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.