



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

60063-100

Date of Issuance:

9/5/25

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Azoxystrobin + Prothioconazole
Turf

Name and Address of Registrant (include ZIP Code):

Sipcam Agro USA, Inc.
2525 Meridian Parkway
Durham, NC 27713

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. You have 18 months from the date of registration to provide these data.

Continues page 2

Signature of Approving Official:

Stephanie Suarez, Ph.D., Acting Product Manager 22
Fungicide Branch, Registration Division (7505T)

Date:

9/5/25

EPA Form 8570-6

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 60063-100."
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The record for this product currently contains the following CSF(s):

- Basic CSF dated 09/16/2024
- Alternate CSF(s) 1-15 dated 9/16/2024

If you have any questions, please contact Thomas Harty at 202-566-0394 or at harty.thomas@epa.gov.

Enclosure

Azoxystrobin	Group	11	Fungicide
Prothioconazole	Group	3	Fungicide

Prothioconazole + Azoxystrobin Turf

[For control of listed diseases on Golf Course Turfgrass]
Intended for use by professional applicators.

Active Ingredients:

Azoxystrobin: methyl (E)-2-[2-[6-(2-cyanophenoxy)pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate 17.54%

Prothioconazole: 15.79%

Other Ingredients: 66.67%

Total: 100.0%

Contains 1.642 lbs. Azoxystrobin and 1.478 lbs. Prothioconazole per gallon. Suspension Concentrate.

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 222-1222 Poison Control Center


Net Contents: ____ [gallons] [gal.] [(Liters)(L)]

EPA Est. No.: _____

Lot number [begins with] [ends with] xx]

EPA Reg. No. 60063-XX

[lot number / label date code]



Manufactured for:
Sipcam Agro USA, Inc.
2525 Meridian Parkway
Durham, NC 27713

[Alternate Brand Name: Carve Xtra]

Read the [entire] label [carefully] before [opening the container] [using this product.]
See additional Precautionary Statements and Directions for Use inside booklet.

Optional Language for label

[Fungicide]

[Open book here] [Open label here] [Pull back book here] [Pull back label here]

[Application Type [T] [Turf]]

[Formulated in the United States of America, with U.S. and imported ingredients.]

[Product of ____] [Note: if manufactured in a country other than U.S., country name will appear here]

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Human flagging is prohibited.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

USER SAFETY REQUIREMENTS

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to freshwater and estuarine/marine fish and aquatic invertebrates, and freshwater/estuaries/marine aquatic plants. Azoxystrobin can be persistent for several months or longer.

DO NOT discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. **DO NOT** discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or regional office of the EPA.

For terrestrial uses **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

Azoxystrobin (and a degradate of azoxystrobin) and degradates of prothioconazole are known to leach through soil into ground water under certain conditions as a result of label use. This product may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Notify State and/or Federal authorities and Sipcam Agro USA, Inc. immediately if you observe any adverse environmental effects due to use of this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Golf courses are not within the scope of the Worker Protection Standard.

DO NOT enter or allow others to enter treated areas until sprays have dried.

PRODUCT INFORMATION

This product is a broad spectrum, preventative fungicide with systemic and curative properties. This product may be applied as a foliar spray in alternating spray programs or in tank mixes with other registered crop protection products. All applications must be made according to the use directions that follow. Equipment must be properly calibrated before use.

RESTRICTIONS

- For use on golf course turf grass only
- For use only by professional applicators
- This product is extremely phytotoxic to certain apple varieties. AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees and apple fruit. **DO NOT** spray this product where spray drift may reach apple trees. **DO NOT** use spray equipment which has been previously used to apply this product to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity to certain apple and crabapple varieties.

SPRAY DRIFT

Groundboom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT - Ground Boom

- For ground equipment, the boom should remain level and have minimal bounce.

RELEASE HEIGHT - Aircraft

- Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

- When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

- Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

- Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.
- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

HANDHELD TECHNOLOGY APPLICATIONS

- Take precautions to minimize spray drift.

MIXING, LOADING, AND APPLYING

This product is intended to be diluted into water and then applied by typical spraying techniques. Always apply this product in sufficient water to obtain thorough, uniform coverage of foliage intended to be protected from disease. For spray volumes, see the turf use charts.

Slowly invert container several times to assure uniform mixture. Measure the required amount of this product and pour into the spray tank during filling. Keep agitator running when filling spray tank and during spray operations.

It is necessary to thoroughly apply the product in order to provide good disease control. **DO NOT** prepare more spray solution than is needed for application. Avoiding spray overlap will reduce the potential for injury.

Tank Mixing

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

When tank mixing this product with other pesticides, observe the more restrictive label limitations and precautions. **DO NOT** exceed any label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

DO NOT combine this product in the sprayer tank with pesticides, surfactants or fertilizers, unless prior use has shown the combination to be physically compatible, non-injurious and effective under similar use conditions.

DO NOT combine this product with other products that contain *Bacillus thuringiensis* subsp. *kurstaki* strain ABTS-351, as the combination may result in phytotoxicity when applied to the crops listed on this label.

DO NOT tank mix this product with oil or with any adjuvants which contain oil as their principal ingredient.

When mixed with EC (emulsifiable concentrate) formulations, this product may be phytotoxic to other crops listed on this label, especially when applied during cool, cloudy conditions that last for several days. Combination in the spray tank with other pesticides, fertilizers, or surfactants is not recommended unless prior use has shown the combination to be physically compatible, effective, and non-injurious under your conditions of use. Adjuvants containing silicone could also have phytotoxic effects. When an adjuvant is used with this product, use a Council of Producers and Distributors of Agrotechnology (CPDA) certified adjuvant.

Compatibility Test for Mix Components: Before mixing components, always perform a compatibility jar test. To determine the physical compatibility, the recommended proportions of products must be added into a suitable container of water in the following sequence:

1. Products packaged in water soluble packaging
2. Wettable powders, wettable granules (dry flowables)
3. Prothioconazole + Azoxystrobin
4. Other liquid flowables
5. Emulsifiable concentrates
6. Water soluble liquids
7. Adjuvants

Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add in the sequence indicated above. Always mix between component additions. After thoroughly mixing, let stand for at least 10-15 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

The safety of all potential tank-mixes, including additives and other pesticides, on all turf types, has not been tested. Before applying any tank-mixture not specifically listed on this label, the safety to the target area must be confirmed. To test for safety, apply the combination to a small area of the target in accordance with the label instructions to ensure that a phytotoxic response will not occur.

INTEGRATED PEST / DISEASE MANAGEMENT

This product provides excellent control of fungal diseases when used according to label directions for control of a broad spectrum of plant diseases. This product is for use in programs that are compatible with the principals of Integrated Pest Management (IPM), including the use of disease resistant varieties, cultural practices, pest scouting, and disease forecasting systems which reduce unnecessary applications of pesticides.

RESISTANCE MANAGEMENT

For resistance management, this product contains prothioconazole, a DMI Group 3 Fungicide, and azoxystrobin, a QoI Group 11 fungicide. Any fungal population may contain individuals naturally resistant to this product, and other Group 3 or Group 11 fungicides. A gradual or total loss of pest control

may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

Fungal isolates with acquired resistance to Group 11 may eventually dominate the fungal population if Group 11 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. Cross resistance has been shown between all members of the Qol fungicides. Since Qol fungicides are a high risk for resistance, this may result in partial or total loss of control of those species.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 3 or Group 11 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological, and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Sipcam Agro at 919-226-1195. You can also contact your pesticide distributor or university extension specialist to report resistance.

Follow the specific resistance management guidance listed in the Turf Use Directions table. If resistance management guidance is not specified in the Turf Use Directions table, then follow the guidance provided in the table below.

Total fungicide applications planned	1	2	3	4	5	6	7	8	9	10	11	12
Applications of Qol fungicides applied alone	1	1	2	2	2	2	2	3	3	3	3	4
Applications of Qol fungicides applied in mixture (tank-mix or formulated)	1	2	2	2	2	3	3	4	4	5	5	6

When multiple applications are required during the year, spray programs for Group 11 (Qol) fungicides must be developed. When two sequential applications of Group 11 fungicides are made, they must be alternated with two or more applications of a fungicide that is not a Group 11 fungicide.

TURF USE DIRECTIONS

During conditions which are favorable to prolonged periods of fungal infection use another registered fungicide for additional applications if maximum amount of this product has been applied. Efficacy for certain diseases may be reduced if resistant isolates to Group 3 or Group 11 fungicides are present. Use this product in an IPM program, alternating fungicides with different modes of action. When environmental conditions are favorable to disease, during period of heavy disease pressure, or with highly susceptible varieties, use the higher listed rates in the rate range and/or shorter spray intervals.

When disease suppression is referred to, suppression can mean either erratic control from good to fair or consistent control at a level below that obtained with the best commercial disease control products.

Note: The application rate of 22.5 fl.oz./A is equivalent to 0.289 lbs./A Azoxystrobin and 0.260 lbs./A Prothioconazole.

Disease	Application rate	Application interval	Useful Information
Anthracnose (<i>Colletotrichum graminicola</i>) Brown Patch (<i>Rhizoctonia solani</i>) Brown Ring (<i>Waitea</i>) Patch (<i>Waitea circinata</i> var. <i>circinata</i>) Dollar Spot (<i>Sclerotinia homoeocarpa</i> syn. <i>Clariireedia homoeocarpa</i>) Gray Leaf Spot (<i>Pyricularia grisea</i>) Mini Ring (<i>Rhizoctonia zeae</i>)*	22.5 fl.oz./A 0.517 fl.oz./1000 sq.ft.	14 – 21 Days	Apply preventatively before conditions are favorable for infection occur. Use the shorter application interval when favorable turf disease conditions persist.
Fairy Ring (caused by Basidiomycete fungi)	22.5 fl.oz./A 0.517 fl.oz./1000 sq.ft.	28 Days	Apply this product as soon as possible after fairy ring symptoms develop. Irrigate with at least 0.5 inches water within 24 hours after application if rainfall is not expected. Severely damaged or thin turf may require reseeding. Fairy ring symptoms may take 2 to 3 weeks to disappear following application. Make an additional application 14-28 days if needed. Minimum Retreatment Interval: 28 days
Large Patch (<i>Rhizoctonia solani</i>)	22.5 fl.oz./A 0.517 fl.oz./1000 sq.ft.	14 to 28 Days	Water-in applications to move fungicide into crown area and root zone. Use the shorter application interval when favorable turf disease conditions persist.
Microdochium Patch (<i>Microdochium nivale</i>)	22.5 fl.oz./A 0.517 fl.oz./1000 sq.ft.	14 to 21 Days	Apply when conditions are conducive to disease development. Use the shorter application interval when favorable turf disease conditions persist.
Snow Mold (<i>Microdochium nivale</i> , <i>Typhula incarnata</i> , <i>T.ishikariensis</i>)	22.5 fl.oz./A 0.517 fl.oz./1000 sq.ft.	14 to 28 Days	Make 1-2 preventative applications before conditions favorable for infection occur, before snow cover. Use the shorter application interval when

			favorable turf disease conditions persist.
Spring Dead Spot (<i>Ophiosphaerella korrae</i> , <i>O. herpotricha</i> , <i>Leptosphaeria korrae</i> , <i>L. narmari</i>)	22.5 fl.oz./A 0.517 fl.oz./1000 sq.ft.	28 Days	Water-in applications to move fungicide into crown area and root zone.
Summer Patch (<i>Magnaporthe poae</i>)	22.5 fl.oz./A 0.517 fl.oz./1000 sq.ft.	14 to 28 Days	Use higher spray-volume or water-in applications to crowns and upper roots for maximum effectiveness. Use the shorter application interval when favorable turf disease conditions persist.
Take-all Root Rot (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	22.5 fl.oz./A 0.517 fl.oz./1000 sq.ft.	28 Days	Apply in adequate water volume or water-in applications to upper root zone.
RESTRICTIONS: <ul style="list-style-type: none"> • MAXIMUM SINGLE APPLICATION RATE: 22.5 fl.oz product/A (0.823 lbs. azoxystrobin/A, 0.141 lbs. prothioconazole/A) • MAXIMUM ANNUAL APPLICATION RATE PER YEAR: 112.5 fl.oz product/A (4.115 lbs. azoxystrobin/A, 0.705 lbs. prothioconazole/A) • MAXIMUM NUMBER OF APPLICATIONS: 5 • DO NOT apply more than 4.68 lbs. azoxystrobin per acre per year. • DO NOT apply more than 0.801 lbs. prothioconazole per acre per year. • For use by professional applicators only. • For use only on golf course turfgrass (greens, tees, fairways, and roughs). • Not for residential use. DO NOT use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational parks, athletic fields, schools, campgrounds, churches, and theme parks. • DO NOT use clippings for animal feed. • DO NOT apply by aerial applications. • DO NOT apply through any type of irrigation equipment (chemigation). • For ground application equipment only. DO NOT use with handheld application equipment, including mechanically pressurized spray guns, backpack or tank pressurized spray guns, spray hawk, or handheld boom applicators. • [Not for sale, distribution, or use in Nassau or Suffolk counties, New York State.] 			

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

STORAGE

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER HANDLING [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

CONTAINER HANDLING [Bulk/Mini-Bulk] [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER HANDLING [Bulk/Mini-Bulk] [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container or pressure rinse promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full of water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

WARRANTY AND LIMITATION OF DAMAGES

CONDITIONS OF SALE: To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. SIPCAM AGRO USA, INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED. To the extent consistent with applicable law, SIPCAM AGRO USA, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND SIPCAM AGRO USA, INC.'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. SIPCAM AGRO USA, INC. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.