

61178-5

01/7/2010

1/83

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Mr. Robert Prince  
President, Microgen, Inc  
33 clinton Rd., Suite 102  
West Caldwell, NJ 07006

JAN -7 2010

Subject: **CCX-151**  
EPA Registration Number 61178-5  
Notification Application Dated December 7, 2009  
EPA Received Date December 14, 2009

Dear Mr. Prince:

This will knowledge receipt of your notification, submitted under the provisions of FIFRA section 3c 9. Based on a review of the submitted material the following comment apply.

**Proposed Amendment:**

- Add labeling claims against Pandemic 2009 H1N1 Influenza A Virus per EPA's Guidance Document dated 10/21/09

**General Comments:**

Based on a review of the material submitted, the following comment apply:

The notification is acceptable. A copy has been inserted in your file for future reference.

Should you have any questions or comments concerning this letter, please contact Drusilla Copeland at (703) 308-6224.

Sincerely,



Velma Noble  
Product Manager (31)  
Regulatory Management Branch I  
Antimicrobials Division (7510P)

CONCURRENCES

SYMBOL	SURNAME	DATE						

Print Form

Please read instructions on reverse before completing form.

Form Approved OM 2070-0060



United States  
Environmental Protection Agency  
Washington, DC 20460

☐ Registration  
☐ Amendment  
☒ Other

OPP Identifier Number

## Application for Pesticide - Section I

1. Company/Product Number Microgen, Inc./61178-5	2. EPA Product Manager Velma Noble	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Microgen, Inc./CCX-151	PM# 31	
5. Name and Address of Applicant (Include ZIP Code) Microgen, Inc. 33 Clinton Rd., Suite 102 West Caldwell, NJ 07006 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

## Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

NOTIFICATION OF (ADDING THE EXACT EPA LABELING RECOMMENDATIONS/CLAIMS AS PER SECTION III OF EPA H1N1 GUIDANCE DOCUMENT FOR PANDEMIC 2009 H1N1 INFLUENZA A VIRUS) AS PER EPA PR NOTICE 98-10.  
This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40CFR152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40CFR152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under section 12 and 14 of FIFRA.

THIS ACTION IS NOT SUBJECT TO A FEE UNDER PRIA II

## Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input checked="" type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt No. per container	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 1, 2 qt.; 1, 2, 5, 15, 55 gal.		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

## Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name ROBERT PRINCE		Title PRESIDENT	
		Telephone No. (Include Area Code) 973-575-9025	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 		3. Title President, Microgen, Inc.	
4. Typed Name Robert Prince		5. Date 12-07-2009	

EPA Form 8570-1 (Rev. 8-94) Previous editions are obsolete.

White - EPA File Copy (original)

Yellow - Applicant Copy

(0)



December 7, 2009

**ATTN: DOCUMENT PROCESSING DESK (NOTIF)**

**PRIA II of 2007: Not subject to Registration Service Fee  
Submission is a NOTIFICATION**

Velma Noble  
Product Manager 31  
Regulatory Management Branch I  
Antimicrobial Division (7504P)  
Office of Pesticide Programs  
Environmental Protection Agency  
Ariel Rios Building  
1200 Pennsylvania Ave NW  
Washington, DC 20460

**Regulatory Action:** NOTIFICATION AS PER EPA PR NOTICE 98-10 (adding pandemic 2009 H1N1 influenza A virus statements as per EPA guidance document).  
**Product Name:** CCX-151  
**EPA Registration No.** 61178 - 5

Dear Ms. Noble:

The purpose of this submission is to submit revised labels (one copy as requested) for agency review.

The changes are the addition of the exact labeling recommendations as outlined in the recent EPA Guidance document on labeling claims against **Pandemic 2009 H1N1 Influenza A Virus**. No other changes are being made to the label at this time.

As noted in the guidance document, the Agency will provide its decision electronically to the registrant upon request and subsequently send the registrant a letter confirming the Agency's decision. Please send your electronic decision to the following email: [robert@microgeninc.com](mailto:robert@microgeninc.com)

**In addition, please address and send the final correspondence for this submission directly to the Microgen Inc. address of record in West Caldwell, NJ 07006.**

If you and/or your staff need any clarification or additional information, please phone or email me directly. I will be more than pleased to get the information back to you as soon as possible.

Sincerely,

Robert Prince

**Robert Prince**  
President  
[robert@microgeninc.com](mailto:robert@microgeninc.com)

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA

**VOICE:** 973-575-9025 **FAX:** 973-575-9075 **E-MAIL:** robert@microgeninc.com  
**WEB:** WWW.MICROGENINC.COM

**Microgen, Inc.**

"Redefining Clean"

E.P.A. Reg. No. 61178-5

33 Clinton Road, Suite 102 West Caldwell, NJ 07006 973-575-9025

**CCX-151**

Net Contents:

Batch No:

## CENTER PANEL

### CCX-151

BROAD SPECTRUM DISINFECTANT, HOSPITAL DISINFECTANT, ANTIMICROBIAL, BROAD SPECTRUM CLEANER & DISINFECTANT, GERMICIDAL, GERMICIDE, GERMICIDAL CLEANER, CLEANS AS IT DISINFECTS, ANTIBACTERIAL, BACTERICIDE, PSEUDOMONACIDAL, HIV, & HBV & HCV VIRUCIDE\*, KILLS COMMON HOUSEHOLD GERMS, HOSPITAL VIRUCIDAL\* DISINFECTANT, CONCENTRATED DISINFECTANT CLEANER, CONCENTRATED FORMULA, CLAIMS AGAINST 140+ ORGANISMS

CLEANER, DISINFECTANT, SANITIZER, DETERGENT, FUNGICIDE, DEODORIZER, VIRUCIDE\*, MILDEWSTAT

#### ACTIVE INGREDIENTS:

Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chloride ..... 13.238%  
Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chloride ..... 13.238%

INERT INGREDIENTS..... 73.524%

TOTAL..... 100.00%

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS: 1 quart, 2 quarts, 1 gallon, 2 gallons, 5 gallons, 15 gallons, 55 gallons, 0,528 gallons (2 liters)

Microgen Inc.  
33 Clinton Road  
Suite 102  
West Caldwell, NJ  
07006  
EPA Reg. No. 61178-5  
EPA Est. No. Varies

#### OPTIONAL MARKETING LANGUAGE

- Contains CCX-151 Brand Hospital Disinfectant Formula Exclusively from MICROGEN, INC.
- Makes 180 ready-to-use gallons – typical yield.
- This product is to be diluted only with the (3M™) (Twist 'n Fill™) (Chemical) dispenser.
- Made in USA
- Use on hard, non-porous surfaces such as floors or walls to clean, disinfect and deodorize.
- 3M and Twist 'n Fill (Dose 'n Fill) are trademarks of 3M.

## LEFT PANEL- NUMBER 1

### PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER. Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Harmful if swallowed.

Wear protective eyewear (goggles, face shield or safety glasses). Wear protective clothing and rubber gloves. Avoid contamination of food. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

### STATEMENT OF PRACTICAL TREATMENT — FIRST AID

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional medical advice, call the following emergency phone number: 000-000-0000.**

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

**If container is 5 gallons or more, the following statement must be used:**

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

### PHYSICAL OR CHEMICAL HAZARDS

**Combustible.** Do not use or store near heat or open flame.

## LEFT PANEL- NUMBER 2

### STORAGE and DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

### STORAGE

Do not store on side. Avoid creasing or impacting of side walls. Store securely in closed original container. Avoid storage at temperature extremes or in sunlight. Avoid shipping or storing below freezing. If product freezes, thaw at room temperature and shake gently to remix components. Use locked storage in an area that will prevent cross-contamination of other pesticides, fertilizer, food and feed. Store in locked area inaccessible to children.

### PESTICIDE DISPOSAL


Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER DISPOSAL

**Triple rinse container (or equivalent) promptly after emptying.** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Discard Rinsate. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.**

**BATCH CODE:**



## RIGHT PANEL-NUMBER 1

This product contains no phosphorous.

This product is a phosphate free, pH neutral formulation designed to provide effective cleaning, deodorizing and disinfection in areas where housekeeping is of prime importance in controlling the hazard of cross contamination on treated surfaces.

This product maximizes (improves) labor results by effectively controlling odors.

This product cleans by removing dirt, grime, blood, urine, fecal matter and other common soils found in animal housing facilities, livestock, swine poultry facilities, grooming facilities, farms, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It (also) eliminates odors leaving surfaces smelling clean and fresh.

This product can be used to disinfect, clean and deodorize terrarium and small animal cages, hot rock, substrate and cage furniture (plastic terrarium ornaments, driftwood, heat caves and water dishes. **(This use not allowed in California)**

This product is for use in kitchens, bathrooms, and other household areas.

This product is a multi-surface cleaner, deodorizer and disinfectant. Use on windows, mirrors, and other non-food contact glass surfaces.

When used as directed, this product will deodorize surfaces in restroom and toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and other places where bacterial growth can cause malodors.

Cross-contamination is of major housekeeping concern. This product has been formulated to aid in the reduction of cross-contamination on treated surfaces not only in hospitals, but also in schools, institutions and industry.

This product delivers non-acid disinfection performance in an economical concentrate.

This product is a concentrated Hospital Use disinfectant that is effective against a broad spectrum of bacteria, is virucidal\*, (and) fungicidal, and eliminates odor causing bacteria when used as directed.

This product is an economical concentrate that can be used with a mop and bucket, trigger sprayers, (or) sponge, or by soaking.

This product will not leave a grit or soap scum.

Will not cause swelling of transducer membrane or harm compressor plates.

This product is a versatile Disinfectant for Veterinarian, Veterinary Practice, Animal Care, Animal Laboratory, and Agricultural and Farm Premise applications.

This product is a complete, chemically balanced disinfectant/sanitizer that provides clear use solutions even in the presence of hard water.

This product inhibits bacterial growth on moist surfaces and deodorizes by killing microorganisms that cause offensive odors. **(Not for use in California)**

This product is for use as a disinfectant on hard, non-porous surfaces (at 703 ppm active).

This product is for use in household and commercial humidifiers. Use of this product will control unpleasant (malodors) odors on commercial humidifiers. **(Not for use in California)**

This product is a versatile sanitizer and broad-spectrum disinfectant formulated for use in Ultrasonic Baths (Ultrasonic cleaning units).

This product is a versatile cleaner, broad-spectrum disinfectant and sanitizer formulated for use on bath and therapy equipment (Whirlpools).

## RIGHT PANEL-NUMBER 2

Use this product to sanitize and disinfect non-porous salon/barber tools and instruments such as combs, brushes, scissors, clippers, trimmers, razors, blades, tweezers and manicure instruments.

This product is a one-step neutral disinfectant that is effective against a broad spectrum of bacteria, is virucidal\* (including HIV-1, HCV & HBV) and inhibits the growth of mold and mildew and their odors when used as directed.

This product may be applied through low-pressure sprayers, and fogging (wet misting) systems.

This product neutralizes musty odors and tough odors from smoke, pet accidents, and spills on contact.

This product is specially formulated to effectively eliminate offensive odors caused by mold and mildew.

This product controls (reduces) (eliminates) (neutralizes) (destroys) odors to make your home (kitchen) (bathroom) sanitary.

This product is effective against household germs and odors by animal waste and bathroom and kitchen odors.

Use this product on the multi-touch surfaces responsible for cross-contamination.

This product is effective at controlling mold and mildew on shower curtains.

This product may be used in work areas such as tool rooms and garages for odor control and light duty cleaning.

This product provides long lasting freshness against tough (pet) odors such as odors from litter boxes and pet accidents.

This product provides effective cleaning strength that will not dull most metal-interlock floor finishes, and does not require a rinse prior to recoat.

This product cleans, shines, deodorizes and disinfects all hard non-porous surfaces listed on the label. It inhibits the growth of mold and mildew, leaving bathrooms and kitchens clean and fresh smelling.

This product is an effective one step sanitizer/cleaner for use on non-food contact surfaces.

When used as directed, this product will deodorize surfaces in toilet areas, behind and under sinks and counters, garbage cans and garbage storage area, and other places where bacterial growth can cause malodors.

This product is for use on non-scratch cleaning of showers and tubs, shower doors and curtains, fixtures and toilet bowls.

This product is an effective antimicrobial cleaner designed for use by wholesale and retail florists, shippers and green houses. *(Not for use in California.)*

When used as directed, this product will disinfect hard, non-porous surfaces such as flower buckets, floors and walls of coolers, design and packing benches and counter tops. *(Not for use in California.)*



## RIGHT PANEL-NUMBER 3

This product may be relied on to deodorize coolers, buckets, garbage pails and other areas where obnoxious odors may develop. *(Not for use in California.)*

Use this product to clean, disinfect and deodorize flower buckets, walls and floors of coolers, shippers, greenhouse packing areas, garbage pails and other areas where obnoxious odors may develop. *(Not for use in California.)*

This product may be used as a general purpose antimicrobial detergent in florist shops, wholesale florist, shippers, green house packing areas and other commercial floriculture places for efficient cleaning and antimicrobial action against certain bacteria which cause: *(Not for use in California.)*

Plugging of stems with slime, which reduces uptake of water for various flowers including roses, chrysanthemums, gladioli and tulips.

Production of ethylene gas, which may injure blooms of the various sensitive flowers including carnations, snapdragons, some orchids, baby's breath, sweet peas, freesia and alstroemeria.

Floor cleaner.

One-step cleaner.

Cleans everyday kitchen messes.

Cleans and shines.

Great for use (on) (in) the (kitchen), (bathroom), (floors) and other household areas. Removes (eliminates) odors. Deodorizes.

Will not harm most surfaces. For a cleaner, fresher household. Cleans without bleaching. Removes stains. Removes dirt. Non-staining. Clear formula.

Eliminates odors caused by bacteria (and) (mildew) (and) (non-fresh foods). Kills odor causing bacteria in the kitchen (bathroom).

Cleans everyday kitchen messes like dirt, grease and food stains on non-food contact surfaces. Cuts through tough grease and grime.

*Escherichia coli (E. coli)*, *Salmonella enterica (Salmonella)*, and *Staphylococcus aureus (Staph)* are common bacteria and are found where food is prepared and stored. This product kills these microorganisms and helps prevent the spread of food borne contamination on kitchen surfaces listed on this label.

Disinfects (and Sanitizes) kitchen surfaces (bathroom surfaces and floors). Sanitizes kitchen surfaces (bathroom surfaces and floors). Kills germs. Kills household bacteria.

Antibacterial.

Kills *Tricophyton mentagrophytes*, agent which causes Athlete's Foot fungus, in the bathroom.

Hospital use Disinfectant. Institutional Disinfectant.

Kills common kitchen (bathroom) germs (and viruses).

Effective non-food contact sanitizer in the presence of 5% serum contamination.

Effective non-food contact sanitizer in the presence of soils.

## RIGHT PANEL-NUMBER 4

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

To be used in hospitals in the following areas as a disinfectant: operating rooms, patient care rooms & facilities, recovery, anesthesia, ER, radiology, X-ray cat labs, newborn nurseries, orthopedics, respiratory therapy, surgi-centers, labs, blood collection rooms, central supply, housekeeping & janitorial rooms, nursing homes, doctor's offices & labs, dentists offices & labs (dental operatories).

This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that: (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

CCX-151 is for use in:

- Hospitals, nursing homes, medical and dental offices and clinics, physician offices, operating rooms, isolation wards, and medical research facilities.
- Patient care rooms & facilities, recovery (rooms), anesthesia, Emergency Rooms, X-ray cat labs, newborn nurseries, orthopedics, whirlpool surfaces, footbath surfaces, respiratory therapy, surgi-centers, labs, blood collection rooms, central supply, housekeeping & janitorial rooms.
- EMS & fire facilities, emergency vehicles, ambulance(s), ambulance equipment/surfaces, police cars.
- Day care centers and nurseries, sick rooms.
- Acute care institutions, alternate care institutions, home healthcare institutions.
- Life care retirement communities.
- Restaurants, restaurants and bars, bars, cafeterias, institutional kitchens, fast food operations and food storage areas.
- Supermarkets, convenience stores, retail and wholesale establishments, department stores, shopping malls, gift shops, video stores, bookstores, dressing rooms and laundries, photocopy centers, bicycle shops, auto repair centers.
- Computer manufacturing sites, toy factories.
- Food establishments, coffee shops, donut shops, bagel stores, pizza parlors, liquor stores.
- Crime scenes and funeral homes, mortuaries, burial vaults, mausoleums, autopsy rooms.
- Police stations, courthouses, correctional facilities, jails, prisons, municipal government buildings, penitentiaries, correctional institutions, bus stations, train stations.
- Institutional facilities, laboratories, factories, business and office buildings, restrooms, hotels and motels, and transportation terminals.
- Public restrooms, public facilities, waysides, travel rest areas, shower rooms, shower stalls, bathrooms.
- Hotel, motels, dormitories.
- Kitchens, bathrooms and other household areas.
- Homes (households).

## RIGHT PANEL-NUMBER 5

- Institutions, schools and colleges, churches, classrooms, community colleges, universities, athletic facilities and locker rooms, exercise rooms, exercise facilities, gyms, gymnasiums.
- Cosmetic manufacturing facilities, medical device manufacturing facilities, biotechnology firms, pharmaceutical manufacturing facilities.
- Heath clubs, spas, tanning spas, tanning beds, footbath surfaces, massage/facial salons, hair/nail/pedicure salons, barber/beauty shops, salons.
- Museums, art galleries, post offices, performance/theater centers, banks, libraries, movie houses, bowling alleys.
- Recycling centers.
- Humidifier water tanks. *(Not for use in California)*
- Campgrounds, playgrounds, recreational facilities, picnic facilities, sports arenas, sports complexes.
- Food processing plants, USDA inspected food-processing facilities, dairy farms, hog farms, equine farms, poultry and turkey farms and egg processing plants, meat/poultry processing plants, meat/poultry producing establishments, mushroom farms, rendering plants.
- Processing facilities for Fish, Wine, Milk, Citrus, Fruits, Vegetable, Ice Cream, and Potatoes, and beverage plants.
- Tobacco plant premise.
- Veterinary clinics, animal life science laboratories, kennels, dog/cat animal kennels, breeding and grooming establishments, pet animal quarters, zoos, pet shops, tack shops and other animal care facilities.
- Household and automotive garages, boats, ships, barges, campers, trailers, mobile homes, cars, trucks, buses, trains, taxis and airplanes.
- Cruise lines, airline terminals, airports, shipping terminals, public transportation.
- Commercial florist and flower shops.
- Basements, cellars, bedrooms, attics, living rooms and porches.
- This product may be used on washable hard non-porous surfaces such as:
- Countertops (Counters), non-food contact surfaces of Slurppy<sup>®</sup> machines, (stoves) stovetops, sinks (bathroom, kitchen), tub surfaces, and exterior surfaces of appliances, refrigerators and ice machines.
- Glass, metal, stainless steel, glazed porcelain, glazed ceramic, granite, marble, plastic (such as polystyrene or polypropylene), sealed limestone, sealed slate, sealed stone, sealed terra cotta, sealed terrazzo, chrome and vinyl.
- Enameled surfaces, painted (finished) woodwork, Formica<sup>®</sup>, vinyl and plastic upholstery.
- Examination tables, X ray tables, washing areas, animal grooming areas.
- Tables, chairs, desks, bed frames, lifts, washable walls, cabinets, doorknobs and garbage cans, cuspidors and spittoons.
- Exhaust fans, refrigerated storage and display equipment, coils and drain pans of air conditioning and refrigeration equipment and heat pumps.
- Large inflatable, non-porous, plastic and rubber structures such as animals, promotional items, moonwalks, slides, obstacle course play and exercise equipment.

## RIGHT PANEL-NUMBER 6


- Hard, non-porous surfaces of picnic tables and outdoor furniture or Non-wooden Picnic tables and outdoor furniture except cushions and wood frames. (Note: Only one statement will be used on label.)
- Telephones and telephone booths.
- Highchairs, baby cribs, diaper changing stations, infant bassinets/cribs/warmers/incubators/care equipment, folding tables.
- Bed railings, bedpans, cervical collars, CPR training mannequins, curing lights, neck braces, oxygen hoods, slit lamps, spine backboards, stretchers and unit stools.
- External lenses vision correction (not for use on contact lenses), light lens covers, optical instruments/implements.
- Drinking fountains.
- Foundations, steps, plumbing fixtures, finished baseboards and windowsills.
- Shower stalls, shower doors and curtains, bathtubs and glazed tiles, chrome plated intakes, toilets, toilet bowls, toilet bowl surfaces, urinals, empty diaper pails, portable and chemical toilets and latrine buckets, porcelain tile and restroom fixtures.
- Ultrasonic baths, whirlpools, whirlpool bathtubs.
- Kennels, kennel runs, cages, kennel/cage floors, conductive flooring.
- Wrestling and gymnastic mats, athletic training tables, physical therapy tables.
- Use this product to clean non-porous personal protective safety equipment, protective headgear, athletic helmets, wrestling/boxing headgear, athletic shoe soles, hard hats, half mask respirators, full face breathing apparatus, gas masks, goggles, spectacles, face shields, hearing protectors and ear muffs. Rinse all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Precaution: Cleaning at 120°F temperature will avoid overheating and distortion of the personal safety equipment that would necessitate replacement.)
- Use this product to clean, sanitize and disinfectant non-porous ambulance equipment and surfaces by rinsing all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Precaution: Cleaning at 120°F temperature will avoid overheating and distortion of the ambulance equipment and surfaces that would necessitate replacement.)
- This product is for Poultry Premise Sanitation (Hatcheries)
 

Egg Receiving Area	Tray Dumping Area	Chick Processing Area
Egg Holding Area	Chick Holding Area	Chick Loading Area
Setter Room	Hatchery Room	Poultry Buildings
- This product is for Swine Premise Sanitation:
 

Farrowing Barns and Areas	Dressing Plants	Blocks
Waterers and Feeders	Loading Equipment	Creep Area
Hauling Equipment	Nursery	Chutes
- This product meets AOAC Use – Dilution Test Standards for hospital disinfectants at 300 ppm water hardness.
- This product meets AOAC efficacy standards for hard surface, non-food contact sanitizers.
- This product has passed the Virucidal\* Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Hepatitis B Virus.

## RIGHT PANEL-NUMBER 7

- This product has passed the Virucidal\* Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Bovine Viral Diarrhea Virus BVDV (surrogate for human Hepatitis C virus).
  - OR
- This product has passed the Virucidal\* Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Hepatitis C Virus (HCV). (*Not for use in California*)
  - OR
- At 0.35 ounce per gallon (or equivalent use dilution of 1:365) in the presence of 5% organic load this product was proven to be effective against (*Duck*) Hepatitis B Virus (HBV) and Bovine Viral Diarrhea Virus BVDV (surrogate for human Hepatitis C virus) with a contact time of 10 minutes.
- This product has passed the Virucidal\* Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces Utilizing Human and *Canine Coronavirus*.
- This product is effective against the control of small flies on floors, walls, drains, countertops, metal surfaces, painted surfaces, glazed porcelain, glazed tile, glass, chrome, rubber, and plastic in restaurants, kitchens, dishwashing areas, and bar and wait station areas.
- Efficacy tests have demonstrated that this product is an effective bactericide, virucide\* and fungicide in the presence of organic soil (5% blood serum).
- For larger areas such as operating rooms and patient care facilities, this product is designed to provide both general cleaning and disinfection.



## RIGHT PANEL-NUMBER 8

### BACTERICIDAL STABILITY OF USE-DILUTION:

Tests confirm that this product, when diluted in 400 ppm hard water and in the presence of 5% soil load, remains effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella enterica* for up to one year when stored in a sealed container at room temperature.

If the use-dilution product becomes visibly dirty or contaminated, the use-dilution must be discarded and a fresh product prepared. Always use clean, properly labeled dry containers when diluting the product. Bactericidal stability of the use-dilution does not apply to open containers such as buckets or pails. Use-dilution product in open containers must be prepared daily or more often if the solution becomes visibly dirty or diluted or contaminated.

### DISINFECTION:

#### PREPARATION OF USE SOLUTION:

For water hardness up to 300 ppm add 0.35 ounces per gallon (2.8 milliliters per liter) of water (1:365) to disinfect hard, non porous surfaces [Glass, metal, stainless steel, glazed porcelain, glazed ceramic, granite, marble, plastic (such as polystyrene or polypropylene), sealed limestone, sealed slate, sealed stone, sealed terra cotta, sealed terrazzo, chrome and vinyl]. Apply solution with a cloth, mop, sponge, hand pump trigger sprayer or other mechanical sprayer devices. Treated surfaces must remain wet for 10 minutes. Let air dry. Prepare a fresh solution for each use. CCX-151 is effective in hard water up to 300 ppm hardness.

This product, in the presence of a 98 % organic soil load, diluted 1:365 (0.35 ounces per gallon) in 791 ppm Hard Water, demonstrated efficacy within 10 minutes against the following organisms: *Staphylococcus aureus*, *Salmonella enterica*.

This product is a Hospital Use Disinfectant at 0.35 ounces per gallon (2.8 milliliters per liter), (1:365 dilution), modified in the presence of 300 ppm hard water (calculated as  $\text{CaCO}_3$ ) and in the presence of organic soil (5% blood serum) for a contact time of 10 minutes.

Remove gross filth or heavy soil. For heavily soiled areas, a pre-cleaning step is required.

This product is Bactericidal according to the AOAC Use Dilution Test Method, Virucidal\* according to the virucidal qualification on hard, inanimate surfaces, modified in the presence of 5% organic serum against the microorganisms listed as follows (insert microorganism list here):

Note that the organisms referenced in the above statement are not associated with blood spills. For blood spills, the surface must be thoroughly cleaned before applying the disinfectant.

## RIGHT PANEL-NUMBER 9

### Disinfection/Fungicidal/\*Virucidal\* Directions:

Apply use solution to hard inanimate, non-porous surfaces thoroughly wetting surfaces with a cloth, mop, sponge or sprayer. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications use a coarse spray device. Spray 6-8 inches from surface and rub with brush, sponge or cloth. Do not breathe spray.

Add 0.35 ounces per gallon (2.8 milliliters per liter) of water to disinfect hard, non-porous surfaces. Treated surfaces must remain wet for 10 minutes. Prepare a fresh solution at least daily or when use dilution becomes diluted or soiled.

### Hospitals, Dental Offices, Nursing Homes and other Health Care Institutions

#### Disinfection/Fungicidal and Virucidal\* Dilution Chart

Ounces of Product	Amount of Water
0.09 ounce	1 quart
0.18 ounce	½ Gallon
0.35 ounce	1 gallon
0.9 ounce	2 ½ gallons
1.75 ounces	5 gallons
3.5 ounces	10 gallons

At 0.35 ounces per one gallon (or equivalent use dilution of 1:365) in the presence of 300 ppm hard water ( $\text{CaCO}_3$ ). This product was proven to be effective against Hepatitis B Virus with a contact time of 10 minutes.

OR

At 0.35 ounces per gallon (or equivalent use dilution of 1:365) in the presence of 5% organic load this product was proven to be effective against Hepatitis B Virus (HBV) and Bovine Viral Diarrhea Virus BVDV (surrogate for human Hepatitis C virus) with a contact time of 10 minutes.

At 0.35 ounces per one gallon (or equivalent use dilution of 1:365) in the presence of 300 ppm hard water ( $\text{CaCO}_3$ ) and 5% organic load this product was proven to be effective against Hepatitis C Virus (HCV) with a contact time of 10 minutes. *(Not for use in California)*

### KILLS HIV, HCV & HBV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS

**PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care setting or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS), Hepatitis C Virus (HCV) and Hepatitis B Virus.



## RIGHT PANEL-NUMBER 10

### **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HCV & HBV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

#### **PERSONAL PROTECTION:**

Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks and eye coverings.

**CLEANING PROCEDURE:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to Federal, State and local regulations for infectious waste disposal. **CONTACT TIME:** Leave surfaces wet for 30 seconds for HIV-1 and 10 minutes for HCV and HBV. The contact time for the viruses, fungi and bacteria listed on this label is 10 minutes except for Polio virus Type 1 (Chat strain) which is 30 minutes.

**SANITIZATION - NON-FOOD CONTACT SURFACES:** Remove all gross filth or heavy soil prior to applying sanitizing solution. To sanitize walls, floors and other hard, nonporous surfaces in such areas as schools, institutions, and industries, use a mop, cloth or immerse item in a solution containing 0.35 ounces per gallon (1:365 dilution) so as to wet all surfaces thoroughly for 1 minute. Drain or air dry. Prepare a fresh solution daily or when solution gets visible dirty.

**Cleansing of Body Surfaces and Body Orifices of Human Remains:** To cleanse away skin secretions and accompanying malodor and to insure the removal of all soil and bloodstains, apply 0.35 ounces of this product to a gallon of water (708 ppm active) to the surfaces and body openings, natural or artificial. Bathe the entire body using sponge or washcloth. A soft brush may be employed on surfaces other than the face. Allow a 10 minute contact time for optimal results. Prepare a fresh solution for application of each remains.

**FUNGICIDAL:** At 0.35 ounces per gallon use-level, (or equivalent use dilution) is effective against the pathogenic fungus *Trichophyton mentagrophytes* (athlete's foot fungus - cause of Ringworm) on inanimate surfaces in the presence of 5% organic soil load and 300 ppm water hardness as  $\text{CaCO}_3$  in locker rooms, dressing rooms, shower and bath areas and exercise facilities. Contact time ~ 10 minutes.

This product, in the presence of a ~ 100% organic soil load, diluted 1:365 (0.35 ounces per gallon) in 395 ppm Hard Water, demonstrated efficacy within 10 minutes against the following pathogenic fungus: *Trichophyton mentagrophytes*. Note that the organism referenced in the previous statement is not associated with blood spills. For blood spills, the surface must be thoroughly cleaned before applying this product.

**VIRUCIDAL\*:** When used on inanimate, hard, non-porous, environmental surfaces at 0.35 ounces per gallon of water for a 10 minute contact time (5% organic soil), except for Poliovirus type 1 (Chat strain): which requires a 30 minute contact time (5% organic soil) and HIV-1 which requires only a 30 second contact time.

This product, in the presence of a 98 % organic soil load, diluted 1:365 (0.35 ounces per gallon) in 400 ppm Hard Water, demonstrated efficacy within 10 minutes against the following virus: *Human Coronavirus*. Note that the organism referenced in the above statement is not associated with blood spills. For blood spills, the surface must be thoroughly cleaned before applying this product.



## RIGHT PANEL-NUMBER 11

**General Deodorization:** To deodorize, add 0.35 ounces of this product per gallon of water (or equivalent use dilution of 1:365). Excess material must be wiped up or allowed to air dry.

**For Use on Finished Floors:** To limit gloss reduction, use 0.35 ounces of this product per gallon of water. Apply with a damp mop or auto scrubber. Allow to air dry.

**For Deodorizing Garbage Cans, Garbage Trucks, Industrial Waste Receptacles and Garbage Handling Equipment:** It is especially important to pre-clean for the product to perform properly. Then, apply a wetting concentration of 1.05 ounces of this product per gallon of water (or equivalent use dilution) for a contact time of 10 minutes.

**For Odors Caused by Dogs, Cats and Other Domestic Animals:** Use on rugs, floors, walls, tile, cages, crates, litter boxes, mats, floor coverings, or any surface soiled by a pet. Test a small inconspicuous area first. Blot problem area. Then follow directions for "General Deodorization".

**Mold and Mildew Control Directions:** Add 0.35 ounces per gallon (2.8 milliliters per liter) of water to control the growth of mold and mildew and their odors on hard, non-porous surfaces. Thoroughly wet all treated surfaces completely. Let air dry. Repeat application weekly or when growth or odor reappear(s).

**To control the growth of mold and mildew on large inflatable, non-porous plastic and rubber structures** (animals, promotional items, moonwalks, slides, obstacle course play and exercise equipment): Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 0.35 ounces per gallon (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

**To control the growth of mold and mildew on non-porous athletic equipment** (wrestling and gymnastic mats, athletic training tables, physical therapy tables, athletic helmets, wrestling/boxing headgear, athletic shoe soles): Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 0.35 ounces per gallon of water (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

**Disinfection/Fungicide/Virucide\* for Barber/Salon Tools Directions:** Immerse pre-cleaned barber/salon tools, such as combs, brushes, razors, clipper and trimmer blades, tweezers, manicure/pedicure tools and scissors, in a 0.35 ounces per gallon solution of the product (or equivalent use dilution). Completely immerse instruments and tools for at least 10 minutes. Rinse thoroughly and dry before use. Prepare fresh solution at least daily or more often if solution becomes cloudy or soiled.

**NOTE:** Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 10 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

**Disinfection of Hard, Non-Porous Surfaces in Footbaths:** To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate surfaces after using the footbath, drain the water and thoroughly clean all hard, non-porous surfaces with soap or detergent, then rinse with water. Saturate surfaces with a use solution of 0.35 ounces per gallon of water (or equivalent use dilution) to exposed surfaces with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow solution to stand for 10 minutes. For spray applications, use a coarse spray device. Do not breathe spray. After the unit has been thoroughly disinfected, rinse all cleaned surfaces with fresh water. Do not use equipment until treatment has been absorbed, set or dried.

## BACK PANEL-NUMBER 1

### DISINFECT TOILET BOWLS

#### NON-ACID TOILET BOWL (AND URINAL) DISINFECTION/CLEANER DIRECTIONS

Remove gross filth prior to disinfection.

**From Concentrate:** Add 0.35 ounces (2.8 milliliters per liter) to the toilet bowl and mix. Brush thoroughly over exposed surfaces and under the rim with a cloth, mop, or sponge. Repeat and allow to stand for 10 minutes and flush.

**From Use Solution:** Empty toilet bowl or urinal and apply 0.35 ounces per gallon (2.8 milliliters per liter) use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer, brush or swab thoroughly and allow to stand for 10 minutes and flush.

**For Heavy Duty Use:** Empty toilet bowl or urinal and apply 1.4 ounces per gallon (11.2 milliliters per liter) use solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer, brush or swab thoroughly and allow to stand for 10 minutes and flush.

**Cleaning/Deodorizing Directions:** Add 0.18 to 0.70 ounces per gallon (1.4 to 5.6 milliliters per liter) of water to clean and deodorize windows, mirrors and non-food contact glass surfaces. When using a coarse spray device, spray 6-8 inches from surface. Rub with sponge or cloth. Do not breathe spray.

**Ultrasonic Bath Disinfectant Directions:** Use this product to disinfect hard (inanimate) non-porous non-critical objects compatible with Ultrasonic cleaning units. Pour fresh solution of 0.35 ounces per gallon of water (or equivalent use dilution) directly into bath chamber. Preclean soiled objects. Place objects into unit and operate for a minimum of 10 minutes, according to manufacturers' use directions. Remove objects and rinse with sterile water (sterile water for injection), or allow to air dry. Replace solution at least daily or when solution becomes visible dirty or discolored.

**For Disinfecting Hard, Non-porous Fiberglass Bath and Therapy Equipment:** To remove body oils, dead tissue, soil and all other buildups or organic matter on inanimate surfaces after using the whirlpool unit, drain the water and refill with fresh water to just cover the intake valve. Add 1.75 ounces of this product for each 5 gallons of water (0.35 ounces per one gallon) in the unit at this point. Briefly start the pump to circulate the solutions. Turn off pump. Wash down the unit sides, seat of the chair lift, and any/all related equipment with a clean swab, brush or sponge. Product to surface contact time must be at least 10 minutes for proper disinfection. After the unit has been thoroughly disinfected, drain solutions from the unit and rinse any/all cleaned surfaces with fresh water. The unit is ready for reuse.

**Disinfection of Hard Non-Porous Surfaces in Whirlpool Units:** After using the whirlpool unit, drain and refill with fresh water to just cover the intake valve. Add 0.35 ounces of this product for each gallon of water at this point. Briefly start the pump to circulate the solution. Turn off the pump. Wash down the unit sides, seat of the chair, lift and any/all related equipment with a clean swab, brush or sponge. Treated surfaces must remain wet for 10 minutes. After the unit has been thoroughly disinfected, drain the solution from the unit and rinse any/all cleaned surfaces with fresh water. Repeat for heavy soiled units.

## BACK PANEL-NUMBER 2

### **SANITIZATION - FOOD CONTACT SANITIZER**

To be used in the following areas as a food contact sanitizer: Cruise Lines, Airlines Terminals, Log Cabins, Coffee Shops, Department Stores, Retail Stores, Bagel Stores, Court Houses, Municipal Government Buildings, Colleges, Universities, Community Colleges, Campgrounds, Travel Rest Areas, Recreational Facilities, Sports Arenas, Donut Shops, Boats/Ships, Hotels/Motels, Dormitories, Movie Houses, Pizza Parlors, Lifecare Retirement Communities, Hospices, Book Stores, Convenient Stores, Liquor Stores, Day Care Centers, Acute Care Institutions, Alternate Care Institutions, Home Health Care Institutions, Food Establishments, Shopping Malls, Churches, Picnic Facilities, Homes, Schools, Institutions, Industries, Meat/Poultry/Food Processing Plants, Dairies, Bars, Restaurants and Cafeterias to sanitize hard, nonporous food contact surfaces as listed and identified below. Use a mop, sponge or cloth to apply sanitizing solution or immerse item in sanitizing solution. This product is an effective sanitizer when diluted in tap water with a contact time of 1 minute.

### **SANITIZATION - FOOD CONTACT SURFACES: (glassware, utensils, cookware and dishware, *Escherichia coli* and *Staphylococcus aureus*)**

1. Remove gross food particles and soil from utensils, glassware, cookware and dishware by a preflush, or pre-scrape and when necessary, pre-soak.
2. Wash with a good detergent.
3. Rinse with potable water.
4. Sanitize in a solution of 0.70 oz. of product per 7 gallons of water (200 ppm active solution). Immerse all utensils for 1 minute or longer if specified by government sanitary code.
5. Place sanitized utensils on a rack or drain board to air dry. Do not rinse or wipe.

### **SANITIZATION - FOOD CONTACT IMMOBILE SURFACES: (food processing equipment, counter tops, tables, picnic tables, appliances, stovetops and food work areas)**

1. Remove all gross food particles and soil by pre-flush or pre-scrape.
2. Clean all surfaces thoroughly using a good detergent or cleaner. Tilt movable surfaces for proper drainage. Rinse with potable water.
3. Mix 0.70 oz. of product per 7 gallons of water (200 ppm active solution).
4. Wet surfaces thoroughly for 1 minute.
5. Let surfaces drain and air dry. Do not rinse or wipe.
6. Fresh sanitizing solution must be prepared daily or more often if solution becomes diluted or soiled.

For mechanical operations: The prepared use solution may be used once for sanitizing and re-used for other purposes such as cleaning.

**To Disinfect Food-Processing and Tobacco Premises:** Before using this product, food products and packaging materials must be removed from area or carefully protected. For floors, walls and storage areas, add 0.35 ounces of this product per gallon of water (or equivalent use dilution). For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. For sprayer applications use a coarse spray device. Allow to remain wet for 10 minutes, then remove excess liquid. After use, all surfaces in the area must be thoroughly rinsed with potable water.

## BACK PANEL-NUMBER 3

**To Disinfect Food Service Establishment Food Contact Surfaces:** Before using this product, food products and packaging materials must be removed from area or carefully protected. For countertops, exterior surfaces of appliances, (and) tables, add 0.35 ounces of this product per gallon of water (or equivalent use dilution). For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. For sprayer applications use a coarse spray device. Allow to remain wet for 10 minutes. Then remove excess liquid and rinse the surface with potable water.

**Disinfecting Potato storage area and equipment:** Remove all potatoes prior to disinfection of potato storage area and equipment. Pre-clean hard surfaces by removing heavy soils or gross filth. Follow general disinfecting directions as outlined in that section. All treated surfaces must be thoroughly rinsed with potable water prior to reuse.

**For Use on Non-Food Contact Surfaces as a General Disinfectant in the Brewery Industry:** Use 0.35 ounces of this product per gallon of water (or equivalent use dilution). Follow the general disinfectant directions above.

**Directions for Fogging:** For use in dairies, beverage and food processing plants. Prior to fogging, food products and packaging material must be removed from the room or carefully protected. After cleaning, fog desired areas using 5.6 ounces per 1000 cubic feet of room area with a product solution containing 0.53 ounces product to 1 gallon of water (or equivalent use dilution) (1,062 ppm). Vacate the area of all personnel for a minimum of 2 hours after fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. All food contact surfaces must be sanitized with an EPA approved food contact sanitizer prior to use. Allow food contact surfaces to drain thoroughly before operations are resumed. Wear a dust mist respirator when mixing the use solution and pouring it into the fogging apparatus.

**NOTE:** The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances must a room or building be entered by anyone within two hours of the actual fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

**FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.**

## BACK PANEL-NUMBER 4

### POULTRY (AND SWINE) PREMISE SANITATION SITE PREPARATION

The first step in any on-going sanitation program should be the removal of gross contamination and debris. This may be accomplished using a shovel, broom, or vacuum depending on the area to be disinfected. The efficacy of even the most efficient germicidal cleaner is reduced in the presence of heavy organic matter. Once the heavy debris is eliminated thoroughly clean all surfaces with soap or detergent and rinse with water.

**Disinfection/Fungicide/Virucide\* of Poultry/Turkey Equipment, Swine Quarters, Animal Quarters and Kennels Directions:** Prior to use, remove all poultry, other animals and their feed from premises, vehicles (trucks) and enclosures such as coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other surfaces of facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 0.35 ounces of this product per gallon of water (or equivalent use dilution). Saturate surfaces with the disinfecting solution for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.

After application, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water (racks, troughs, automatic feeders, fountains and waterers) must be thoroughly scrubbed with soap or detergent then rinsed with potable water before reuse.

**HATCHERIES:** For disinfection/fungicide/virucide, use 0.35 ounces of this product per gallon of water. Apply to hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces exposed to solution for 10 minutes or more and allow to air dry.

**VEHICLES:** Clean all vehicles including mats, crates, cabs, and wheels with high-pressure water and this product. For sprayer applications, use a coarse spray. Use 0.35 ounces of this product per gallon of water for disinfection/fungicide/virucide. Leave all treated surfaces exposed to solution for 10 minutes or more and allow to air dry.

### DRESSING PLANT AND RENDERING PLANT DISINFECTION – FUNGICIDE – VIRUCIDE

**USE DIRECTIONS:** Cover or remove all food and packaging materials. Remove all gross soils. Disinfect walls and floors in poultry and animal dressing plants. Disinfect offal rooms, exterior walls and loading platforms of dressing plants. Saturate all surfaces with the use solution, 0.35 ounces of this product per gallon of water. Scrub to loosen all soils. Allow to soak for 10 minutes and thoroughly rinse all wetted and cleaned surfaces with potable water.

## BACK PANEL-NUMBER 5

### FARM PREMISE DISINFECTION DIRECTIONS

#### For use in Equine, Dairy and Hog Farms:

1. Remove all animals and feed from premises, vehicles and enclosures.
2. Remove all litter and manure from floors, walls and surfaces of barn, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces with the disinfecting solution at 0.35 ounces per gallon of water for a period of 10 minutes.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, scrappers used in removing litter and manure.
7. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set and dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

#### For Treatment of Animal Housing Facilities:

1. Remove all animals and feed from premises, vehicles and enclosures.
2. Remove all litter and manure from floors, walls and surfaces of barn, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces (floors, walls, cages and other washable hard, non-porous environmental surfaces) with the disinfecting solution at 0.35 ounces per gallon of water (or equivalent use dilution) for a period of 10 minutes. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels, scrappers used in removing litter and manure.
7. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set and dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

**Terrarium and Small Animal Cage and Cage Furniture Disinfection (*This use not allowed in California*):** Animals frequently defecate on hot rocks and other cage furniture items inside your terrarium. This can result in high bacteria and ammonia levels that can lead to possible infection/disease in your animals. When used regularly this product can eliminate these high bacteria/ammonia levels in your cage and on your cage furniture items.

1. Remove all animals.
2. Thoroughly clean all surfaces and objects (hot rocks, caves, cage furniture, feeding and watering dishes, and appliances) including the substrate in the terrarium or cage with soap or detergent and rinse with water.
3. Saturate all surfaces (floors, walls, cages and other washable hard, non-porous environmental surfaces) with the disinfecting solution at 0.35 ounces per gallon of water (or equivalent use dilution) for a period of 10 minutes. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. Then wipe surfaces dry.
4. Saturate gravel as above and let stand for 10 minutes. Place in bucket of clean water and swirl for 15-30 seconds. Thoroughly air dry before returning to terrarium.
5. Do not return animals to the habitat until it is dry and ventilated.
6. Thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.
7. Clean terrarium at least once weekly or more as needed.
8. Do not apply this product directly on to the small animal. If this product comes into contact with the small animal's skin, then immediately wash the material off of the animal with lukewarm water. If the small animal ingests the product, contact your veterinarian immediately.

**Note:** Substrates for desert terrariums (i.e. gravel) must be completely dry before returning to terrarium to avoid high humidity levels. Always replace substrate if a foul odor persists.

## **BACK PANEL-NUMBER 6**

**Veterinary Clinics/Animal Life Science Laboratory/Zoos/Pet Shop/Kennels/Breeding and grooming Establishment/Tack Shops Disinfection Directions:** For cleaning and disinfecting the following hard non-porous surfaces: equipment, utensils, instruments, cages, kennels, stables, stalls and catteries. Remove all animals and feeds from premises, animal transportation vehicles, crates etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or transversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 0.35 ounces per gallon of water (or equivalent dilution) for a period of 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, scrappers, used in removing litter and manure. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, automatic feeders, waterers and other equipment which dispenses food or water with soap or detergent, and rinse with potable water before reuse.

### **SANITIZING HATCHERY ROOMS USING FOGGING DEVICES**

Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Close room off so fog is confined to room to be treated. Mix 1 quart of this product to 2½ gallons water. Insert the nozzle of the fogger through a suitable opening into the room. With the setting in maximum output, fog for one minute for each 4000 cubic feet of space in the room. When fogging is completed ventilate buildings and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly clean all surfaces with soap or detergent and rinse with water. Wear a dust mist respirator when mixing the use solution and pouring it into the fogging apparatus.

### **SANITIZING INCUBATORS AND HATCHERS USING FOGGING DEVICES**

Mix 4.2 ounces of this product to 122 ounces of water. Fog 3-8 ounces of this into setters and hatchers immediately after transfer. Repeat daily in setters and every 12 hours in hatchers. Discontinue hatcher treatments at least 24 hours prior to pulling the hatch. It is acceptable to fog setters and hatchers with 0.35 ounces per gallon solution of this product on an hourly or every other hour basis. Vacate the area of all personnel for a minimum of 2 hours after fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. If this is done, fog for 30-90 seconds once per hour or once every two hours. When fogging is complete, ventilate buildings, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed set or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse. Wear a dust mist respirator when mixing the use solution and pouring it into the fogging apparatus. Only for treatment of setters and hatchers after poultry/chicks/eggs have been removed. Not for treatment of hatchers which contain chicks/eggs.

**NOTE:** The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances must a room or building be entered by anyone within two hours of the actual fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

**FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.**

## BACK PANEL-NUMBER 7

**COMMERCIAL FLORIST USE DIRECTIONS** (*Not for use in California*): To clean, disinfect and deodorize hard non-porous surfaces in one step, prepare use solution by adding (mixing) 0.35 ounces per gallon of water (1:365 dilution). For heavy-duty use, add (mix) 1.4 ounces per gallon of water (1:91).

Remove all leaves, petals, garbage and refuse. Pre-clean surfaces using pressurized water where possible. Apply use solution to hard (inanimate) non-porous surfaces thoroughly wetting surfaces as recommended and required, with a cloth, mop, brush, sponge or sprayer.

For heavily soiled areas, a preliminary cleaning is required.

For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray.

Treated surfaces must remain wet for 10 minutes. Allow to air dry.

Prepare a fresh solution at least daily or when use solution becomes visibly dirty.

**FOR CONTROL OF SMALL FLIES ON NON-FOOD CONTACT SURFACES:** For control of small flies: *Drosophila* spp. and the *Phoridae* family. To control flies on non-food contact surfaces such as floors, walls, countertops, metal surfaces, painted surfaces, glazed porcelain, glazed tile, glass, chrome, rubber, and plastic in restaurants, kitchens, dishwashing areas, and bar and wait station areas. Remove food and food packaging prior to use. Cover exposed food-handling surfaces. After-removing gross filth, apply a solution of 0.35 ounces of this product per gallon of water (or equivalent use dilution) to surfaces and locations where flies may breed. Spray surfaces thoroughly or apply by pouring, mopping or sponging onto the surface. Repeat application 1-2 times per week or as needed. Do not contaminate food and food packaging.

**FOR CONTROL OF SMALL FLIES IN DRAINS:** For control of small flies: *Drosophila* spp. and the *Phoridae* family. Spray or pour solution of 0.35 ounces per gallon of water (or equivalent use dilution) into drain during time of lowest level of drain use. Add 2.45 ounces of solution daily to each drain to maintain fly control. Apply around the edge of the drain and coat all sides of inside of drain.

**Waterbed Conditioner:** When used as a waterbed conditioner, controls the growth of odor-causing and slime-forming bacteria. Prevents bubbles, preserves plasticizers, conditions vinyl interior, sequesters minerals.

If bed has not been treated properly, drain bed completely. Add 15 gallons of water, mix vigorously, drain bed again. Fill bed with water and follow dosage directions.

**Dosage:** To control growth of odor-causing and slime-forming bacteria, add 1.33 fluid ounces in a free flow waterbed of 90 to 180 gallon capacity. Add 2.7 fluid ounces in a waveless/fiber waterbed mattress of 90 to 180 gallon capacity. Repeat application every 4 to 6 months.



## BACK PANEL-NUMBER 8

### SANITIZING NON-FOOD CONTACT SURFACE DIRECTIONS

**Preparation of sanitizer use solution:** Add 0.35 ounces (of this product) per gallon of water (or equivalent use dilution) to sanitize hard, non-porous environmental surfaces. This product is an effective sanitizer against the following:

*Klebsiella pneumoniae*  
*Staphylococcus aureus*

Apply sanitizer use solution to pre-cleaned, hard, non-porous surfaces with a cloth, mop, sponge, sprayer or by immersion.

For sprayer applications, use a coarse pump or trigger sprayer. Spray 6-8 inches from surface and rub with brush, sponge or cloth. Do not breathe spray.

Treated surfaces must remain wet for 1 minute, then wipe with sponge, mop or cloth or allow to air dry.

OR

**Sanitizing Non-Food Contact Surface Directions:** Add 0.35 ounces per gallon (2.8 milliliters per liter) of water to sanitize hard non-porous environmental surfaces. Treated surfaces must remain wet for 1 minute. Allow to air dry.

This product is not for use to disinfect, sanitize or clean eating utensils, glassware and dishes.

**To Sanitize (and Deodorize):** Apply use solution to hard inanimate, non-porous surfaces thoroughly wetting surfaces with a cloth, mop, sponge or sprayer. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications use a coarse spray device. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray. Let stand for 1 minute. Then wipe.

**LAUNDRY ADDITIVE (RESIDUAL BACTERIOSTATIC AND RESIDUAL SELF SANITIZING ACTIVITY UNDER CONDITIONS OF HIGH RELATIVE HUMIDITY OR WET CONTAMINATION) AGAINST ODOR-CAUSING BACTERIA FOR INSTITUTIONAL, INDUSTRIAL AND HOSPITAL USE.** This product sanitizes laundry such as bedspreads, sheets, pillowcases, diapers, towels, and other wet linens by controlling and/or reducing the growth of odor-causing bacteria. It can be used in industrial and institutional areas such as motels, hotel chains, nursing homes and hospitals. This product is used as an addition to the final rinse cycle.

Add 1.4 fluid ounces of this product per 100 lbs. of dry laundry to the final rinse cycle water (200 ppm). If the product is to be diluted prior to adding it to the final rinse cycle, use 0.18 ounce per gallon of water and then add to the washwheel in the final rinse cycle.

**SHOE BATH SANITIZER:** To prevent cross contamination from area to area in animal areas, and the packaging and storage areas of food plants, shoe baths containing one inch of freshly made solution must be placed at all entrances to buildings, hatcheries and at all the entrances to the production and packaging rooms. Scrape waterproof shoes and place in 0.35 ounces of this product per gallon (2.8 milliliters per liter) of water solution for 1 minute prior to entering area. Change the sanitizer solution in the bath at least daily or sooner if solution appears dirty.

**SHOE FOAM DIRECTIONS** (*Not for use in California*): To prevent cross contamination from area to area in animal areas, and the packaging and storage areas of food plants, apply a foam layer approximately 0.5 to 2 inches thick made from a solution of 0.35 to 0.5 ounces per gallon of water (or equivalent use dilution) (708 to 1029 ppm active) at all entrances to buildings, hatcheries, production and packaging rooms by using a foam generating machine or aerator to apply foam layer. Follow the foaming directions as specified by the manufacture of the foam generator/aerator. Scrape waterproof shoes. Stand and/or walk through foamed area for 1 minute prior to entering area. Foam area must be washed and replaced at least daily or when it appears dirty.

## BACK PANEL-NUMBER 9

*(For food processing or other facilities that have installed entryway sanitizing systems)*

**ENTRYWAY SANITIZING SYSTEMS:** To prevent cross contamination from area to area, set the system to deliver (0.35 oz. per gallon of water) of sanitizing solution. The (spray) (foam) must cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor.

Do not mix other foam additives to the sanitizing solution.

**SALON/BARBER INSTRUMENTS AND TOOLS SANITIZER DIRECTIONS:** Pre-clean soiled instruments and tools. Completely immerse water safe non-porous instruments and tools in a solution of 0.35 ounces per gallon (or equivalent use dilution) of water for at least 1 minute. Rinse thoroughly and dry before use. Fresh solution must be prepared at least daily or more often if solution becomes cloudy, or soiled.

**NOTE:** Plastics may remain immersed until ready to use. Stainless steel shears and metal instruments must be removed after 10 minutes, rinsed, dried, and kept in a clean non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

**Ultrasonic Bath Disinfection Directions:** Use this product to disinfect hard, (inanimate) non-porous, non-critical objects compatible with Ultrasonic cleaning units. Pour a fresh solution of 0.35 ounces per gallon of water (or equivalent use dilution) directly into bath chamber. Pre-clean soiled objects. Place objects into unit and operate for a minimum of 10 minutes, according to manufacturers' use directions. Remove objects and rinse with sterile water (sterile water for injection), or allow to air dry. Replace solution at least daily or when solution becomes visible dirty or discolored.

**Disinfection of Hard, Non-Porous Surfaces in Whirlpool Units:** After using the whirlpool unit, drain and refill with fresh water to just cover the intake valve. Add 0.35 ounces of this product for each gallon of (or equivalent use dilution) water at this point. Briefly start the pump to circulate the solution. Turn off the pump. Wash down the unit sides, seat of the chair, lift and any/all related equipment with a clean swab, brush or sponge. Treated surfaces must remain wet for 10 minutes. After the unit has been thoroughly disinfected, drain the solution from the unit and rinse any/all cleaned surfaces with fresh water. Repeat for heavy soiled units.

**Humidifier Directions** *(Not applicable in CA):* Thoroughly clean water tank and filters. Add 0.07 ounces of this product per gallon of refill water. Not for use in heat or atomizing type humidifiers.

OR

**Humidifier Bacteria/Algae Treatment** *(Not for use in California):* Formulated for use in portable humidifiers. Thoroughly clean water tank and filters before each heating season, or sooner if necessary. For every gallon of water in the humidifier tank add 0.07 ounces of this product. When you refill the tank, add 0.07 ounces of this product for every gallon of water added.

Not for use in heat vaporizing or atomizing type humidifiers.

## BACK PANEL-NUMBER 10

**Special Instructions for Cleaning Carpet Against Odor Causing Bacteria:** This product may be used in industrial, institutional, commercial and residential areas such as homes, motels & hotel chains, nursing homes, schools and hospital. For use on wet, cleanable synthetic fibers. Do not use on wool. Vacuum carpet thoroughly prior to cleaning. Test fabric for color fastness.

**For portable extraction units:** Mix 0.18 ounce of this product per gallon of water.

**For truck mounted extraction machines:** Mix 4.2 ounces of the product per gallon of water and meter at 4 gallons per hour.

**For rotary floor machines:** Mix 0.35 ounces of this product per gallon of water and apply at the rate of 300-500 sq. ft. per gallon.

Do not mix this product with other cleaning products. Follow the cleaning procedures specified by the manufacturer of the cleaning equipment. After using this product, set the carpet pile and protect the carpet from furniture legs and bases while drying. Do not over wet. If applied to stain resistant nylon carpet, apply a fabric protector according to the carpet manufacturer's directions.

## BACK PANEL-NUMBER 11

### CARPET DEODORIZER AGAINST ODOR-CAUSING BACTERIA, FOR HOME, INSTITUTIONAL, INDUSTRIAL AND HOSPITAL USE

This product cleans and deodorizes the carpet (by controlling/reducing the growth of odor-causing bacteria). It can be used in industrial, institutional, commercial and residential areas such as homes, motels, hotels chains, nursing homes and hospitals. *(Note to reviewer: statement in parenthesis not allowed for use in California.)*

Vacuum carpet thoroughly prior to application. Mix 0.18 ounce of product per gallon of water. Follow the injection and/or extraction procedures as specified for any conventional steam cleaning equipment you are using. For rotary floor machines, mix 0.35 ounces per gallon of water and spray on carpet at a rate of 300-500 sq. ft. per gallon.

For use on washable synthetic fibers. Do not use on wool. Test color fastness of carpet before use. Apply diluted product to a small concealed spot, then rub with a clean white cloth. If color changes or transfers to cloth, a water-based product should not be used.

After using the product, set carpet pile in one direction with a stiff brush. Place aluminum foil under the legs of furniture while carpet is drying. Over-wetting can cause carpet to shrink. Manufacturer assumes no responsibility for over-wetting misuse.

**Note:** This product should not be mixed with other cleaning products.

### FOOD PROCESSING PLANTS USING FOGGING DEVICES

For use in dairies, beverage and food processing plants. Prior to fogging, food products and packaging material must be removed from the room or carefully protected. Wear a dust mist respirator when mixing the use solution and pouring it into the fogging apparatus. After cleaning, fog desired areas using 5.6 ounces per 1000 cubic feet of room area with a solution containing 0.5 ounce of product to 1 gallon of water (1,062 ppm). Vacate the area of all personnel for a minimum of 2 hours after fogging. All food contact surfaces must be thoroughly rinsed prior to reuse with potable water then sanitized with an EPA approved food contact sanitizer.

**NOTE:** The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances must a room or building be entered by anyone within two hours of the actual fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. If the building must be entered, then the individuals entering the building must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

**FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.**

### MUSHROOM FARM INDUSTRY USE DIRECTIONS

**Site Preparation:** The first step in any on going sanitation program should be the removal of gross contamination and debris. This may be accomplished by using a shovel, broom, or vacuum, depending on the area to be disinfected.

**Disinfection:** Use 0.35 ounces of this product per gallon of water (or equivalent dilution). Wet all surfaces thoroughly. Treated surfaces should be allowed to remain wet for 10 minutes. Let air-dry. For heavily soiled areas, pre-clean first. Prepare a fresh solution for each use.

**For Heavy Duty Cleaning:** When greater cleaning is desired, use 0.70 ounces of this product per gallon of water (or equivalent dilution). Heavily soiled areas may require repeated cleaning before treatment.

**DO NOT APPLY TO THE MUSHROOM CROP, COMPOST OR CASING.** Rinse treated surfaces with potable water before they contact the crop, compost or casing.

## MICROORGANISM LIST INSERT

**Disinfection Performance:** At 0.35 ounces of this product to one gallon of water use level, this product is bactericidal and fungicidal on hard inanimate surfaces modified in the presence of 5% organic serum with a 10 minute contact time against:

### Isolates From AIDS Patients

1. *Aspergillus niger*
2. *Candida albicans*
3. *Cryptococcus neoformans*
4. *Pseudomonas aeruginosa*
5. *Staphylococcus aureus*
6. *Streptococcus pneumoniae*

### Gram Positive Clinical Isolates

7. *Enterococcus faecalis*
8. *Micrococcus luteus*
9. *Staphylococcus aureus*
10. *Staphylococcus aureus* (Toxic shock)
11. *Staphylococcus epidermidis*
12. *Staphylococcus saprophyticus*
13. *Streptococcus haemolyticus*
14. *Streptococcus pyogenes*

### Gram Negative Clinical Isolates

15. *Acinetobacter calcoaceticus* var. *anitratus*
16. *Acinetobacter calcoaceticus* var. *lwoffii*
17. *Bordetella bronchiseptica*
18. *Brevundimonas diminuta*
19. *Burkholderia cepacia*
20. *Enterobacter agglomerans*
21. *Enterobacter cloacae*
22. *Enterobacter gergoviae*
23. *Enterobacter liquefaciens*
24. *Escherichia coli* (Urinary)
25. *Escherichia coli* (Wound)
26. *Flavobacterium meningosepticum*
27. *Hafnia alvei*
28. *Klebsiella oxytoca*
29. *Klebsiella pneumoniae*
30. *Morganella morganii*
31. *Proteus mirabilis*
32. *Proteus vulgaris*
33. *Pseudomonas aeruginosa*
34. *Pseudomonas fluorescens*
35. *Pseudomonas pseudomallei*
36. *Pseudomonas putida*
37. *Pseudomonas stutzeri*
38. *Serratia marcescens*
39. *Sphingomonas paucimobilis*

**Other Bacteria**

40. *Actinobacillus pleuropneumoniae*
41. *Actinomyces pyogenes*
42. *Bacillus cereus*
43. *Bacteroides fragilis*
44. *Corynebacterium ammoniagenes* (*Brevibacterium ammoniagenes*)
45. *Bordetella bronchiseptica*
46. *Burkholderia pickettii*
47. *Campylobacter jejuni*
48. *Chryseomonas luteola*
49. *Corynebacterium pseudotuberculosis*
50. *Enterobacter aerogenes*
51. *Enterococcus faecalis*
52. *Enterococcus faecium*
53. *Enterococcus hirae*
54. *Escherichia coli*
55. *Escherichia coli* strain 0157:H7
56. *Escherichia vulneris*
57. *Haemophilus influenzae*
58. *Klebsiella pneumoniae*
59. *Listeria monocytogenes*
60. *Pasteurella haemolytica*
61. *Pseudomonas aeruginosa*
62. *Rhodococcus equi*
63. *Salmonella enterica*
64. *Salmonella schottmuelleri*
65. *Salmonella typhi*
66. *Shigella dysenteriae*
67. *Staphylococcus aureus*
68. *Staphylococcus auricularis*
69. *Staphylococcus capitis*
70. *Staphylococcus hominis*
71. *Staphylococcus simulans*
72. *Stenotrophomonas maltophilia*
73. *Streptococcus equi* var. *equi*
74. *Streptococcus equi* var. *zooepidermicus*
75. *Streptococcus pneumoniae* (PRSP)
76. *Streptococcus pyogenes*
77. *Streptococcus salivarius*
78. *Yersinia enterocolitica*

**Pathogenic Fungi**

79. *Trichophyton mentagrophytes*

**Environmental Fungi**

80. *Aspergillus candidus*
81. *Aspergillus niger*
82. *Penicillium chermesinum*
83. *Penicillium oxalicum*
84. *Penicillium spinulosum*
85. *Ulocladium* sp.

**Antibiotic Resistant Gram Negative Bacteria**

86. *Pseudomonas aeruginosa* (Sulfa, Cefatoxime, Nitrofurantoin, Tetracycline, Amikacin, Ampicillin, Cephalothin and Bactine Resistant)
87. *Escherichia coli* (Ampicillin, Tetracycline, Penicillin and Sulfa Resistant)
88. *Klebsiella oxytoca* (Ampicillin, Sulfanilimide and Tetracycline Resistant)
89. *Klebsiella pneumoniae* type 1 (Ampicillin, Tetracycline, Cephalothin and Sulfa Resistant)
90. *Morganella morganii* (Penicillin and Tetracycline Resistant)
91. *Enterobacter agglomerans* (Ampicillin and Sulfanylimide Resistant)
92. *Salmonella choleraesuis* (Antibiotic Resistant)
93. *Enterobacteriaceae* with extended beta-lactamase resistance (Ampicillin and Piperacillin Resistant)

**Antibiotic Resistant Gram Positive Bacteria**

94. *Enterococcus faecalis* (Vancomycin Resistant-VRE)
95. *Enterococcus faecium* (Vancomycin Resistant-VRE)
96. *Staphylococcus aureus* (Methicillin-MRSA, Community Associated Methicillin Resistant - CA-MRSA PVL Positive)
97. *Staphylococcus aureus* (CA-MRSA Genotype USA 400)
98. *Staphylococcus aureus* (Penicillin G, Penicillin, Ampicillin, Cefazolin, Cefatoxime, Chloramphenicol, Ciprofloxacin, Clindimycin, Erythromycin, Oxacillin, Rifampin, Tetracycline Resistant)
99. *Staphylococcus aureus* (Vancomycin Resistant – VRSA)
100. *Staphylococcus aureus* (Vancomycin Resistant Intermediate-VISA)
101. *Staphylococcus epidermidis* (Ampicillin and Drug Resistant)

**Virucidal\* Performance:** At 0.35 ounces per gallon (1:365) use level, this product was evaluated in the presence of 5% serum with a 10 minute contact time unless otherwise noted below and found to be effective against the following viruses on hard, non-porous environmental surfaces:

*Respiratory illnesses attributable to Pandemic 2009 H1N1 are caused by influenza A virus. This product (Product Name) is a broad-spectrum hard surface disinfectant that has been shown to be effective against (influenza A virus tested and listed on the label) and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).*

*This product has demonstrated effectiveness against influenza A virus and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 influenza A virus.*

*This product has demonstrated effectiveness against (influenza A virus tested and listed on the label) and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).*

*Kills Pandemic 2009 H1N1 influenza A virus (formerly called swine flu).*

*Kills Pandemic 2009 H1N1 influenza A virus.*

#### **Human Viruses**

102. Adenovirus type 2
103. Cytomegalovirus
104. HBV (Hepatitis B Virus)
105. HCV (Hepatitis C Virus)
106. Herpes Simplex type 1 Virus
107. Herpes Simplex type 2 Virus
108. HIV-1 (AIDS Virus)
109. Human Coronavirus
110. Influenza A/Brazil Virus
111. Influenza A/Victoria (H3N2) Virus
112. Influenza A2-Asian Virus
113. Influenza B Virus (Allen strain)
114. Influenza C Virus (Taylor strain)
115. Measles Virus
116. Parainfluenza Virus type 1
117. Poliovirus type 1 (Chat strain) 30 minutes contact time
118. Respiratory Syncytial Virus
119. Rotavirus
120. Vaccinia Virus



**Animal Premise Virucidal\* Performance:** At 0.35 ounces per gallon (1:365) use level, this product was evaluated in the presence of 5% serum with a 10 minute contact time and found to be effective against the following viruses on hard, non-porous environmental surfaces:

*Respiratory illnesses attributable to Pandemic 2009 H1N1 are caused by influenza A virus. This product (Product Name) is a broad-spectrum hard surface disinfectant that has been shown to be effective against (influenza A virus tested and listed on the label) and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).*

*This product has demonstrated effectiveness against influenza A virus and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 influenza A virus.*

*This product has demonstrated effectiveness against (influenza A virus tested and listed on the label) and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).*

*Kills Pandemic 2009 H1N1 influenza A virus (formerly called swine flu).*

*Kills Pandemic 2009 H1N1 influenza A virus.*

#### **Non-Human Viruses**

121. Avian Influenza/Turkey/Wisconsin Virus
122. Canine Coronavirus
123. Canine Distemper Virus
124. Canine Herpesvirus
125. Equine Herpesvirus
126. Equine Influenza Virus
127. Feline Calicivirus
128. Norovirus
129. Feline Infectious Peritonitis Virus
130. Infectious Bovine Rhinotracheitis (IBR) Virus
131. Newcastle Disease Virus
132. Porcine Parvovirus
133. Porcine Respiratory & Reproductive Syndrome Virus (PRRSV)
134. Porcine Rotavirus
135. Pseudorabies Virus
136. Transmissible Gastroenteritis (TGE) Virus
137. T1 bacteriophage
138. T4 bacteriophage
139. Vesicular Stomatitis Virus (VSV)
140. Bovine Viral Diarrhea Virus (BVDV)
141. Avian Influenza Virus (H5N1)
142. Influenza A Virus (H1N1) (formerly called swine flu)