FM 1? 01282-4 194

SEVIN® brand 10% carbaryl insecticide BAIT GRANULES

ACCEPTED
SEP 23 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

FOR FIELD, FRUIT AND VEGETABLE CROPS
- LAWN, GARDEN & PREMISE INSECT CONTROL
- FOR FLOWERS, ORNAMENTAL PLANT BEDS, TURFGRASS LAWNS,
- USE FOR OUTDOOR PEST CONTROL
- CONTROLS FIRE ANTS

ACTIVE INGREDIENT	
Carbaryl (1-naphthyl N-methylcarbamate)	10.04% by wt
INERT INGREDIENTS	89.96% by wt
TOTAL	100.00% by wt

EPA REG NO. 61282-4 EPA EST. NO 61282-WI-1

SEVIN is the registered trademark of Rhone-Poulenc for carbaryl insecticides

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS
CAUTION
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

MAY BE HARMFUL IF SWALLOWED Avoid breathing dust. Do not take internally. Avoid contact with eyes, skin or clothing. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO PHYSICIAN below.

STATEMENT OF PRACTICAL TREATMENT

GENERAL
Contact a physician immediately in all cases of ingestion or suspected poisoning. Transport to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.

IF SWALLOWED: Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Move from contaminated atmosphere and call a physician.

NOTE TO PHYSICIAN
Carbaryl is a carbamate insecticide which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Specific treatment consists of parasympathetic atropine sulfate. Caution should be maintained to prevent overparasympathetic. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized. Then intramuscularly every 30 to 60 minutes, if needed, to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning equipment or disposal of wastes.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

STORAGE AND DISPOSAL

STORAGE Store unused product in original container only in cool, dry area out of reach of children and animals. Do not store in areas where temperatures regularly exceed 100° F.

PESTICIDE DISPOSAL Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If bag is burned, stay out of smoke.

IMPORTANT PRE HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing residues SEVIN residues when carbaryl is used in accordance with label directions. If SEVIN carbaryl insecticide is used in accordance with label directions these crops, including bean vines, carrot tops, sweet corn silage, sweet corn tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in illegal residues in milk or meat.

Do not graze rotational food and feed crops not listed on this or other carbaryl labels in carbaryl treated soil.

Do not use reclaimed irrigation water from areas treated with carbaryl on upland crops for which carbaryl tolerances are not established.

Broadcast applications may be made with either ground equipment or aircraft. Treatments may be repeated as necessary. Dosages from 10-20 pounds per acre of SEVIN brand 10% Bait Carbaryl insecticide will control cutworms, armyworms, crickets, darkling ground beetles, grasshoppers, mole crickets and sowbugs (pillbugs).

HACCO, INC.

FIELD AND VEGETABLE CROPS

CROP	DOSAGE/ ACRE	PRE-HARVEST INTERVAL	SPECIFIC DIRECTIONS
Wheat	5-15 Pounds	None Forage/ 21 Days Grain	Do not make more than 2 applications after grain heads emerge from boot. Do not apply within 21 days of harvest of grain. No limit on emergence on green wheat used as pasture or forage.
Cucumbers squash melons peanuts	10 pounds	None	
Peas	pounds	Days	
Alfalfa Cotton	15 pounds	Days	
Beans carrots corn eggplants okra peppers potatoes tobacco tomatoes and sweet corn	10-20 pounds	None	
Pastures Rangeland Wasteland	10-20 pounds	None	Treatment may be repeated as necessary. Lower rate is suggested for early instar grasshoppers. Roses and armyworms on small plants or sparse vegetation. Use higher rate when grasshoppers and crickets are mature and armyworms are in late instar or when vegetation is dense.
Asparagus strawberries apples cherries peaches pears plums	10 pounds	Day	
Grapes	20 pounds	Days	
Broccoli Brussels sprouts cabbage cauliflower collards mixed lettuce garden beets (roots) horseradish parsnips radishes rutabagas and turnips (roots) apricots and nectarines	20 pounds	3 Days	
Sugar beets cilantro endive escarole garden beets tops kale leaf lettuce parsley spinach Swiss chard and turnips tops	20 pounds	14 Days	
Figs	10-20 pounds	Days	

LAWNS, RECREATIONAL, AND ORNAMENTAL TURF AREAS

For control of turfgrass pests, apply granules uniformly to the turfgrass. For optimum control, follow specific directions for proper watering after application.

SITE	INSECT	OUNCES OF SEVIN brand 10% BAIT /1000 SQUARE FEET	SPECIFIC DIRECTIONS
Turfgrass	Armyworms Crickets Cutworms Darkling ground beetles Grasshoppers Mole crickets Sowbugs	4 - 8	Treat entire area to obtain uniform coverage. Using particular attention to bunding permeters, edges of sidewalks and along porch and patio foundations. Use the lower rate before insects flared up and before turf is damaged. Use the higher rate when insects are numerous and turf damage is imminent. For best results mow lawn, remove clippings and apply material when grass blades are dry. Water lightly to carry granules to turf for 15 minutes after application. Keep children and pets off treated area until grass is dry. Repeat application may be made at 30 day intervals or as needed for control of Sod Webworms, Chinch Bugs or Ants.
Ornamental plants	Armyworms Crickets Cutworms Darkling ground beetles Grasshoppers Mole crickets Sowbugs	4 - 8	Apply uniformly to soil around plants, shrubs and trees. For best results, thoroughly water area to be treated prior to application.
Lawns, Recreational and Ornamental Turf Areas	Ants Centipedes Chiggers Chinch bugs Earwigs Elders sheller European chafer Fall armyworm Flea hopper Fleas Green June beetle Green June beetle grubs Lawn moths Sod webworm Leafhoppers Lucerne moth Sphinx Millipedes Sowbugs Springtails Ticks Webworms Armyworm	4 - 8	Treat entire area to obtain uniform coverage. Using particular attention to bunding permeters, edges of sidewalks and along porch and patio foundations. Use the lower rate before insects flared up and before turf is damaged. Use the higher rate when insects are numerous and turf damage is imminent. For best results mow lawn, remove clippings and apply material when grass blades are dry. Water lightly to carry granules to turf for 15 minutes after application. Keep children and pets off treated area until grass is dry. Repeat application may be made at 30 day intervals or as needed for control of Sod Webworms, Chinch Bugs or Ants.

FOUNDATION "INSECT BARRIER" TREATMENT, FLOWER BEDS, ORNAMENTAL PLANTINGS

Apply 1/2 Lb per 1,000 Sq. Ft. (22 X 22) to all soil areas for use as contact and bait control of Earwigs, Cutworms, Armyworms, Crickets, Sowbugs, Millipedes, Ants and for contact control of Fleas around the outside of buildings, under porches and foundation crawl spaces, in and around flower and ornamental plantings, along sod cracks next to buildings and as an insect barrier in a band 5 feet wide surrounding the foundation. For the insect barrier, sprinkle 1/4 Lb. of granules in a 5 foot wide band 100 feet in length (500 Sq. Ft.).

IMPORTED FIRE ANTS

For control of Imported Fire Ants, apply one to two ounces to each mound. Immediately irrigate with a minimum of one gallon of water to flush the chemical from the granules to the inside of the mound. Do not use any less gallons of water as flooding of the mound will result in poor control. Apply water gently to avoid destroying ants and causing migration from the mound. For best results, apply in the early morning or late evening hours when ants are most active. Treat new mounds as they appear.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label, (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions, and (c) that the directions, warnings and other statements on the label were prepared with reasonable evaluation of reasonable risks of effectiveness, of toxicity to non-target animals, and to plants, and of residues on food crops, and upon reports of tests ascertained. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES EXPRESS OR IMPLIED AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NET WEIGHT
50 POUNDS (22.68 KILOS)

BEST AVAILABLE COPY