

**SNAIL AND SLUG LG  
PELLETED BAIT**

**ATTRACTS AND KILLS  
SNAILS AND SLUGS!**

**BERRY CROPS-VEGETABLES,  
SETS, & SEEDLINGS  
AROUND PATIOS & FLOWER BOXES  
DICHONDRA & TURFGRASS LAWNS  
FRUIT & NUT TREES  
FLOWERS, ORNAMENTALS AND  
LANDSCAPE PLANTINGS**

**ACCEPTED**  
**MAR 30 1995**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
or amended, for the pesticide  
registered under  
EPA Reg. No. 61282-10

**ACTIVE INGREDIENT:**

Metalddehyde (2,4,6,8-tetramethyl-  
1,3,5,7-tetraoxocyclo-  
octane)..... 3.5%  
INERT INGREDIENTS:..... 96.5%  
TOTAL 100.0%

EPA REG. NO. 61282-10  
EPA EST. NO. 61282-WI-1

**KEEP OUT OF REACH OF CHILDREN 12 pt.\*  
CAUTION 18 pt.\***

**STATEMENTS OF PRACTICAL TREATMENT**  
If swallowed: Call a physician or  
Poison Control Center immediately.  
Drink 1 or 2 glasses of water and  
induce vomiting by touching back of  
throat with finger. Do not induce  
vomiting or give anything by mouth  
to an unconscious or convulsing  
person. If on skin: Wash with soap  
and warm water. If in eyes: Flush  
with flowing water. Get medical  
attention if irritation persists.

**THIS PESTICIDE MAY BE FATAL TO  
DOGS OR OTHER PETS IF EATEN  
KEEP PETS OUT OF TREATED AREA  
SEE SIDE/BACK PANEL FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS**

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS AND  
DOMESTIC ANIMALS**  
**DO NOT TAKE INTERNALLY, DO NOT GET  
IN EYES OR ON SKIN. CAUTION:** May  
be harmful if swallowed, inhaled or

\*Assumes size of front panel greater  
than or equal to 30 sq. inches.

absorbed through the skin. Do not  
breathe dust, get in eyes, on skin  
or on clothing. Wash thoroughly with  
soap and warm water after handling.  
Wash contaminated clothing with soap  
and warm water before re-use. If  
poisoning occurs, obtain prompt  
medical aid. Keep bait away from dogs  
and pets. Bait may be attractive to  
dogs. Confine pets during  
application to prevent them from  
believing they are being fed.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, to  
areas where surface water is present  
or to intertidal areas below the  
mean high water mark. Do not  
contaminate water when disposing of  
equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to  
use this product in a manner  
inconsistent with its labeling.  
Read entire label and use in  
accordance with precautionary  
statements and directions. Do not  
apply directly onto plants. Do not  
allow bait to contact the edible  
portions of any food crop.

**STORAGE AND DISPOSAL 12 pt.\***

Do not contaminate water, food or  
feed by storage or disposal. **STORAGE:**  
Store product in original labeled  
container in a cool, dry, locked  
place out of reach of children. Keep  
container tightly closed when not in  
use. Do not store in direct  
sunlight. **DISPOSAL:** Do not reuse  
empty container. Securely wrap  
original container in several layers  
of newspaper and discard in trash.

**USE PRECAUTIONS**

Read entire label before using. Use  
only as a baiting treatment to be  
applied directly to the soil area  
beneath plants as listed on this  
label. Do not allow bait to come in  
contact with parts of plant that may  
be used for food or feed purposes.  
The grower is responsible for  
residues on his crops, as well as  
for damages caused by drift from his  
property to that of others. Do not  
allow to drift to areas occupied by  
unprotected humans or beneficial  
animals or onto adjoining food,  
fiber or pasture crops. **NOTE** Snail  
& Slug LG Pelleted Bait should never  
be applied to dry soil. Applications  
should be made following an  
irrigation for best results. Apply  
at any time of the day but evening  
applications are preferred. Do not  
re-water for 48 hours after  
application.

GENERAL INFORMATION

Snails and Slugs generally feed during the night and damage many varieties of plants and plant seedlings. They inhabit damp moist areas of the soil surface around decaying refuse, organic matter, beneath boards, rocks and trash piles and hide at the base of growing plants outdoors and in the greenhouse. Their presence can be detected by observing the shiny trails which they leave on sidewalks, driveways and patios.

The Use Profile for Snail and Slug LG includes the following Sites and the Listed Plants in Terrestrial Food Crops and Terrestrial Non-Food Crops For Home Greenhouse Flowers/Ornamentals and Domestic Outdoor Areas. Uses are as follows.

HOW MUCH AND WHEN TO USE

For control of Snails and Slugs in the following areas, use 1 lb. to treat 1,100 to 1,750 Sq. Ft., unless otherwise stated. Snails and Slugs attack the tender growing tips of plants devouring the entire tip or sometimes eating holes in the new leaves which are yet unfolded. When the leaves expand and open, this feeding appears as a "row of holes" in the leaf, all in a line. This is one "sign" of Snail or Slug damage to your plants. Apply when this first plant damage appears or is anticipated and repeat as needed for control. Pour Pellets directly from container. Do not dispense by hand unless wearing protective gloves. Apply down wind; do not breathe dust.

FOR THE BERRY PATCH

Blackberries, Blueberries, Boysenberries, Currants, Dewberries, Gooseberries, Loganberries, Raspberries, Strawberries  
For soil surface coverage. Apply 1 lb. to treat 1,100 to 3,600 sq.ft.

TO PROTECT VEGETABLES, SETS AND SEEDLINGS IN THE GARDEN

Apply 1 lb. to treat 1,100 to 1,750 sq. ft.  
Alfalfa, Artichoke, Asparagus, Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupe, Carrots, Cauliflower, Cereal Crops, Celery, Clover, Cole Crops, Collards, Corn, Cowpeas, Cucumbers, Eggplant, Endive, Garlic, Ginseng, Horseradish, Kale, Kohlrabi, Leeks, Legumes, Lettuce, Melons, Muskmelons, Mustard Greens, Okra, Onions, Parsnips, Peas, Peppers, Pimentos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas, Salsify, Shallots, Spinach, Squash, Sweet Potatoes,

Swiss Chard, Tomatoes, Turnips, Watermelons.

FOR HERB GARDENS

Anis, Caraway, Catnip, Chicory, Chives, Coriander, Dill, Mustard, Oregano, Parsley, Peppermint, Sage, Savory, Spearmint, Thyme. Apply 1 lb. to treat 1,100 to 1,750 sq. ft.

FOR CONTROL OF SNAILS & SLUGS AROUND PATIOS, PORCHES, PREMISE AND POOL AREAS

Snail & Slug LG Pelleted Bait attracts and kills snails and slugs. Sprinkle pellets on moist areas where these pests have been seen or their "shiny" trails (from the night before) are seen on the sidewalk or patio stones. Keep pets out of area while baiting and until pellets have expanded and disintegrated into the soil surface. Apply 1 pound of pellets to cover 1,750 sq. ft. (35 ft. x 50 ft. area). Repeat in 5 to 7 days or as needed for control.

HOME DICHONDRA AND TURFGRASS LAWNS  
Water infested area thoroughly before application. Broadcast SNAIL & SLUG LG Pelleted Bait directly on lawn or turf. Spread evenly. Apply 1 lb. to treat 1,100 to 1,750 sq. ft. Do not re-water for 48 hours.

HOME ORCHARDS

Almonds, Apples, Apricots, Avocados, Bananas, Cherries, Citrus (Oranges, Lemons, Limes, Grapefruit, etc.), Grapes, Nectarines, Peaches, Pears, Pecans, Plums, Stone Fruits, Walnuts  
Broadcast liberally under trees, and any other places where slugs and snails may congregate. Apply 1 lb. to treat 1,100 to 1,750 sq. ft.

TO PROTECT FLOWERS, FOLIAGE PLANTS AND SEEDLINGS GROWING IN BENCHES OR FLATS IN THE HOME GREENHOUSE, LATHE HOUSE, OR COLD FRAME

Orchids, Ornamental Flowering Plants and Foliage Plants. Do not apply directly to plants unless prior experience has shown the plant to be tolerant to this product. Most plants are tolerant however some slight tip burning has been noticed on extremely tender orchids. One pound will cover up to 1,100 sq. ft. Do not re-water for 48 hours.

FOR HOME FLOWER, ORNAMENTAL AND LANDSCAPE PLANTINGS - HOME ORNAMENTAL GARDENS, PLANTED BORDERS AND GROUND COVERS

Including Azaleas, Begonias, Camellias, Chrysanthemums, Daffodils, Daisies, Geraniums, Honeysuckle, Hydrangeas, Ice Plant, Impatiens,

Iris, Marigolds, Pansies, Petunias, Rhododendron, Rose, Sweet Peas, Snapdragons, Spathiphyllum, Tulips. Snails and Slugs attack the tender growing tips of plants, sometimes eating holes in the new leaves which are yet unfolded. When the leaves expand and open, this feeding appears as a "row of holes" in the leaf, all in a line. This is a "sign" of Snail or Slug damage. Water the infested area thoroughly before application. Broadcast or sprinkle Snail and Slug LG Pelleted Bait evenly over the infested areas and along edges of flower beds adjacent to border or ground cover plants. Also apply on ground at base of fences and walls in shady, protected locations. Apply 1 lb. to treat 1,100 to 1,750 sq. ft. Do not re-water for 48 hours.

NOTICE: Sellers warranty shall be limited to the terms of the label and subject thereto. The buyer and user assumes any risk to persons or property arising out of the use or handling of this product and accepts the product on these conditions.

NET WEIGHT: (1,2,2½,5,10,25) POUNDS

HACCO, INC.  
P.O. BOX 7190  
MADISON, WI 53707