

61282-10

11/20/2003

1/2

SNAIL AND SLUG L&G PELLETED BAIT

ATTRACTS AND KILLS SNAILS and SLUGS!

**BERRY CROPS-VEGETABLES
SETS, & SEEDLINGS
AROUND PATIOS & FLOWER BOXES
DICHONDRA & TURFGRASS LAWNS
FRUIT & NUT TREES AND
LANDSCAPE PLANTINGS**

ACTIVE INGREDIENT:

Metaaldehyde (2, 4, 6, 8-tetramethyl- 1, 3, 5, 7,-tetraoxycyclooctane	3.5%
INERT INGREDIENTS:	96.5%
TOTAL	100.0%

EPA REG. NO. 61282-10

EPA EST NO. 61282-WI-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

Have label with you when obtaining treatment advice.

If Swallowed	<ul style="list-style-type: none"> Call a poison control center, doctor, or 1-800-301-7976 immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.
If on Skin	<ul style="list-style-type: none"> Wash with plenty of soap and water.
If in Eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing.
<ul style="list-style-type: none"> Call a poison control center, doctor or 1-800-301-7976 for treatment advice. 	

**THIS PESTICIDE MAY BE FATAL TO
DOGS OR OTHER PETS IF EATEN
KEEP PETS OUT OF TREATED AREA
SEE SIDE/BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS**

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

DO NOT TAKE INTERNALLY, DO NOT GET IN EYES OR ON SKIN. CAUTION: May be harmful if swallowed, inhaled or absorbed through the skin. Do not breathe dust, get in eyes, on skin or on clothing. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and warm water before re-use. If poisoning occurs, obtain prompt medical aid. Keep bait away from dogs and pets. Bait may be attractive to dogs. Confine pets during application to prevent them from believing they are being fed.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label and use in accordance with precautionary statements and directions. Do not apply directly onto plants. Do not allow bait to contact the edible portions of any food crop.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children. Keep container tightly closed when not in use. Do not store in direct sunlight.

CONTAINER DISPOSAL: If empty: Do not reuse empty container. Place in trash or offer for recycling if available. If partly filled: Call local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.


USE PRECAUTIONS

Read entire label before using. Use only as a baiting treatment to be applied directly to the soil area beneath plants as listed on this label. Do not allow bait to come in contact with parts of plant that may be used for food or feed purposes. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others. Do not allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. NOTE: Snail & Slug LG Pelleted Bait should never be applied to dry soil. Applications should be made following an irrigation for best results. Apply at any time of the day, but evening applications are preferred. Do not re-water for 48 hours after application.

GENERAL INFORMATION

Snails and Slugs generally feed during the night and damage many varieties of plants and plant seedlings. They inhabit damp moist areas of the soil surface around decaying refuse, organic matter, beneath boards, rocks and trash piles and hide at the base of growing plants outdoors and in the greenhouse. Their presence can be detected by observing the shiny trails which they leave on sidewalks, driveways and patios.

The Use profile for Snail and Slug LG includes the following Sites and the Listed Plants in Terrestrial Food Crops and Terrestrial Non-Food Crops for Home Greenhouse Flowers/Ornamentals and Domestic Outdoor Areas. Uses are as follows.



HOW MUCH AND WHEN TO USE

For control of Snails and Slugs in the following areas, use 1 lb. to treat 1,100 to 1,750 sq. ft., unless otherwise stated. Snails and Slugs attack the tender growing tips of plants devouring the entire tip or sometimes eating holes in the new leaves which are yet unfolded. When the leaves expand and open, this feeding appears as a "row of holes" in the leaf, all in a line. This is one "sign" of Snail or Slug damage to your plants. Apply when this first plant damage appears or is anticipated and repeat as needed for control. Pour Pellets directly from container. Do not dispense by hand unless wearing protective gloves. Apply down wind; do not breathe dust.

FOR THE BERRY PATCH

- Blackberries, Blueberries,
- Boysenberries, Currants, Dewberries,
- Gooseberries, Loganberries,
- Raspberries, Strawberries

For soil surface coverage. Apply 1 lb. to treat 1,100 to 3,600 sq. ft.

TO PROTECT VEGETABLES, SETS AND SEEDLINGS IN THE GARDEN

Apply 1 lb. to treat 1,100 to 1,750 sq. ft.

- Alfalfa, Artichoke, Asparagus, Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupe, Carrots, Cauliflower, Cereal Crops, Celery, Clover, Cole Crops, Collards, Corn, Cowpeas, Cucumbers, Eggplant, Endive, Garlic, Ginseng, Horseradish, Kale, Kohlrabi, Leeks, Legumes, Lettuce, Melons, Muskmelons, Mustard Greens, Okra, Onions, Parsnips, Peas, Peppers, Pimentos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas, Salsify, Shallots, Spinach, Squash, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips, Watermelons.

FOR HERB GARDENS

- Anis, Caraway, Catnip, Chicory, Chives, Coriander, Dill, Mustard, Oregano, Parsley, Peppermint, Sage, Savory, Spearmint, Thyme. Apply 1 lb. to treat 1,100 to 1,750 sq. ft.

FOR CONTROL OF SNAILS & SLUGS AROUND PATIOS, PORCHES, PREMISE AND POOL AREAS

Snail & Slug LG Pelleted Bait attracts and kills snails and slugs. Sprinkle pellets on moist areas where these pests have been seen or their "shiny" trails (from the night before) are seen on the sidewalk or patio stones. Keep pets out of area while baiting and until pellets have expanded and disintegrated into the soil surface. Apply 1 lb. of pellets to cover 1,750 sq. ft. (35 ft. x 50 ft. area). Repeat in 5 to 7 days or as needed for control.

HOME DICHONDRA AND TURFGRASS LAWNS

Water infested area thoroughly before application. Broadcast Snail & Slug LG Pelleted Bait directly on lawn or turf. Spread evenly. Apply 1 lb. to treat 1,100 to 1,750 sq. ft. Do not re-water for 48 hours.

HOME ORCHARDS

Almonds, Apples, Apricots, Avocados, Bananas, Cherries, Citrus (Oranges, Lemons, Limes, Grapefruit, etc.), Grapes, Nectarines, Peaches, Pears, Pecans, Plums, Stone Fruits, Walnuts. Broadcast liberally under trees, and any other places where slugs and snails may congregate. Apply 1 lb. to treat 1,100 to 1,750 sq. ft.

TO PROTECT FLOWERS, FOLIAGE PLANTS AND SEEDLINGS GROWING IN BENCHES OR FLATS IN THE HOME GREENHOUSE, LATHE HOUSE, OR COLD FRAME

Orchids, Ornamental Flowering Plants and Foliage Plants. Do not apply directly to plants unless prior experience has shown the plant to be tolerant to this product. Most plants are tolerant however some slight tip burning has been noticed on extremely tender orchids. One pound will cover up to 1,100 sq. ft. Do not re-water for 48 hours.

FOR HOME FLOWER, ORNAMENTAL AND LANDSCAPE PLANTINGS - HOME ORNAMENTAL GARDENS, PLANTED BORDERS AND GROUND COVERS

Including Azaleas, Begonias, Camellias, Chrysanthemums, Daffodils, Daisies, Geraniums, Honeysuckle, Hydrangeas, Ice Plant, Impatiens, Iris, Marigolds, Pansies, Petunias, Rhododendron, Rose, Sweet Peas, Snapdragons, Spathiphyllum, Tulips. Snails and Slugs attack the tender growing tips of plants, sometimes eating holes in the new leaves which are yet unfolded. When the leaves expand and open, this feeding appears as a "row of holes" in the leaf, all in a line. This is a "sign" of Snail or Slug damage. Water the infested area thoroughly before application. Broadcast or sprinkle Snail and Slug LG Pelleted Bait evenly over the infested areas and along edges of flower beds adjacent to border or ground cover plants. Also apply on ground at base of fences and walls in shady, protected locations. Apply 1 lb. to treat 1,100 to 1,750 sq. ft. Do not re-water for 48 hours.

NOTICE: Sellers warranty shall be limited to the terms of the label and subject thereto. The buyer and user assumes any risk to persons or property arising out of the use or handling of this product and accepts the product on these conditions.

NET WEIGHT: (1,2,2½, 5,10,25) POUNDS

HACCO, INC.
P.O. BOX 7190
MADISON, WI 53707