

PM 12 11 2 - 1 8-17

MAR 23 1995

Judith A. Thompson  
Hacco, Inc.  
P.O. Box 7190  
Madison, WI 53707

Dear Ms. Thompson:

Subject: Revised labeling to add aerial application  
Snail and Slug AG Pelleted Bait  
EPA Registration No. 61282-11  
Amendment application of January 12, 1995

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable provided that you make the labeling changes indicated below before you release the product for shipment bearing the amended labeling:

Change the word "formulation" under the directions for use to "formation".

A stamped copy is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.  
Product Manager (19)  
Insecticide-Rodenticide Branch  
Registration Division (7505C)

Enclosure

1985

**SNAIL and  
SLUG AG  
PELLETED BAIT**

**ATTRACTS AND KILLS  
SNAILS and SLUGS!**

**BERRY CROPS - CITRUS GROVES  
APPLE & PEACH ORCHARDS  
IN GREENHOUSES - LATH HOUSES  
VEGETABLE, HERB & ORNAMENTAL  
GARDENS  
DICHOONDRA & TURFGRASS LAWNS**

**ACTIVE INGREDIENT:**  
Metaldehyde (2,4,6,8,-tetramethyl-  
1,3,5,7,-tetraoxycyclo-  
octane) ..... 3.5%  
**INERT INGREDIENTS:** ..... 96.5%  
Total..... 100.0%

EPA REG. NO. 61282-11  
EPA EST. NO. 61282-WI-1  
KEEP OUT OF REACH OF CHILDREN (12  
pt.)\*

**CAUTION (12 pt.)\***  
**STATEMENTS OF PRACTICAL TREATMENT**  
**IF SWALLOWED:** Call a physician or  
Poison Control Center immediately.  
Drink 1 or 2 glasses of water and  
induce vomiting by touching back of  
throat with finger. Do not induce  
vomiting or give anything by mouth  
to an unconscious or convulsing  
person.  
**IF ON SKIN:** Wash with soap and warm  
water. **IF IN EYES:** Flush with flowing  
water. Get medical attention if  
irritation persists.

**THIS PESTICIDE MAY BE FATAL  
TO DOGS OR OTHER PETS IF EATEN  
KEEP PETS OUT OF TREATED AREA**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS  
DO NOT TAKE INTERNALLY.  
DO NOT GET IN EYES OR ON SKIN.**

**CAUTION:** May be harmful if swallowed,  
inhaled or absorbed through the skin.  
Do not breathe dust, get in eyes, on  
skin or on clothing. If poisoning  
occurs, obtain prompt medical aid.  
Keep bait away from dogs and pets.  
Bait may be attractive to dogs.  
Confine pets during application to  
prevent them from believing they are  
being fed.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

**MAR 23 1995**

\*Assumes a front panel of over  
30 sq. in.  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
61282-11

**PERSONAL PROTECTIVE EQUIPMENT**  
Applicators and other handlers must  
wear long-sleeved shirt and long  
pants, waterproof gloves and shoes  
plus socks.  
Follow manufacturer's instructions  
for cleaning and maintaining Personal  
Protective Equipment (PPE). If no  
such instructions for washables, use  
detergent and hot water. Keep and  
wash PPE separately from other  
laundry.  
When handlers use closed systems,  
enclosed cabs, or aircraft in a  
manner that meets the requirements  
listed in the Worker Protection  
Standard (WPS) for agricultural  
pesticides (40 CFR 170.240(d)(4-6))  
the handler PPE requirements may be  
reduced or modified as specified in  
WPS.

**USER SAFETY RECOMMENDATIONS**  
Users should wash hands before  
eating, drinking, chewing gum,  
using tobacco, or using the  
toilet. Remove clothing  
immediately if pesticide gets  
inside. Then wash thoroughly and  
put on clean clothing.

**ENVIRONMENTAL HAZARDS**  
Do not apply directly to water, to  
areas where surface water is present  
or to intertidal areas below the  
mean high water mark. Do not  
contaminate water when disposing of  
equipment washwaters.

**DIRECTIONS FOR USE**  
It is a violation of Federal law to  
use this product in a manner  
inconsistent with its labeling.  
Read entire label before using this  
product.  
Do not apply this product in a way  
that will contact workers or other  
persons, either directly or through  
drift. Only protected handlers may  
be in the area during application.  
For any requirements specific to  
your State or Tribe, consult the  
agency responsible for pesticide  
regulation.  
Apply as a soil treatment. Terres-  
trial food and non-food crops may be  
broadcast by air or ground equipment  
unless specified otherwise on the  
label. Apply 24 to 40 pounds of  
pelleted bait per acre unless  
otherwise stated. Use a band  
treatment between the rows after  
formulation of edible parts. Do not  
contaminate edible portions of the  
plant.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes plus socks.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area being treated must be vacated and not re-entered by persons without shoes.

**USE PRECAUTIONS**

Read entire label before using. Use only as a baiting treatment to be applied directly to the soil area beneath plants as listed on this label. Do not allow bait to come in contact with parts of plant that may be used for food or feed purposes. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others. Do not allow to drift to areas occupied by beneficial animals or onto adjoining

food, fiber or pasture crops. NOTE: Snail & Slug AG Pelleted Bait should never be applied to dry soil. Applications should be made following an irrigation for best results. Apply anytime of the day, but evening applications are preferred. Do not re-water for 48 hours after application.

**GENERAL INFORMATION**

Snails and Slugs generally feed during the night and damage many varieties of plants and plant seedlings. They inhabit damp moist areas of the soil surface around decaying refuse, organic matter, beneath boards, rocks and trash piles and hide at the base of growing plants outdoors and in the greenhouse. Their presence can be detected by observing the shiny trails left on sidewalks, driveways and patio blocks. The Use Profile for Snail and Slug AG includes the following Sites and the Listed Crops and/or Areas in - Terrestrial Food Crops; Terrestrial Non-Food Crops; Greenhouse Food Crops; Greenhouse Flowers/Ornamentals; Domestic and Nondomestic Outdoor Areas; and Indoor uses. Uses are as follows:

**APPLICATION RATE**

For control of SNAILS and SLUGS in the following areas, use 1 lb. to treat 1,100 to 1,750 Sq. Ft. (equivalent to 24 to 40 pounds of pelleted bait per acre), unless otherwise stated. Apply when first plant damage appears or is anticipated and repeat as needed for control.

**VEGETABLE AND FOOD CROPS**

Apply 1 lb. to treat  
1,100 to 1,750 sq. ft.

Alfalfa, Artichoke, Asparagus, Beans, Beets, Black-eyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupe, Carrots, Cauliflower, Cereal Crops, Celery, Clover, Cole Crops, Collards, Corn, Cowpeas, Cucumbers, Eggplant, Endive, Garlic, Ginseng, Horseradish, Kale, Kohlrabi, Leeks, Legumes, Lettuce, Melons, Muskmelons, Mustard Greens, Okra, Onions, Parsnips, Peas, Peppers, Pimentos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas, Salsify, Shallots, Spinach, Squash, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips, Watermelons, Wheat.

Artichokes: By air or ground application before tight bud stage, apply 25 to 70 lbs./acre. After tight bud stage, apply by ground application only.

#### FRUIT AND NUT CROPS

Almonds, Apples, Apricots, Avocados, Bananas, Cherries, Citrus (Oranges, Lemons, Limes, Grapefruit, etc.), Grapes, Nectarines, Peaches, Pears, Pecans, Plums, Stone Fruits, Walnuts.  
Broadcast liberally under trees, around irrigation pipes, farm buildings and any other place where slugs and snails may congregate. Apply 1 lb. to treat 1,100 to 1,750 sq. ft.

#### BERRY CROPS

Blackberries, Blueberries, Boysenberries, Currants, Dewberries, Gooseberries, Loganberries, Raspberries, Strawberries.

For soil surface coverage; apply 1 lb. to treat 1,100 to 3,600 sq. ft. (12 to 39 lbs./acre).

#### HERBS

Anis, Caraway, Catnip, Chicory, Chives, Coriander, Dill, Mustard, Oregano, Parsley, Peppermint, Sage, Savory, Spearmint, Thyme

Apply 1 lb. to treat 1,100 to 1,750 sq. ft.

#### FLOWER AND ORNAMENTAL PLANTINGS

ORNAMENTAL GARDENS, PLANTED BORDERS AND GROUND COVERS: Including Azaleas, Begonias, Camellias, Chrysanthemums, Daffodils, Daisies, Geraniums, Honeysuckle, Hydrangeas, Ice Plant, Impatiens, Iris, Marigolds, Pansies, Petunias, Rhododendron, Rose, Sweet Peas, Snapdragons, Spathiphyllum, Tulips.  
Water the infested area thoroughly before application. Broadcast or sprinkle Snail and Slug AG Pelleted Bait evenly over the infested areas and along edges of flower beds adjacent to border or ground cover plants. Also apply on ground at base of fences and walls in shady, protected locations. Apply 1 lb. to treat 1,100 to 1,750 sq. ft. Do not re-water for 48 hours.

#### DICHONDRA AND TURFGRASS LAWNS

Water infested area thoroughly before application. Broadcast SNAIL & SLUG AG Pelleted Bait directly on lawn or turf. Spread evenly. Apply 1 lb. to treat 1,100 to 1,750 sq. ft. Do not re-water for 48 hours.

#### GREENHOUSES AND LATHE HOUSES

Distribute pellets on and beneath benches, but do not place in pots or flats unless plants are well established. Keep pellets from coming in contact with plants. If soil is dry, irrigate prior to, or sprinkle lightly, after distributing pellets. For crops such as Tobacco, Asparagus, Beans, Beets, Cabbage, Carrots, Celery, Collards, Cucumbers, Eggplant, Endive, Kale, Kohlrabi, Lettuce, Melons, Mustard, Peas, Peppers, Radish, Spinach, Swiss Chard, Tomatoes, Turnips.  
Apply 1 lb. to treat 500 to 600 sq.

ft. Do not re-water for 48 hours.

NOTE: This product has proven to have a reasonable residual control, however for heavily infested greenhouses a second application may be needed in 7 to 10 days.

#### GREENHOUSE FLOWERS & ORNAMENTALS Orchids, Ornamental Flowering Plants and Foliage Plants.

Do not apply directly to plants unless prior experience has shown the plant to be tolerant to this product.

Most plants are tolerant however some slight tip burning has been noticed on extremely tender orchids. One pound will cover up to 1,100 sq. ft. Do not re-water for 48 hours.

#### SEED GRASSES

Alfalfa, Barley, Hay, Oats, Wheat and other grasses grown for seed. Apply up to 45 lbs. per acre.

#### SEED FOOD CROPS

Wheat, Alfalfa, Beans, Clover, Corn (Sweet), Grasses (Barley, Hay, Oats, Wheat), Peas, Potato. Apply 1 lb. to treat 1,100 to 1,750 sq. ft.

#### DOMESTIC AND NON-DOMESTIC OUTDOOR USES

Domestic Dwellings, Greenhouses, Lathe Houses, Lawns, Including Dichondra, Ornamental Grasses, Ornamental Plants (Including Nursery Stock), Turf

#### INDOOR USES

Farm Buildings: Do not allow pets or other animals access to bait. After control is achieved thoroughly sweep and clean the area.

#### STORAGE AND DISPOSAL (12 pt.)\*

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children. Keep container tightly closed when not in use. Do not store in direct sunlight.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State & local authorities, by burning. If burned stay out of smoke.

**NOTICE OF WARRANTY**

Manufacturer and Seller warrant that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Manufacturer and Seller. In no case shall Manufacturer and Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer.

Except as expressly provided herein, Manufacturer and Seller make no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage.

NET CONTENTS: 5 (10, 25, 50) Pounds

MANUFACTURED BY:  
HACCO, INC.  
P.O. Box 7190  
Madison, WI 53707 U.S.A.