

PM

19

61282-16

Page 1 of 11

JUL - 6 1994

Ms. Judith A. Thompson
HACCO, Inc.
P.O. Box 7190
Madison, WI 53707

Dear Ms. Thompson:

Subject: Label amendment dated April 12, 1994 submitted in
response to PR Notice 93-7
Sevin Brand 5% Carbaryl Insecticide Bait Granules
EPA Registration No. 61282-16

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments listed below. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Standard (WPS). Under the WPS labeling regulations of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed printed labeling AND

- WITHIN one year from the date of this acceptance

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,

DHE

Dennis H. Edwards, Jr.
Product Manager 13
Insecticide-Rodenticide Branch
Registration Division (7505C)

Attachment

3 of 11
ACCEPTED
with COMMENTS
in EPA Letter Dated

SEVIN brand 5% Carbaryl Insecticide BAIT Granules

For Field, Fruit and Vegetable Crops
Lawn, Garden & Premise Insect Control
For Flowers, Ornamental Plant Beds, Turfgrass Lawns,
Use for Outdoor Pest Control
To Attract and Kill Insect Pests Listed on this Label

JUL - 6 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
61282-16

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate)..... 5%

INERT INGREDIENTS: 95%

Total 100%

EPA Reg. No. 61282-16

EPA Est. No. 61282-WI-1

SEVIN is the registered trademark of Rhone-Poulenc for carbaryl
insecticides.

Self-Verified
Label

KEEP OUT OF REACH OF CHILDREN {12 pt¹}

CAUTION {18 pt¹}

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

MAY BE HARMFUL IF SWALLOWED. Avoid breathing dust. Do not take
internally. Avoid contact with eyes, skin, or clothing.

ANTIDOTE STATEMENT

ATROPINE SULFATE IS HIGHLY EFFECTIVE AS AN ANTIDOTE. See NOTE TO
PHYSICIAN below.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear long-sleeved shirt and long
pants, waterproof gloves and shoes plus socks.
Follow manufacturer's instructions for cleaning and maintaining
Personal Protective Equipment (PPE). If no such instructions for
washables, use detergent and hot water. Keep and wash PPE separately
from other laundry.

ENGINEERING CONTROLS STATEMENTS:

When handlers use closed systems, enclosed cabs, or aircraft in a
manner that meets the requirements listed in the Worker Protection
Standard (WPS) for agricultural pesticides {40 CFR 170.240(d)(4-6)}
the handler PPE requirements may be reduced or modified as specified
in the WPS.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum,
using tobacco, or using the toilet. Remove clothing immediately
if pesticide gets inside. Then wash thoroughly and put on clean
clothing.

STATEMENT OF PRACTICAL TREATMENT

Carbaryl is an N-Methyl Carbamate Insecticide.

¹Assumes a front panel of over 30 sq. in.

BEST AVAILABLE COPY

GENERAL

Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and **SHOW A COPY OF THIS LABEL TO THE PHYSICIAN.**

IF SWALLOWED: Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger.

IF IN EYES: Flush eyes with plenty of water. Get medical attention, if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Move from contaminated atmosphere and call a physician.

NOTE TO PHYSICIAN

Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible. Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes, if needed, to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedative should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates and toxic to birds and other wildlife. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Self-Verified
Label

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves and shoes, plus socks.

Self-Verified
Label

STORAGE AND DISPOSAL

STORAGE: Store unused product in orig. container only, in cool, dry area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100°F.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities by burning. If bag is burned, stay out of smoke.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION.

Tolerances established under Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when carbaryl is used in accordance with label directions. If SEVIN carbaryl insecticide is used in accordance with label direction, these crops, including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in illegal residues in milk or meat.

Do not plant rotational food and feed crops not listed on this or other carbaryl labels in carbaryl treated soil.

Do not use reclaimed irrigation water from crops treated with carbaryl on upland crops for which carbaryl tolerances are not established.

March 31, 1994

6 7 11

FIELD, FRUIT AND VEGETABLE CROPS

Broadcast applications may be made with either ground equipment or aircraft. Treatments may be repeated as necessary. Dosages from 20-40 pounds per acre of SEVIN brand 5% Carbaryl Insecticide Bait will control cutworms, armyworms, crickets, darkling ground beetles, grasshoppers, molecrickets and sowbugs (pillbugs).

CROP	DOSAGE/ ACRE	PRE- HARVEST INTERVAL	SPECIFIC DIRECTIONS
Wheat	20-30 pounds (7 1/3 - 11 ozs. per 1000 sq.ft.)	None (Forage) 21 Days (Grain)	Do not make more than 2 applications after grain heads emerge from boot. Do not apply within 21 days of harvest grain. No time limitation on green wheat used as pasture or forage.
Cucumbers, squash, melons, peanuts	20 pounds (7 1/3 ozs. per 1000 sq. ft.)	None	
Peas	30 pounds (11 ozs. per 1000 sq.ft.)	3 Days	
Alfalfa, Birdsfoot Trefoil (forage and Hay), Clover (pasture, forage), Cotton	30 pounds (11 ozs. per 1000 sq.ft.)	7 Days	
Beans, carrots, corn (field, sweet, pop) eggplants, okra, peppers, potatoes, Sorghum forage, Soybeans, Sweet potatoes, tobacco, tomatoes	20-40 pounds (7 1/3 - 14 2/3 ozs. per 1000 sq. ft.)	None	
Pastures, Rangeland, Wasteland	20-40 pounds (7 1/3 - 14 2/3 ozs. per 1000 sq. ft.)	None	Treatment may be repeated as necessary. Lower rate is suggested for early instar grasshoppers, crickets and armyworms on small plants or sparse vegetation. Use higher rate when grasshoppers and crickets are mature and armyworms are in late instars or when vegetation is dense.
Asparagus, strawberries, apples, cherries, grapes, peaches, pears, plums	20-40 pounds (7 1/3 - 14 2/3 ozs. per 1000 sq. ft.)	1 Day	

self-Verified
label

March 31, 1994

7 4 11

(Field, Fruit and Vegetable Crops, continued)

CROP	DOSAGE/ ACRE	PRE- HARVEST INTERVAL	SPECIFIC DIRECTIONS
Broccoli, Brussels sprouts, cabbage, cauliflower, cowpeas, Kohlrabi, head lettuce, garden beets (roots), horseradish, parsnips radishes, rutabagas, salsify (roots) and turnips (roots), apricots and nectarines	20-40 pounds (7 1/3 - 14 2/3 ozs. per 1000 sq. ft.)	3 Days	
Sugar beets, celery, Chinese Cabbage, collards, dandelion endive (escarole), garden beets (tops), Hanover Salad Kale, leaf lettuce, mustard greens parsley, salsify (tops), spinach, Swiss chard and turnips (tops)	20-40 pounds (7 1/3 - 14 2/3 ozs. per 1000 1000 sq.	14 Days	
Citrus	20-40 pounds (7 1/3 - 14 2/3 ozs. per 1000 sq. ft.)	5 Days	
Sorghums (milo, grain sorghum, sweet sorghum, and hybrids)	20-40 pounds (7 1/3 - 14 2/3 ozs. per 1000 sq. ft.)	21 (Grain)	

Self-
certified
Label

(Field, Fruit and Vegetable Crops, continued)

CROP	DOSAGE/ ACRE	PRE- HARVEST INTERVAL	SPECIFIC DIRECTIONS
Oats, barley and rye (fall treatment)	20-40 pounds (7 1/3 - 14 2/3 ozs. per 1000 sq. ft.)		Do not feed treated fodder to dairy animals or animals being finished for slaughter or allow them to graze on crops treated with this product.
Sunflowers	20-40 pounds (7 1/3 - 14 2/3 ozs. per 1000 sq. ft.)	60 (crop for oil, food or feed purposes) 0 (crop for seeding purposes only)	

Self-Verified
1

POULTRY HOUSES

SITE	INSECT	OUNCES OF SEVIN brand 5% BAIT/1000 SQUARE FEET	SPECIFIC DIRECTIONS
Poultry Houses, (chickens, turkeys, ducks, geese, gamebirds, and pigeons)	Darkling beetle (Lesser mealworm or litter beetle)	48	<p>Apply bait uniformly to floor or litter area using appropriate equipment such as chest-carried fertilizer/seed spreader. Do not incorporate. It is not necessary to remove birds during application, but uniform coverage may be difficult to achieve with birds present. Do not contaminate feeding or watering units.</p> <p>Optimum treatment timings are: 1. Ten to 14 days after the birds are placed in the house and 2. two weeks after timing "#1" above. One application at each of the two timings listed above should be made for heavy infestations. If only one applica- tion can be made, timing #1 should be followed.</p> <p>For best control of high darkling beetle populations: In addition to a bait application, walls, cracks, posts, crevice and insulation around ventilation areas should be treated with a sprayable or flowable formulation of SEVIN brand Carbaryl Insecticide. Spray applications should be made immediately following removal of the flock from the house or immediately prior to introducing the flock into the house. All label directions and restrictions for the application of the liquid SEVIN brand Carbaryl Insecticide formulation used should be carefully followed.</p> <p>Avoid spraying of nests, eggs and feeding and watering units.</p> <p>Do not treat poultry, game birds or their premises within 7 days of slaughter.</p>

9 9 11

LAWNS, RECREATIONAL, AND ORNAMENTAL TURF AREAS

For control of turfgrass pests, apply granules uniformly to the turf grass. For optimum control, follow specific directions for proper watering after application.

Self-Verified
Label

SITE	INSECT	OUNCES OF SEVIN brand 5% BAIT/1000 SQUARE FEET	SPECIFIC DIRECTIONS
Turfgrass	Armyworms, Crickets, Cutworms, Darkling ground beetle, Grass- hoppers, Mole crickets, Sowbugs	8 to 16	Treat entire area to obtain uniform coverage, giving particular attention to building perimeters, edges of sidewalks, and along porch and patio foundations. For best results, mow lawn, remove clippings, and thoroughly water area to be treated prior to application.
Ornamental plants	Armyworms, Crickets, Cutworms, Darkling ground beetle, Grass- hoppers, Mole crickets, Sowbugs	8 to 16	Apply uniformly to soil around plants, shrubs, and trees. For best results, thoroughly water area to be treated prior to application.
Lawns, Recreational & Ornamental Turf Areas	Ants, Centipedes, Chiggers, Chinch bugs, Earwigs, Essex skipper, European chafer, Fall army- worm, Fiery skipper, Fleas Green June beetle, Green June beetle grubs, Lawn moths (sod web- worm),	44 to 60	Treat entire area to obtain uniform coverage, giving particular attention to building perimeters, edges of sidewalks, and along porch and patio foundations. Use the lower rate before insects "build-up" and before turf is damaged. Use the higher rate when insects are numerous and turf damage is imminent. For best results mow lawn, remove clippings and apply material, then allow sprinklers to run for 15 minutes after application. Keep children and pets off treated area until grass is dry. Repeat application may be made at 30 day intervals or as needed for control of Sod Webworms, Chinch Bugs or Ants.

LAWNS, RECREATIONAL, AND ORNAMENTAL TURF AREAS (CONT.)

SITE	INSECT	OUNCES OF SEVIN brand 5% BATT/1000 SQUARE FEET	SPECIFIC DIRECTIONS
	Leaf- hoppers Lucerne moth larvae, Millipedes, Spittle- bugs, Spring- tails, Ticks, Yellow- striped armyworm		

Self-Verified
Label

FOUNDATION "INSECT BARRIER" TREATMENT
FLOWER BEDS, ORNAMENTAL PLANTINGS

Apply 1 pound per 1,000 Sq. Ft. (32'x32') to all soil areas for use as a contact and bait control of Earwigs, Cutworms, Armyworms, Crickets, Sowbugs, Millipedes, Ants; and for contact control of Fleas; around the outside of buildings, under porches and foundation crawl spaces, in and around flower and ornamental plantings, along soil cracks next to buildings and as an insect barrier in a band 5 feet wide surrounding the foundation. For the insect "Barrier" sprinkle 1/2 lb. of granules in a 5 foot wide band 100 feet in length (500 Sq. Ft.).

IMPORTED FIRE ANTS

For control of Imported Fire ants, apply two to four ounces to each mound. Immediately irrigate with a minimum of one gallon of water to flush the chemical from the granule to the inside of the mound. Do not use over two gallons of water, as flooding of the mound will result in poor control. Apply water gently to avoid disturbing ants and causing migration from the mound. For best results, apply in the early morning or late evening hours when ants are least active. Treat new mounds as they appear.

LIMITED WARRANTY AND DISCLAIMER

MANUFACTURER AND SELLER WARRANT THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER AND SELLER. IN NO CASE SHALL MANUFACTURER AND SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

SEVIN brand Carbaryl Insecticide Bait Granules

Page 9 of 9

March 31, 1994

11 of 11

EXCEPT AS EXPRESSLY PROVIDED HEREIN, MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.

NET WEIGHT
POUNDS

MANUFACTURED BY
HACCO, INC.
BOX 7190
MADISON, WISCONSIN 53707 U.S.A.

Self-Verified
Label