


503551

PM 13 61282-25 A 1211

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460</p>	<p>EPA Reg. Number: 61282-25</p>		<p>Date of Issuance: APR 18 1996</p>
	<p>Term of Issuance: Conditional</p>		
	<p>Name of Pesticide Product: Diazinon Lawn &amp; Garden WBC</p>		
<p>NOTICE OF PESTICIDE: <u>  X  </u> Registration <u>      </u> Reregistration (under FIFRA, as amended)</p>			
<p>Name and Address of Registrant (include ZIP Code):  HACCO, Inc. P.O. Box 7190 Madison, WI 53707</p>			
<p><small>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</small></p>			
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> <li>1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.</li> <li>2. Make the labeling changes listed below before you release the product for shipment:             <ol style="list-style-type: none"> <li>a. Add the phrase "EPA Reg. No. 61282-25".</li> <li>b. In the first group of lawn pests, delete the comma between "Lawn" and "Chinch" in the Pests Column, and in the rate column add a comma to "1000 sq. ft." so that it reads "1,000 sq. ft.", as in all other entries.</li> </ol> </li> </ol>			
<p>Signature of Approving Official:</p>		<p>Date: APR 18 1996</p>	

page 2

EPA Reg. No. 61282-25

- c. In the second group of lawn pests, revise "Ant" to "Ants".
- d. In the third group of lawn pests, bring "Chiggers, Fleas" up one line, i.e., eliminate the blank line between those pest names and "Brown Dog Ticks" to avoid any possible misinterpretation that the 3 gallon rate is for Brown Dog Ticks, while the 24 gallon rate is for Chiggers and Fleas.
- e. For the crop Almonds, correct the spelling of "Parlatoria".
- f. For the crop Turnips and Turnip Tops, add a comma after "Cabbageworms" in the Pest column.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosures

PEEL HERE →

3211

# DIAZINON LAWN & GARDEN WBC

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

APR 18 1996

Water-Based Concentrate.

This product must not be used on golf courses and sod farms.  
For outdoor use on the home lawn and the home garden only.  
Spray concentrate. Contains diazinon insecticide.

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

61282-25

## PROTECTS:

Lawns, Vegetables,  
Fruits and Nuts,  
Roses, Flowers,  
Trees, Shrubs

## CONTROLS:

Ants, Aphids, Chinch Bugs,  
Cutworms, Fire Ants<sup>1</sup>,  
Flea Beetles, Grubs, Mites,  
Sod Webworms and many  
others.  
<sup>1</sup>Aids in control

## Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)  
phosphorothioate ..... 22.4%  
Inert Ingredients: ..... 77.6%  
Total: ..... 100.0%

EPA Reg. No. 61282-

EPA Est. No. 61282-WI-1

**KEEP OUT OF REACH OF CHILDREN.**  
**CAUTION**

See end of inside booklet for additional precautionary statements. See inside booklet for  
complete Directions for Use and Notice of Warranty.

NET CONTENTS: One Pint (473 mL)

HACCO, INC.  
P.O. Box 7190  
Madison, WI 53707

## Diazinon Lawn & Garden WBC

### DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

**IMPORTANT:** Read the entire Directions for Use and Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

#### Conditions of Warranty

The directions for use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Manufacturer or the Seller. All such risks shall be assumed by the buyer. Manufacturer and Seller warrant that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. Manufacturer and Seller make no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Manufacturer or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Manufacturer and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of HACCO, Inc.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

#### General Information

This product is a water-based concentrate product which must be diluted with water before application. To avoid having spray mixture left over, mix only the amount needed. Refer to the section, How to Mix and Apply. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruits and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. This product works fast and applications should be repeated only as directed to maintain control.

Notes: (1) Do not use inside the home. (2) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia, or orchids, since injury may occur. (3) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (4) Do not apply to food crops grown in greenhouses.

**Precautions:** (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

**How to mix and apply:**

Step #1: Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

Step #2: Shake this product well before using. Use clean sprayer. Carefully measure and mix the amount of this product and water as indicated in the rate chart.

Step #3: Spray as directed in rate chart. Thorough coverage is important.

See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

Step #4: Flush sprayer with clean water after each use.

#### Measurement Conversions:

1 tablespoon = 3 teaspoons

1 fl. oz. = 6 teaspoons or 2 tablespoons

6 fl. oz. = 3/4 cup

1 1/2 tablespoons in 3 gals. of water = 1 1/2 teaspoons in 1 gal. of water

2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

#### Lawn Pest Control

##### Precautions

1. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.
3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl.
4. Avoid overlapping of sprays.

5. Where irrigation (watering) is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion, and St. Augustinegrass. To control pests listed below, apply recommended rates.

Pests	RATE		Remarks
	Diazinon Lawn & Garden WBC	Water	
Lawn, Chinch Bugs	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1000 sq. ft.	3 gals. 24 gals.	Thoroughly wet down grass a few hours before applying. In the North, if Chinch Bugs are the only problem, a lower rate of application can be used: 3 teaspoons in 3 gals. of water covers 125 sq. ft. or 4 fl. oz. in 24 gals. of water per 1,000 sq. ft.
Ant, Armyworms, Bermudagrass Mites, Clover Mites, Springtails (Collembola), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs)	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Thoroughly wet down grass a few hours before applying. For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active.  Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect.  For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.
Brown Dog Ticks, Chiggers, Fleas	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Note: Do not apply this product to animals.  For brown dog ticks, spray grass and under shrubbery, particularly near house.
Hyperodes Weevil	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Treat problem areas in mid-April and again in Mid-May. Water grass thoroughly after application.
Millipedes	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Thoroughly wet grass a few hours before applying.
Rhodesgrass Scales	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Apply when crawlers first emerge.
White Grubs of Japanese Beetle, European Chafer, Southern Chafer	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. 24 gals.	Apply anytime between late July and early October. Water grass thoroughly after application.

# Diazinon Lawn & Garden WBC

5711

## Nuisance Pests Outside the Home

<p>Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (<i>Collembola</i>)</p>	<p>Mix 2 fl. oz. in 1 gal. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.</p> <p>To prevent entry into houses, spray a 5-ft. band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 ft. Repeat application as necessary.</p> <p>Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate.</p>
<p>Fire Ants</p>	<p>To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.</p> <p>Note: For best results, apply in cool weather 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear</p>

## Insect control on Ornamental Trees, Shrubs, and Flowers

Spray when listed pests first appear. Repeat as necessary.

Trees and Shrubs	Insects	Rate per 1 Gal. of Water	Timing/Special Directions
<p>Arborvitae, Azaleas, Birch, Boxwood, Camellias, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental, Plum, Poplar, Rhododendron, Scotch Pine, Spirea, Spruce, Sycamore, Wax Plant (Hoya), Willow, Yellowwood, Yew</p>	<p>Aphids, Apple-and-Thorn Skeletonizers, Bagworms, European Pine Sawfly, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak), Hemlock Chermes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Oak Loopers, Oblique-banded Leafrollers, Obscure Root Weevils, Omnivorous Leafhoppers, Pear Slugs, Scale Crawlers (Cottoncuscution, Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft), Tent Caterpillars, Webworms (Cotoneaster, Fall, Juniper, Mimosa), Whiteflies</p>	<p>2 tea-spoons</p>	<p>Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage with spray.</p> <p>Mimosa Webworms: apply when adults first appear. Make 2 additional applications at 4 to 5-week intervals.</p>

Roses and Flowers	Insects	Rate per 1 Gal. of Water	Timing/Special Directions
<p>Azaleas, • Camellias, • Chrysanthemums, Carnations, Gladioli, Lilac, • Lilies, Marigolds, • Pansies, • Petunias, Rhododendron, • Roses</p>	<p>Aphids, Dipterous Leafminers, Euonymus Scale Crawler, • Flea Beetles, • Leafhoppers, • Mealybugs, Mites • Carnation Bud, • Cyclamen, European Red, Two-Spotted Spider, • Thrips, Whiteflies</p>	<p>2 tea-spoons</p>	<p>Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.</p>

## Fruits and Nuts

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted.

Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

Crop	PHI*
Almonds	• •
Apricots	21
Blackberries	7
Blueberries	7
Boysenberries	7
Cherries	21
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums	21
Prunes	21
Raspberries	7
Strawberries	5
Walnuts	45

\* PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

• • Dormant spray only.

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

**Diazinon Lawn & Garden WBC**

6/11

Crops	Pest	Rate	Timing/Special Directions
Almonds (CA only)	San Jose Scale, Pariatoria Scale	2 teaspoons in 1 gal. of water	Apply only as dormant spray. Do not apply more than 6 gals. per 500 sq. ft. of home orchard.  This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.
Apricots	Aphids, Apricot Mealybugs, Mites (Brown, Two- spotted Spider), Scale Crawlers (Olive, San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Berries	Aphids, Blueberry Maggots, Cherry Fruitworms, Cranberry Fruitworms, Thrips, Two-spotted Spider Mites	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Allow a minimum of 14 days between applications. Do not apply more than 4 gals. per 500 sq. ft. of blueberry patch. Do not make more than 2 applications per season.  Allow a minimum of 7 days between the last application and harvest.
Caneberries, (CA, OR, WA only): Blackberries, Boysen- berries, Dewberries, Raspberries	Aphids, Leafhoppers, Raspberry Fruitworms, Two-Spotted Spider Mites, Thrips	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 5 gals. per 500 sq. ft. of caneberry patch. Do not make more than 5 applica- tions per season a minimum of 14 days apart. Do not apply within 7 days of harvest.  This product may be used on caneberries only in CA, OR and WA. Do not use this product on caneberries grown in other states.
Cherries	Black Cherry Aphids, Cherry Fruitflies, Cherry Fruitworms (Maggots), Fruitree Leaf- rollers, Mites (Cherry Rust), Scale Crawlers (San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 21 days of harvest.

Grapes	Aphids, Grape Berry Moths, Grape Leafhoppers, Leafhoppers	1½ teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 3 gals. per 500 sq. ft. of home vineyard. Do not make more than 5 applica- tions per season a minimum of 7 days apart. Do not apply within 28 days of harvest.
Nectarines	Aphids, Mites (Brown, Two- Spotted Spider), Scale Crawlers (Olive, San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Peaches	Aphids, Mites (Brown, Two- Spotted Spider), Scale Crawlers (Olive, San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply within 21 days of harvest.
Plums and Prunes	Aphids (Leafcurl Plum, Mealy Plum, Thistle), Mites (Brown, European Red)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 21 days of harvest.
Strawberries	Aphids, Two-Spotted Spider Mites, Strawberry Leaf- rollers	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 3 gals. per 500 sq. ft. of strawberry patch. Do not make more than 3 applica- tions per season a minimum of 7 days apart. Do not apply within 5 days of harvest.
Walnuts (CA only)	Aphids, Scale Crawlers (Walnut, Italian Pear, Frosted), Spider Mites, Twig Borers	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 6 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applica- tions per season a minimum of 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest.  This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.

# Diazinon Lawn & Garden WBC

7/11

## Vegetables

For foliar or soil applications, this product should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil.

Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

### Pre-Harvest Intervals (PHI\*) for Foliar Applications

Crops	PHI* (Days)
Beans <sup>1</sup> (succulent varieties only: Lima, Pole and Snap)	7
Broccoli	7
Cabbage	21
Cantaloupes	3
Carrots	14
Casabas	3
Chiffonade	7
Chinese Broccoli, Chinese Cabbage, Collards	10
Crenshaws	3
Cucumbers	7
Endive (Escarole)	14
Honeydew Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Onions (Bulb and Green), Parsnips	14
Peas <sup>1</sup>	7
Pumpkins	5
Persian Melons	3
Potatoes (Irish)	35
Radishes, Rutabagas, Spinach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Tomatoes	1
Turnips (Roots and Tops)	14
Watermelons	3

\* PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

<sup>1</sup> Do not use this product on dried pea or bean varieties, such as pinto beans, dried limas, split peas, or dried black-eyed peas.

## Vegetables

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

Crop	Pest	Rate	Timing/Special Directions
Beans, Lima, Pole, and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms (snap beans only)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Notes: (1) Do not apply this product to dried bean varieties, such as pinto beans or dried limas. (2) To protect bees, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom.			
Beets, Red (Table)	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.  Note: Resistant leaf-miners will not be controlled by diazinon or similar type insecticides.
Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 21 days of harvest.  Note: Resistant leaf-miners will not be controlled by diazinon or similar type insecticides.

## Diazinon Lawn &amp; Garden WBC

Cabbage (continued)	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Root Maggots	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Carrots	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Cauliflower	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 appli- cations per season. Do not apply within 7 days of harvest.  Note: Resistant leaf- miners will not be controlled by diazinon or similar type insecticides.
Chinese Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 appli- cations per season. Do not apply within 10 days of harvest.  Note: Resistant leaf- miners will not be controlled by diazinon or similar type insecticides.

Chinese Cabbage (Bok Choy and Napa)	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest.  Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Collards	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest.  Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Corn (see sweet corn)			
Cucumbers	Aphids, Thrips Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden.	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Endive (Escarole)	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden.	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecticides.
Kale	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden.	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest.  Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecticides.
Lettuce	Aphids, Flea Beetles, Leafhoppers Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden.	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.



**Diazinon Lawn & Garden WBC**

Lettuce (Continued)	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Musk- melons, Persians, and hybrids of these) and Water- melons	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Onions (Bulb and Green)	Onion Thrips	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
Parsnips *	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecticides.
Peas (succulent only)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just days before planting. Immediately work into soil 4-8 inches.
Notes: (1) Do not apply this product to dried pea varieties, such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom.			

Peppers	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest.
Potatoes	Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest.  Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecticides.
Radishes	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 3 applications per season. Do not apply within 14 days of harvest.  Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
Rutabagas	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
Spinach *	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecticides.
Squash, Summer	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.

# Diazinon Lawn & Garden WBC

Squash, Winter	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Sweet Corn	Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last application.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Note: To protect bees, do not apply to corn foliage during the pollen shed period.			
Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies, ( <i>Drosophila</i> spp.)	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Turnips and Turnip Tops	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.  Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

Turnips and Turnip Tops (continued)	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
-------------------------------------	----------------------------------	---	---

## STORAGE AND DISPOSAL

**STORAGE:** Store in original container in an area inaccessible to children and pets. If product freezes, place container in a warm room (68°F or above) and allow to thaw. Shake well before using.

**DISPOSAL:** Do not pour the concentrate, the spray mixture, or the rinse water down the drain or sewer. Do not reuse empty container. Triple rinse the empty container, pour the rinse water into the sprayer, then spray the area to be treated. Securely wrap empty container in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

## PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

### CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

### Statement of Practical Treatment

**If swallowed:** Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.  
**If in eyes:** Flush eyes with plenty of water. Call a physician if irritation persists.  
**If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.  
**If on skin:** Wash with plenty of soap and water. Get medical attention.  
**Note to Physician:** This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.

### Environmental Hazards

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1, and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where irrigation (watering) is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least ¼ inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

11/74

# DIAZINON LAWN & GARDEN WBC

Water-Based Concentrate.

This product must not be used on golf courses and sod farms.  
For outdoor use on the home lawn and the home garden only.  
Spray concentrate. Contains diazinon insecticide.

## PROTECTS:

Lawns, Vegetables,  
Fruits and Nuts,  
Roses, Flowers,  
Trees, Shrubs

## CONTROLS:

Ants, Aphids, Chinch Bugs,  
Cutworms, Fire Ants<sup>1</sup>,  
Flea Beetles, Grubs, Mites,  
Sod Webworms and many  
others.

<sup>1</sup>Aids in control

## Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)  
phosphorothioate ..... 22.4%

Inert Ingredients: ..... 77.6%

Total: ..... 100.0%

EPA Reg. No. 61282- *EL*

EPA Est. No. 61282-WI-1

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

See booklet for additional precautionary statements, complete Directions for Use and Notice of  
Warranty. Do not use this product unless booklet has been read.

NET CONTENTS: One Pint (473 mL)

HACCO, INC.  
P.O. Box 7190  
Madison, WI 53707