

## DIAZINON LAWN & GARDEN WBC

Water Based Concentrate

This product must not be used on golf courses and sod farms. For outdoor use on the home lawn and the home garden only. Spray concentrate. Contains diazinon insecticide.

**PROTECTS:** Lawns, Vegetables, Roses, Flowers, Trees, Shrubs, **CONTROLS:** Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants, Flea Beetles, Grubs, Mites, Sod Webworms and many others as listed in Directions For Use. (Aids in Control)

**Active Ingredient:**

**Diazinon:** O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)

phosphorothioate.....22.4%

**Inert Ingredients:**.....77.6%

**Total:**.....100.0%

EPA Reg. No. 61282-25

### KEEP OUT OF REACH OF CHILDREN

### CAUTION

HACCO, INC. PO Box 7190 Madison, WI 53707

### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

### CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home or in greenhouses. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fishponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

### First Aid

This product contains diazinon, an organophosphate insecticide that inhibits cholinesterase

**If swallowed** Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious or convulsing person.

**If in eyes** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

<b>If inhaled</b>	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
<b>If on Skin or Clothing</b>	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
<b>For a Medical Emergency involving this product, call 1-800-228-5635, ext. 136 or call collect 612-851-8180, ext. 136</b>	
<b>NOTE TO PHYSICIAN</b>	
This product contains diazinon, an organophosphate that inhibits cholinesterase. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, but should be administered only in conjunction with atropine. If ingested induce emesis or lavage stomach.	

### ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1, and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4-inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

### DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

**IMPORTANT:** Read the entire Directions for Use and Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

### CONDITIONS OF WARRANTY

**It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of HACCO, Inc., the Manufacturer or Seller. In no case shall HACCO, Inc., the**

**Manufacturer or Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Except as expressly provided herein, HACCO, Inc., the Manufacturer or Seller makes no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. Buyer's or user's exclusive remedy, and HACCO, Inc.'s, the Manufacturer's or Seller's total liability, shall be for damages not exceeding the cost of the product.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Reformulation or repackaging of this product is prohibited.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

### General Information

This product is a water-based concentrate which must be diluted with water before application. To avoid having spray mixture left over, mix only the amount needed. Refer to the section, How to Mix and Apply. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many hard-to-kill sucking and chewing insects. This product works fast and applications should be repeated only as directed to maintain control. **Notes:** (1) Do not use inside the home. (2) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia, or orchids, since injury may occur. (3) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (4) Do not apply this product in greenhouses.

**Precautions:** (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

### How to mix and apply:

**Step #1:** Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

**Step #2:** Shake this product well before using. Use clean sprayer. Carefully measure and mix the amount of this product and water as indicated in the rate chart.

**Step #3:** Spray as directed in rate chart. Thorough coverage is important. See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

**Step #4:** Flush sprayer with clean water after each use.

61282-25

07/24/2002

1/4

ACCEPTED

JUL 24 2002

**Measurement Conversions:**

- 1 tablespoon = 3 teaspoons
- 1 fluid ounce = 6 teaspoons or 2 tablespoons
- 6 fluid ounces = 3/4 cup
- 1-1/2 tablespoons in 3 gallons of water = 1-1/2 baspoons in 1 gallon of water
- 2 tablespoons in 3 gallons of water = 2 teaspoons in 1 gallon of water

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticides registered under

EPA Reg. No. 6/282-25

**Lawn Pest Control**

**Precautions**

1. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.
3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase
4. Avoid overlapping of sprays.
5. Where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product, however, stop watering before puddling occurs.

Use on Bent, Bermuda, Bluegrass, Dichonra, Fescue, Merion, and St. Augustinegrass. To Control pests listed below, apply recommended rates.

Pests	Diazinon Lawn and Garden WBC	Water	Remarks
Lawn Chinch Bugs	1-1/2 tablespoons per 125 sq. ft.	3 gallons	Thoroughly wet down grass a few hours before applying. In the North, if Chinch Bugs are the only problem, a lower rate of application can be used: 3 teaspoons in 3 gallons of water covers 125 sq. ft. or 4 fl. oz. in 24 gallons of water per 1,000 sq. ft.
Ants, Armyworms, Bermudagrass Mites, Clover Mites, Springtails ( <i>Collembola</i> ), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs)	1-1/2 tablespoons per 125 sq. ft.	3 gallons	Thoroughly wet down grass a few hours before applying.  For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active.  Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect.  For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.
Brown Dog Ticks, Chiggers, Fleas	1-1/2 tablespoons per 125 sq. ft.	3 gallons	<b>Note:</b> Do not apply this product to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Hyperodes, Weevil	1-1/2 tablespoons per 125 sq. ft.	3 gallons	Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application.
Millipedes	1-1/2 tablespoons per 125 sq. ft.	3 gallons	Thoroughly wet grass a few hours before applying.
Rhodegrass, Scales	1-1/2 tablespoons per 125 sq. ft.	3 gallons	Apply when crawlers first emerge.
White Grubs of Japanese Beetle, European Chafer, Southern Chafer	1-1/2 tablespoons per 125 sq. ft.	3 gallons	Apply anytime between late July and early October. Water grass thoroughly after application.

**Nuisance Pests Outside the Home**

Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails ( <i>Collembola</i> )	Mix 2 fl. oz. in 1 gallon of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entrance-ways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.  To prevent entry into houses, spray a 5-ft. band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 ft. Repeat applications as necessary.  Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate.
--	--

**Fire Ants**

To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gallon of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gallon per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness. **Note:** For best results, apply in cool weather 65-80 F. or in early morning or late evening hours. Treat new mounds as they appear.

Insect control on Ornamental Trees, Shrubs and Flowers. Spray when listed pests first appear. Repeat as necessary.

Trees and Shrubs	Insects	Rate per 1 gallon of water	Timing / Special Directions
Arborvitae, Azaleas, Birch, Boxwood, Camellias, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental, Plum, Poplar, Rhododendron, Scotch Pine, Spirea, Spruce, Sycamore, Wax Plant (Hoya), Willow, Yellow- wood, Yew	Aphids, Apple and Thorn Skeletonizers, Bagworms, European Pine Sawfly, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak), Hemlock Chermes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Oak Loopers, Oblique-banded Leafrollers, Obscure Root Weevils, Omnivorous Leafhoppers, Pear Slugs, Scale Crawlers (Cottonycusion, Euonymus, Lecanium, Pine Needle, Pitt, San Jose, Soft), Tent Caterpillars, Webworms (Cotoneaster, Fall, Juniper, Mimosa), Whiteflies.	2 teaspoons	Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage with spray.  Mimosa Webworms: apply when adults first appear. Make 2 additional applications at 4 to 5 week intervals.
Roses and Flowers	Insects	Rate per 1 gallon of water	Timing / Special Directions
Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses	Aphids, Dipterous Leafminers, Euonymus Scale Crawler, Flea Beetles, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-Spotted Spider), Thrips, Whiteflies.	2 teaspoons	Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.

**Vegetables**

For foliar or soil applications, this product should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil.

Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply this product in greenhouses.

**Pre-Harvest Intervals (PHI\*) for Foliar Applications**

Crops	PHI* (Days)
Beans <sup>1</sup> (succulent varieties only: Lima, Pole and Snap)	7
Broccoli	7
Cabbage	21
Cantaloupes	3
Carrots	14
Casaba Melons	3
Chinese Broccoli, Chinese Cabbage	10
Crenshaw Melons	3
Cucumbers	7
Honeydew Melons	3
Lettuce	14
Muskmelons	3
Onion (Bulb and Green)	14
Peas <sup>1</sup>	7
Peppers	5
Persian Melons	3
Potatoes (Irish)	35
Squash, Summer	7
Squash, Winter	3
Sweet corn	7
Tomatoes	1
Watermelons	3

\* PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

<sup>1</sup> Do not use this product on dried pea or bean varieties, such as pinto beans, dried limas, split peas, or dried black-eyed peas.

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

Crop	Pest	Rate	Timing / Special Directions
Beans, Lima, Pole, and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms. 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.

## Vegetables (cont'd)

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

Crop	Pest	Rate	Timing / Special Directions
Beans, Lima, Pole, and Snap (succulent varieties only) (cont'd.)	Wireworms (snap beans only)	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
	<b>Notes:</b> (1) Do not apply this product to dried bean varieties, such as pinto beans or dried limas. (2) To protect bees, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom.		
Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. <b>Note:</b> Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 21 days of harvest. <b>Note:</b> Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Root Maggots	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Carrots	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. <b>Note:</b> Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Chinese Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. <b>Note:</b> Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Chinese Cabbage (Bok Choy and Napa)	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 10 days of harvest. <b>Note:</b> Resistant leafminers will not be controlled by diazinon or similar type insecticides.

## Vegetables (cont'd)

For application to the foliage, thoroughly spray all plant parts. Application of the soil should be mixed into the soil immediately after application.

Crop	Pest	Rate	Timing / Special Directions
Peas (succulent only) (cont'd.)	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just days before planting. Immediately work into soil 4-8 inches.
	<b>Notes:</b> (1) Do not apply this product to dried pea varieties, such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if the peas or weeds in the treatment area are in bloom.		
Peppers	Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 5 days of harvest.
Potatoes	Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 35 days of harvest. <b>Note:</b> Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Squash, Summer	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Squash, Winter	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Sweet Corn	Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworm, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
	<b>Note:</b> To protect bees, do not apply to corn foliage during the pollen shed period.		
Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies, ( <i>Drosophila</i> spp.)	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest.

Corn (see Sweet Corn)			
Cucumbers	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest.
Lettuce	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and hybrids of these) and Watermelons	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 3 days of harvest.
Onions (bulb and green)	Onion Thrips	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest.
Peas (succulent only)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	1 teaspoon in 1 gallon of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Allow at least 7 days between the last application and harvest.

Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gallon of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.

### STORAGE AND DISPOSAL

**STORAGE:** Store in original container in an area inaccessible to children and pets. If product freezes, place container in a warm room (68 F or above) and allow to thaw. Shake well before using.

**DISPOSAL:** Do not pour the concentrate, the spray mixture, or the rinse water down the drain or sewer. Do not reuse empty container. Triple rinse the empty container, pour the rinse water into the sprayer, then spray the area to be treated. Securely wrap empty container in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

*THIS SPACE  
INTENTIONALLY  
LEFT BLANK*

### DIAZINON LAWN & GARDEN WBC

Water Based Concentrate

This product must not be used on golf courses and sod farms.  
For outdoor use on the home lawn and the home garden only.  
Spray concentrate. Contains diazinon insecticide.

**Active Ingredient:**

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate .....22.4%

Inert Ingredients: .....77.6%

Total: .....100.0%

EPA Reg. No. 61282-25  
KEEP OUT OF REACH OF CHILDREN  
CAUTION

See inside booklet for additional precautionary statements,  
directions for use and notice of warranty.  
HACCO, INC. PO BOX 7190 MADISON, WI 53707

5/10

Booklet

Container Label — 6/6

## DIAZINON LAWN & GARDEN WBC

Water Based Concentrate

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

**PROTECTS:** Lawns, Vegetables, Roses, Flowers, Trees, Shrubs  
**CONTROLS:** Ants, Aphids, Chinch Bugs, Cutworms, Fire Ants,\* Flea Beetles, Grubs, Mites, Sod webworms and many others as listed in Directions For Use. \*Aids in control

Active Ingredient:

Diazinon: 0,0-Diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate.....22.4%

Inert Ingredients.....77.6%

Total.....100.0%

EPA Reg # 61282-25

EPA Est # 61282-WI-1 (Lot begins 51) 34704-CO-1 (Lot begins 06)

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See booklet for additional precautionary statements, complete Directions for Use and Notice of Warranty. Do not use this product unless booklet has been read.  
HACCO, INC. • PO BOX 7190 • MADISON, WI 53707

**NET CONTENTS: 16 fl. oz. (473 mL)**

